

A Course in Interacting Particle Systems

J.M. Swart

January 7, 2022

Contents

1	Introduction	9
1.1	General set-up	9
1.2	The voter model	11
1.3	The contact process	13
1.4	Ising and Potts models	16
1.5	Phase transitions	20
1.6	Variations on the voter model	23
1.7	Branching and coalescing particles	25
1.8	Periodic behavior	29
2	Continuous-time Markov chains	35
2.1	Poisson point sets	35
2.2	Transition probabilities and generators	38
2.3	Poisson construction of Markov processes	40
2.4	Examples of Poisson representations	43
2.5	Countable state space	44
3	The mean-field limit	47
3.1	Processes on the complete graph	47
3.2	The mean-field limit of the Ising model	48
3.3	Analysis of the mean-field model	50
3.4	Functions of Markov processes	54
3.5	The mean-field contact process	59
3.6	The mean-field voter model	61
3.7	Exercises	63
4	Construction and ergodicity	65
4.1	Introduction	65
4.2	Feller processes	67
4.3	Poisson construction	77
4.4	Generator construction	90
4.5	Ergodicity	96
4.6	Application to the Ising model	99
4.7	Further results	103

5	Monotonicity	107
5.1	The stochastic order	107
5.2	The upper and lower invariant laws	112
5.3	The contact process	115
5.4	Other examples	118
5.5	Exercises	119
6	Duality	123
6.1	Introduction	123
6.2	Additive systems duality	124
6.3	Cancellative systems duality	132
6.4	The backward in time process	137
6.5	Markov process duality	139
6.6	Additive duality revisited	140
6.7	Cancellative duality revisited	145
6.8	Other dualities	147
6.9	Invariant laws of the voter model	154
6.10	Homogeneous invariant laws	156
6.11	Equality of critical points	162
7	Oriented percolation	165
7.1	Introduction	165
7.2	Oriented percolation	166
7.3	Survival	169
7.4	K-dependence	171

Preface

Interacting particle systems, in the sense we will be using the word in these lecture notes, are countable systems of locally interacting Markov processes. Each interacting particle system is defined on a lattice: a countable set with (usually) some concept of distance defined on it; the canonical choice is the d -dimensional integer lattice \mathbb{Z}^d . Situated on each point in this lattice, there is a continuous-time Markov process with a finite state space (often even of cardinality two) whose jump rates depend on the states of the Markov processes on near-by sites. Interacting particle systems are often used as extremely simplified ‘toy models’ for stochastic phenomena that involve a spatial structure.

An attractive property of interacting particle systems is that they are easy to simulate on a computer.¹ Although the definition of an interacting particle system often looks very simple, and problems of existence and uniqueness have long been settled, it is often surprisingly difficult to prove anything nontrivial about its behavior. With a few exceptions, explicit calculations tend not to be feasible, so one has to be satisfied with qualitative statements and some explicit bounds. Despite intensive research for over more than forty years, some easy-to-formulate problems still remain open while the solutions of others have required the development of nontrivial and complicated techniques.

Luckily, as a reward for all this, it turns out that despite their simple rules, interacting particle systems are often remarkably subtle models that capture the sort of phenomena one is interested in much better than might initially be expected. Thus, while it may seem outrageous to assume that “Plants of a certain type occupy points in the square lattice \mathbb{Z}^2 , live for an exponential time with mean one, and place seeds on unoccupied neighboring sites with rate λ ” it turns out that making the model more realistic often does not change much in its overall behavior. Indeed, there is a general philosophy in the field, that is still insufficiently understood, that says that interacting particle systems come in ‘universality classes’ with the property that all models in one class have roughly the same behavior.

As a mathematical discipline, the subject of interacting particle systems is still relatively young. It started around 1970 with the work of R.L. Dobrushin and F. Spitzer, with many other authors joining in during the next few years. By 1975, general existence and uniqueness questions had been settled, four classic models had been introduced (the exclusion process, the stochastic

¹To get started doing this yourself, look at my simulation library that is available from <http://staff.utia.cas.cz/swart/simulate.html>.

Ising model, the voter model and the contact process), and elementary (and less elementary) properties of these models had been proved. In 1985, when Liggett's published his famous book [Lig85], the subject had established itself as a mature field of study. Since then, it has continued to grow rapidly, to the point where it is impossible to accurately capture the state of the art in a single book. Indeed, it would be possible to write a book on each of the four classic models mentioned above, while many new models have been introduced and studied.

While interacting particle systems, in the narrow sense indicated above, have apparently not been the subject of mathematical study before 1970, the subject has close links to some problems that are considerably older. In particular, the Ising model (without time evolution) has been studied since 1925 while both the Ising model and the contact process have close connections to percolation, which has been studied since the late 1950-ies. In recent years, more links between interacting particle systems and other, older subjects of mathematical research have been established, and the field continues to receive new impulses not only from the applied, but also from the more theoretical side.

The present notes are loosely based on an older set of lecture notes for courses that I gave at Charles University in Prague in 2009 and 2011. Another input came from slides for a course I gave at Verona University in 2014. Compared to the lecture notes of 2011, most of the text has been rewritten. Many figures have been added, as well as a chapter on the mean-field limit. The old lecture notes were organized around three classical models: the contact process, the Ising model, and the voter model. Instead, the present notes are organized around methods: the mean-field limit, graphical representations, monotone coupling, duality, and comparison with oriented percolation. Compared to the older notes, some results have been removed, in particular about the Ising model, whose study requires rather different techniques from the other models. Another omission are positive correlations. On the other hand, a wide range of interacting particle systems not (or barely) mentioned in the previous lecture notes are now used as examples throughout the notes, to give a better impression of the modern literature of the subject.

I am indebted to Tibor Mach for a careful reading of the lecture notes from 2011 that led to a large number of typos being corrected. I would like to thank Aernout van Enter for a number of corrections and suggestions that helped me improve the text from 2016. Sam Olesker-Taylor and Jan Niklas Latz pointed out some typos in the versions of 2020 and 2021.

These lecture notes have first been posted on the arXiv in 2017 and have been updated in 2020 and 2022. The update in 2020 corrected the proof of

Lemma 4.31, which was wrong in the original version, and made some other minor corrections. In the update of 2022 several additions have been made. The main existence and uniqueness result in Chapter 4 has been formulated as a pathwise uniqueness result for solutions of the evolution equation (4.14). Chapter 6 has also been radically rewritten and in Chapter 1 a section has been added about periodic behavior. In the other chapters several smaller points have been clarified and some pictures have been added.

Chapter 1

Introduction

1.1 General set-up

Let S be a finite set, called the *local state space*, and let Λ be a countable set, called the *lattice*. We let S^Λ denote the Cartesian product space of Λ copies of S , i.e., elements x of S^Λ are of the form

$$x = (x(i))_{i \in \Lambda} \quad \text{with} \quad x(i) \in S \quad \forall i \in \Lambda.$$

Equivalently, S^Λ is nothing else than the set of all functions $x : \Lambda \rightarrow S$.

Interacting particle systems are continuous-time Markov processes $X = (X_t)_{t \geq 0}$ with a state space of the form S^Λ . Thus, $(X_t)_{t \geq 0}$ is a Markov process such that at each time $t \geq 0$, the state of X is of the form

$$X_t = (X_t(i))_{i \in \Lambda} \quad \text{with} \quad X_t(i) \in S \quad \forall i \in \Lambda.$$

We call $X_t(i)$ the *local state* of X at time t and at the *position* i . Positions $i \in \Lambda$ are also often called *sites*.

The time evolution of continuous-time Markov processes is usually characterized by their *generator* G , which is an operator acting on functions $f : \mathcal{S} \rightarrow \mathbb{R}$, where \mathcal{S} is the state space. For example, in the case of Brownian motion, the state space is \mathbb{R} and the generator is the differential operator $G = \frac{1}{2} \frac{\partial^2}{\partial x^2}$. In the case of an interacting particle system, the state space is of the form $\mathcal{S} = S^\Lambda$ and the generator can usually be written in the form

$$Gf(x) = \sum_{m \in \mathcal{G}} r_m \{f(m(x)) - f(x)\} \quad (x \in S^\Lambda). \quad (1.1)$$

Here \mathcal{G} is a set whose elements are *local maps* $m : S^\Lambda \rightarrow S^\Lambda$ and $(r_m)_{m \in \mathcal{G}}$ is a collection of nonnegative constants called *rates*, that say with which

Poisson intensity the local map m should be applied to the configuration X_t . The precise definitions will be given in later chapters, but at the moment it suffices to say that if we approximate $(X_t)_{t \geq 0}$ by a discrete-time Markov chain where time is increased in steps of size dt , then

$r_m dt$ is the probability that the map m
is applied during the time interval $(t, t + dt]$.

Often, the lattice Λ has the structure of an (undirected) graph. In this case, we let E denote the corresponding *edge set*, i.e., a set of unordered pairs $\{i, j\}$ called *edges*, with $i, j \in \Lambda$, $i \neq j$, that in drawings of the graph are connected by a line segment. We let

$$\mathcal{E} := \{(i, j) : \{i, j\} \in E\}$$

denote the corresponding set of all *ordered* pairs (i, j) that correspond to an edge. We call

$$\mathcal{N}_i := \{j \in \Lambda : \{i, j\} \in E\} \tag{1.2}$$

the *neighborhood* of the site i .

Many well-known and well-studied interacting particle systems are defined on the d -dimensional integer lattice \mathbb{Z}^d . We denote the origin by $0 = (0, \dots, 0) \in \mathbb{Z}^d$. For any $i = (i_1, \dots, i_d) \in \mathbb{Z}^d$, we let

$$\|i\|_1 := \sum_{k=1}^d |i_k| \quad \text{and} \quad \|i\|_\infty := \max_{k=1, \dots, d} |i_k| \quad (i \in \mathbb{Z}^d)$$

denote the ℓ_1 -norm and supremumnorm, respectively. For $R \geq 1$, we set

$$E^d := \{\{i, j\} : \|i - j\|_1 = 1\} \quad \text{and} \quad E_R^d := \{\{i, j\} : 0 < \|i - j\|_\infty \leq R\}. \tag{1.3}$$

Then (\mathbb{Z}^d, E^d) is the integer lattice equipped with the *nearest neighbor* graph structure and (\mathbb{Z}^d, E_R^d) is the graph obtained by connecting all edges within $\|\cdot\|_\infty$ -distance R with an edge. We let \mathcal{E}^d and \mathcal{E}_R^d denote the corresponding sets of ordered pairs (i, j) .

Before we turn to rigorous mathematical theory, it is good to see a number of examples. It is easy to simulate interacting particle systems on a computer. In simulations, the infinite graphs (\mathbb{Z}^d, E^d) or (\mathbb{Z}^d, E_R^d) are replaced by a finite piece of \mathbb{Z}^d , with some choice of the boundary conditions (e.g. periodic boundary conditions).

1.2 The voter model

For each $i, j \in \Lambda$, the *voter model map* $\text{vot}_{ij} : S^\Lambda \rightarrow S^\Lambda$ is defined as

$$\text{vot}_{ij}(x)(k) := \begin{cases} x(i) & \text{if } k = j, \\ x(k) & \text{otherwise.} \end{cases} \quad (1.4)$$

Applying vot_{ij} to a configuration x has the effect that local state of the site i is copied onto the site j . The *nearest neighbor voter model* is the interacting particle system with generator

$$G_{\text{vot}}f(x) = \frac{1}{|\mathcal{N}_0|} \sum_{(i,j) \in \mathcal{E}^d} \{f(\text{vot}_{ij}(x)) - f(x)\} \quad (x \in S^\Lambda). \quad (1.5)$$

Here \mathcal{N}_0 is the neighborhood of the origin and $|\mathcal{N}_0| = 2d$ denotes its cardinality. Similarly, replacing the set of oriented edges \mathcal{E}^d by \mathcal{E}_R^d and replacing \mathcal{N}_0 by the appropriate set of neighbors in this new graph, we obtain the *range R voter model*.

In the context of the voter model, the local state $x(i)$ at a site i is often called the *type* at i . The voter model is often used to model biological populations, where organisms with different genetic types occupy sites in space. Note that since each site j has $|\mathcal{N}_j| = |\mathcal{N}_0|$ neighbors, the total rate of all maps vot_{ij} with $i \in \mathcal{N}_j$ is one. In view of this, an alternative way to describe the dynamics in (1.5) is to say that with rate 1, the organism living at a given site dies, and is replaced by a descendant chosen with equal probability from its neighbors.

An alternative interpretation, that has given the voter model its name, is that sites represent people and types represent political opinions. With rate one, an individual becomes unsure what political party to vote for, asks a randomly chosen neighbor, and copies his/her opinion.

In Figure 1.1, we see the four snapshots of the time evolution of a two-dimensional nearest-neighbor voter model. The initial state is constructed by assigning i.i.d. types to the sites. Due to the copying dynamics, we see patches appear where every site in a local neighborhood has the same type. As time proceeds, these patches, usually called *clusters*, grow in size, so that eventually, for any $N \geq 1$, the probability that all sites within distance N of the origin are of the same type tends to one.¹

It turns out that this sort of behavior, called *clustering*, is dimension dependent. The voter model clusters in dimensions 1 and 2, but not in

¹In spite of this, for the model on the infinite lattice, it is still true that the origin changes its type infinitely often.




Figure 1.1: Four snapshots of a two-dimensional voter model with periodic boundary conditions. Initially, the types of sites are i.i.d. Time evolved in these pictures is 0, 1, 32, and 500.

dimensions 3 and more. In Figure 1.2, we see the four snapshots of the time evolution of a three-dimensional voter model. The model is simulated on a cube with periodic boundary conditions, and the types of the middle layer are shown in the pictures. In this case, we see that even after a long time, there are still many different types near the origin.²

²On a finite lattice, such as we use in our simulations, one would eventually see one type take over, but the time one has to wait for this is very long compared to dimensions 1 and 2. On the infinite lattice, the probability that the origin has a different type from its right neighbor tends to a positive limit as time tends to infinity.




Figure 1.2: Four snapshots of a three-dimensional voter model with periodic boundary conditions. Initially, the types of sites are i.i.d. Time evolved in these pictures is 0, 4, 32, and 250.

1.3 The contact process

The contact process is another interacting particle system with a biological interpretation. For this process, we choose the local state space $S = \{0, 1\}$. We interpret a site such that $X_t(i) = 1$ as *occupied* by an organism, and a site such that $X_t(i) = 0$ as *empty*. Alternatively, the contact process can be seen as a model for the spread of an infection. In this case, sites with $X_t(i) = 1$ are called *infected* and sites with $X_t(i) = 0$ are called *healthy*.

For each $i, j \in \Lambda$, we define a *branching map* $\mathbf{bra}_{ij} : \{0, 1\}^\Lambda \rightarrow \{0, 1\}^\Lambda$ as

$$\mathbf{bra}_{ij}(x)(k) := \begin{cases} x(i) \vee x(j) & \text{if } k = j, \\ x(k) & \text{otherwise.} \end{cases} \quad (1.6)$$

Note that this says that if prior to the application of \mathbf{bra}_{ij} , the site i is occupied, then after the application of \mathbf{bra}_{ij} , the site j will also be occupied, regardless of its previous state. If initially i is empty, then nothing happens. We interpret this as the organism at i giving *birth* to a new organism at j , or the infected site i *infecting* the site j . If j is already occupied/infected, then nothing happens.

For each $i \in \Lambda$, we also define a *death map* $\mathbf{death}_i : \{0, 1\}^\Lambda \rightarrow \{0, 1\}^\Lambda$ as

$$\mathbf{death}_i(x)(k) := \begin{cases} 0 & \text{if } k = i, \\ x(k) & \text{otherwise.} \end{cases} \quad (1.7)$$

If the map \mathbf{death}_i is applied, then an organism at i , if there is any, dies, respectively, the site i , if it is infected, *recovers* from the infection.



Figure 1.3: Four snapshots of a two-dimensional contact process. Initially, only a single site is infected. The infection rate is 2, the death rate is 1, and time evolved in these pictures is 1, 5, 10, and 20.

Recalling (1.3), the (nearest neighbor) contact process with *infection rate*

$\lambda \geq 0$ and *death rate* $\delta \geq 0$ is the interacting particle system with generator

$$G_{\text{cont}}f(x) := \lambda \sum_{(i,j) \in \mathcal{E}^d} \{f(\text{bra}_{ij}(x)) - f(x)\} + \delta \sum_{i \in \mathbb{Z}^d} \{f(\text{death}_i(x)) - f(x)\} \quad (x \in \{0,1\}^{\mathbb{Z}^d}). \quad (1.8)$$

This says that infected sites infect each healthy neighbor with rate λ , and infected sites recover with rate δ .

In Figure 1.3, we see the four snapshots of the time evolution of a two-dimensional contact process. Occupied sites are black and empty sites are white. Initially, only the origin is occupied. The infection rate is 2 and the death rate is 1. In this example, the infection spreads through the whole population, eventually reaching a steady state³ where a positive fraction of the population is infected. Of course, starting from a single infected site, there is always a positive probability that the infection dies out in the initial stages of the epidemic.

Unlike the voter model, the behavior of the contact process is roughly similar in different dimensions. On the other hand, the proportion λ/δ of the infection rate to the death rate is important for the behavior. By changing the speed of time, we can without loss of generality choose one of the constants λ and δ to be one, and it is customary to set $\delta := 1$. In Figure 1.4, we have plotted the *survival probability*

$$\theta(\lambda) := \mathbb{P}^{1_{\{0\}}} [X_t \neq 0 \ \forall t \geq 0] \quad (1.9)$$

of the one-dimensional contact process, started in $X_0 = 1_{\{0\}}$, i.e., with a single infected site at the origin, as a function of the infection rate λ . For reasons that we cannot explain here, this is in fact the same as the probability that the origin is infected in equilibrium.

It turns out that for the nearest-neighbor contact process on \mathbb{Z}^d , there exists a *critical value* $\lambda_c = \lambda_c(d)$ with $0 < \lambda_c < \infty$ such that $\theta(\lambda) = 0$ for $\lambda \leq \lambda_c$ and $\theta(\lambda) > 0$ for $\lambda > \lambda_c$. The function θ is continuous, strictly increasing and concave on $[\lambda_c, \infty)$ and satisfies $\lim_{\lambda \rightarrow \infty} \theta(\lambda) = 1$. One has

$$\lambda_c(1) = 1.6489 \pm 0.0002. \quad (1.10)$$

Proving these statements is not easy, however. For example, continuity of the function θ in the point λ_c was proved only in 1990 [BG90], seventeen years

³In fact, on the finite square used in our simulations, one can prove that the infection dies out a.s. However, the time one has to wait for this is exponentially large in the system size. For the size of system shown in Figure 1.3, this time is already too long to be numerically observable.




Figure 1.4: Survival probability of the one-dimensional contact process.

after the introduction of the model in [CS73, Har74]. The best⁴ rigorous upper bound on the constant from (1.10) is $\lambda_c(1) \leq 1.942$ which is proved in [Lig95].

Krone [Kro99] introduced a *two-stage contact process*. In this model, the local state space is $\{0, 1, 2\}$ where 0 represents an empty site, 1 a young organism, and 2 an adult organism. In a branching event, an adult organism produces a young organism on an empty neighboring site. In addition, young organisms can grow up. Both young and adults can die, the young possibly at a higher rate than the adults. The behavior of this model is similar to that of the contact process.

1.4 Ising and Potts models

In a *stochastic Ising model*, sites in the lattice \mathbb{Z}^d are interpreted as atoms in a crystal, that can have two possible local states, usually denoted by -1 and $+1$. In the traditional interpretation, these states describe the direction of the magnetic field of the atom, and because of this, the local state $x(i)$ of a site i is usually called the *spin* at i . More generally, one can consider *stochastic Potts models* where each “spin” can have $q \geq 2$ possible values. In this case, the local state space is traditionally denoted as $S = \{1, \dots, q\}$,

⁴There exists a sequence of rigorous upper bounds on the constant from (1.10) that is known to converge to the real value, but these bounds are so difficult to calculate that the best bound that has really been achieved by this method is much worse than the one in [Lig95].

the special case $q = 2$ corresponding to the Ising model (except for a small difference in notation between $S = \{-1, +1\}$ and $S = \{1, 2\}$).

Given a state x and site i , we let

$$N_{x,i}(\sigma) := \sum_{j \in \mathcal{N}_i} 1_{\{x(j) = \sigma\}} \quad (\sigma \in S) \quad (1.11)$$

denote the number of neighbors of the site i that have the spin value $\sigma \in S$. In the Ising and Potts models, sites like or dislike to have the same spin value as their neighbors, depending on a parameter $\beta \in \mathbb{R}$. Adding a so-called *Glauber dynamics* to the model,⁵ sites update their spin values with rate one, and at such an event choose a new spin value with probabilities that depend on the values of their neighbors. More precisely, the *stochastic Potts model with Glauber dynamics* is the interacting particle system that evolves in such a way that

$$\text{site } i \text{ flips to the value } \sigma \text{ with rate } r_i^\sigma(x) := \frac{e^{\beta N_{x,i}(\sigma)}}{\sum_{\tau \in S} e^{\beta N_{x,i}(\tau)}}. \quad (1.12)$$

More formally, we can write the generator as

$$G_{\text{Potts}} f(x) := \sum_{i \in \mathbb{Z}^d} \sum_{\sigma \in S} r_i^\sigma(x) \{f(m_i^\sigma(x)) - f(x)\}, \quad (1.13)$$

where $m_i^\sigma : S^\Lambda \rightarrow S^\Lambda$ are maps defined by

$$m_i^\sigma(x)(j) := \begin{cases} \sigma & \text{if } j = i, \\ x(j) & \text{otherwise.} \end{cases} \quad (1.14)$$

The attentive reader may notice that the way we have written the generator in (1.13) is different from the way we have written our generators so far, since unlike the rates r_m in (1.1), the rates $r_i^\sigma(x)$ depend on the state x . This will be explained in more detail in Chapter 4. In particular, in Section 4.6, we will see that it is possible to rewrite the generator in (1.13) in a way that fits the general form (1.1) (with rates that do not depend on the state x) but for the Potts model, unlike the models we have seen so far, this way of writing the generator is less natural and more complicated.

⁵The terms *Ising model* and *Potts model* refer only to certain Gibbs measures. A *stochastic Ising model* or *Potts model* is any interacting particle system that has these Gibbs measures as its invariant laws (usually reversible). There exist several different ways to invent a dynamics with this property. This will be explained in a bit more detail in Section 5.4. In this section, we stick to Glauber dynamics.

Returning to our informal description in (1.12), we notice that for $\beta > 0$, sites prefer to have spin values that agree with as many neighbors as possible, i.e., the model is *ferromagnetic*. For $\beta < 0$, the model is *antiferromagnetic*. These terms reflect the situation that in some materials, neighboring spins like to line up, which can lead to long-range order that has the effect that the material can be magnetized. Antiferromagnetic materials, on the other hand, lack this effect.

Alternatively, Potts models can also be interpreted as social or economic models, where sites represent people or firms and spin values represent opinions or the state (financially healthy or not) of a firm [BD01].




Figure 1.5: Four snapshots of a $q = 4$, $\beta = 1.2$ Potts model with Glauber dynamics and periodic boundary conditions. Initially, the types of sites are i.i.d. Time evolved in these pictures is 0, 4, 32, 500.

In Figure 1.5 we see four snapshots of a two-dimensional nearest-neighbor Potts model with four possible spin values. We have used periodic boundary conditions, and the value of the parameter β is 1.2. Superficially, the behavior is similar to that of a voter model, in the sense that the system forms clusters

of growing size that in the end take over any finite neighborhood of the origin. Contrary to the voter model, however, even in the middle of a large cluster that is predominantly of one color, sites can still flip to other values as is clear from (1.12), so in the simulations we see many small islands of different colors inside large clusters where one color dominates. Another difference is that clustering happens only when the value of the parameter β is large enough. For small values of β , the behavior is roughly similar to the voter model in dimensions $d \geq 3$. There is a critical value $0 < \beta_c < \infty$ where the model changes from one type of behavior to the other type of behavior. In this respect, the model is similar to the contact process.

To make this critical value visible, imagine that instead of periodic boundary conditions, we would use frozen boundary conditions where the sites at the boundary are kept fixed at one chosen color, say color 1. Then the system has a unique invariant law (equilibrium), in which for sufficiently large values of β the color 1 is (much) more frequent than the other colors, but for low values of β all colors occur with the same frequency. In particular, for the Ising model, where the set of possible spin values in $\{-1, +1\}$, we let

$$m_*(\beta) := \begin{array}{l} \text{the expectation of } x(0) \text{ with } +1 \text{ boundary} \\ \text{conditions, in the limit of large system size.} \end{array} \quad (1.15)$$

This function is called the *spontaneous magnetization*. For the Ising model in two dimensions, the spontaneous magnetization can be explicitly calculated, as was first done by Onsager [Ons44]. The formula is

$$m_*(\beta) = \begin{cases} (1 - \sinh(\beta)^{-4})^{1/8} & \text{for } \beta \geq \beta_c := \log(1 + \sqrt{2}), \\ 0 & \text{for } \beta \leq \beta_c. \end{cases} \quad (1.16)$$

This function is plotted in Figure 1.6. In this case, the critical point β_c is known explicitly.

For Ising models in dimensions $d \geq 3$, the graph of $m_*(\beta)$ looks roughly similar to Figure 1.6, with $\beta_c \approx 0.442$ in dimension 3 [GPA01], but no explicit formulas are known.

In dimension one, one has $m^*(\beta) = 0$ for *all* $\beta \geq 0$. More generally, one-dimensional Potts models do not show long range order, even if β is very large.⁶ By this we mean that in equilibrium, the correlation between the spin values at 0 and a point $i \in \mathbb{Z}$ tends to zero as $i \rightarrow \infty$ for any value of β (even

⁶This was first noticed by Ising [Isi25], who introduced the model but noticed that it was uninteresting, incorrectly assuming that what he had proved in dimension 1 would probably hold in any dimension. Peierls [Pei36] realized that dimension matters and proved that the Ising model in higher dimensions does show long range order.




Figure 1.6: The spontaneous magnetization of the two-dimensional Ising model.

though the decay is slow if β is large). In Figure 1.7, we compare the time evolution of a one-dimensional Potts model (with a large value of β) with the time evolution of a one-dimensional voter model. In the voter model, the cluster size keeps growing, but in the Potts model, the typical cluster size converges to a finite limit.

1.5 Phase transitions

Figures 1.4 and 1.6 are examples of a phenomenon that is often observed in interacting particle systems. As a parameter governing the dynamics crosses a particular value, the system goes through an abrupt change in behavior. This is called a *phase transition* and the value of the parameter is called the *point of the phase transition* or, in the mathematical literature, *critical point*. As we will see in a moment, in the physics literature, the term critical point has a more restricted meaning. The term “phase transition” of course also describes the behavior that certain materials change from a gas, fluid, or solid phase into another phase at a particular value of the temperature, pressure etc., and from the theoretical physicist’s point of view, this is indeed the same phenomenon.

In both Figure 1.4 and 1.6, the point of the phase transition in fact separates two regimes, one where the interacting particle systems (on the infinite lattice) has a unique invariant law (below λ_c and β_c) and another regime where there are more invariant laws (above λ_c and β_c). Indeed, for the contact process, the delta measure on the empty configuration is always an invariant law, but above λ_c , a second, nontrivial invariant law also appears. Potts models have q invariant laws (one corresponding to each color)




Figure 1.7: Time evolution of a one-dimensional voter model (above) and a one-dimensional Potts model (below) with a high value of β .

above the critical point.⁷ Multiple invariant laws are a general phenomenon associated with phase transitions.

Phase transitions are classified into *first order* and *second order* phase transitions.⁸ Second order phase transitions are also called *continuous* phase

⁷More precisely, they have q invariant laws that have the additional property that they are also translation invariant in space. Depending on the dimension, there may exist additional invariant laws that are not translation invariant.

⁸This terminology was introduced by Paul Ehrenfest. The idea is that in first order phase transitions, the first derivative of the free energy has a discontinuity, while in a second order phase transitions, the first derivative of the free energy is continuous and only the second derivative makes a jump.

transitions. The phase transitions in Figures 1.4 and 1.6 are both second order, since the functions θ and m_* are continuous at the critical points λ_c and β_c , respectively. Also, second order phase transitions are characterized by the fact that at the critical point, there is only one invariant law. By contrast, if we would draw the function $m_*(\beta)$ of a Potts model for sufficiently large values of q (in dimension two, for $q > 4$), then the plot of m_* would make a jump at β_c and the system would have multiple invariant laws at this point, which means that this phase transition is first order.

It can be difficult to prove whether a given phase transition is first or second order. While for the two-dimensional Ising model, continuity of the magnetization follows from Onsager's solution [Ons44], the analogous statement for the three-dimensional Ising model was only proved recently [ADS15] (70 years after Onsager!).

For the Ising model, it is known (but only partially proved) that

$$m_*(\beta) \propto (\beta - \beta_c)^c \quad \text{as } \beta \downarrow \beta_c,$$

where c is a *critical exponent*, which is given by

$$c = 1/8 \text{ in dim 2, } \quad c \approx 0.326 \text{ in dim 3, } \quad \text{and } \quad c = 1/2 \text{ in dim } \geq 4.$$

For the contact process, it has numerically been observed that

$$\theta(\lambda) \propto (\lambda - \lambda_c)^c \quad \text{as } \lambda \downarrow \lambda_c,$$

with a critical exponent

$$\begin{aligned} c &\approx 0.276 \text{ in dim 1, } \quad c \approx 0.583 \text{ in dim 2,} \\ c &\approx 0.813 \text{ in dim 3, } \quad \text{and } \quad c = 1 \text{ in dim } \geq 4. \end{aligned}$$

In theoretical physics, (nonrigorous) *renormalization group theory* is used to explain these critical exponents and calculate them. According to this theory, critical exponents are *universal*. For example, the nearest-neighbor model and the range R models with different values of R all have different values of the critical point, but the critical exponent c has the same value for all these models.⁹ Also, changing from the square lattice to, e.g., the triangular lattice has no effect on c .

Critical exponents are associated only with second order phase transitions. At the critical point of a second order phase transition, one observes

⁹Universality in the range R does not always hold. It has been proved that the $q = 3$ ferromagnetic Potts model in dimension two has a first order phase transition for large R [GB07], while the model with $R = 1$ is known to have a second order phase transition [DST17].

critical behavior, which involves, e.g., power-law decay of correlations. For this reason, physicists use the term “critical point” only for second order phase transitions.

So far, there is no mathematical theory that can explain critical behavior, except in high dimensions (where one uses a technique called the *lace expansion*) and in a few two-dimensional models (that have a conformally invariant scaling limit that can be described using the Schramm-Loewner equation).

1.6 Variations on the voter model

Apart from the models discussed so far, lots of other interacting particle systems have been introduced and studied in the literature to model a plethora of phenomena. Some of these behave very similarly to the models we have already seen (and even appear to have the same critical exponents), while others are completely different. In this and the next sections, we take a brief look at some of these models to get an impression of the possibilities.

The *biased voter model* with *bias* $s \geq 0$ is the interacting particle system with state space $\{0, 1\}^{\mathbb{Z}^d}$ and generator (compare (1.5))

$$\begin{aligned} G_{\text{bias}}f(x) := & \frac{1}{2d} \sum_{(i,j) \in \mathcal{E}^d} \{f(\text{vot}_{ij}(x)) - f(x)\} \\ & + \frac{s}{2d} \sum_{(i,j) \in \mathcal{E}^d} \{f(\text{bra}_{ij}(x)) - f(x)\}, \end{aligned} \quad (1.17)$$

where vot_{ij} and bra_{ij} are the voter and branching maps defined in (1.4) and (1.6). The biased voter model describes a situation where one genetic type of an organism (in this case, type 1) is more fit than the other type, and hence reproduces at a larger rate. Alternatively, this type may represent a new idea or opinion that is more attractive than the current opinion. Contrary to the normal voter model, even if we start with just a single individual of type 1, there is a positive probability that type 1 never dies out and indeed takes over the whole population, as can be seen in Figure 1.8.

Fix $i \in \mathbb{Z}^d$ and for any $x \in \{0, 1\}^{\mathbb{Z}^d}$, let

$$f_\tau(x) := \frac{1}{|\mathcal{N}_i|} \sum_{j \in \mathcal{N}_i} 1_{\{x(j) = \tau\}} \quad (\tau = 0, 1)$$

be the frequency of type τ in the neighborhood \mathcal{N}_i . In the standard voter model, if the present state is x , then the site i changes its type with the




Figure 1.8: Time evolution of a one-dimensional biased voter model with bias $s = 0.2$.

following rates:

$$\begin{aligned} 0 &\mapsto 1 && \text{with rate } f_1(x), \\ 1 &\mapsto 0 && \text{with rate } f_0(x). \end{aligned}$$

In the biased voter model, this is changed to

$$\begin{aligned} 0 &\mapsto 1 && \text{with rate } (1 + s)f_1(x), \\ 1 &\mapsto 0 && \text{with rate } f_0(x). \end{aligned}$$

Another generalization of the voter model, introduced in [NP99], is defined by the rates

$$\begin{aligned} 0 &\mapsto 1 && \text{with rate } f_1(x)(f_0(x) + \alpha f_1(x)), \\ 1 &\mapsto 0 && \text{with rate } f_0(x)(f_1(x) + \alpha f_0(x)), \end{aligned} \tag{1.18}$$

where $0 \leq \alpha \leq 1$ is a model parameter. Another way of expressing this is to say that if the individual at i is of type τ , then this individual dies with rate

$$f_\tau(x) + \alpha f_{1-\tau}(x), \tag{1.19}$$

and once an individual has died, just as in the normal contact process, it is replaced by a descendant of a uniformly chosen neighbor.

If $\alpha = 1$, then the rate of dying in (1.19) is one and we are back at the standard voter model, but for $\alpha < 1$, individuals die less often if they are

surrounded by a lot of individuals of the other type. In biology, this models *balancing selection*. This is the effect that individuals that differ from their neighbors experience less competition, which results in a selective drive for high biodiversity.

In the social interpretation of the voter model, we may interpret (1.19) as saying that persons change their mind *less* often if they disagree with a lot of neighbors, i.e., the model in (1.18) has “rebellious” behavior.

Numerical simulations, shown in Figure 1.9, suggest that in one dimension and for ranges $R \geq 2$, the model in (1.18) exhibits a phase transition in α . For α sufficiently close to 1, the model behaves essentially as a voter model, with clusters growing in time, but for small values of α (which represent strong rebellious behavior), the cluster size tends to a finite limit.

1.7 Branching and coalescing particles

For each $i, j \in \mathbb{Z}^d$, we define a *coalescing random walk map* $\mathbf{rw}_{ij} : \{0, 1\}^{\mathbb{Z}^d} \rightarrow \{0, 1\}^{\mathbb{Z}^d}$ by

$$\mathbf{rw}_{ij}(x)(k) := \begin{cases} 0 & \text{if } k = i, \\ x(i) \vee x(j) & \text{if } k = j, \\ x(k) & \text{otherwise.} \end{cases} \quad (1.20)$$

Applying \mathbf{rw}_{ij} to a configuration x has the effect that if the site i is occupied by a particle, then this particle jumps to the site j . If there is already a particle at j , then the two particles coalesce.

The interacting particle system with generator

$$G_{\mathbf{rw}}f(x) = \frac{1}{|\mathcal{N}_0|} \sum_{(i,j) \in \mathcal{E}^d} \{f(\mathbf{rw}_{ij}(x)) - f(x)\} \quad (x \in \{0, 1\}^{\mathbb{Z}^d}) \quad (1.21)$$

describes a system of coalescing random walks, where each particle jumps with rate 1 to a uniformly chosen neighboring site, and two particles on the same site coalesce; see Figure 1.10. Likewise, replacing the coalescing random walk map by the *annihilating random walk map* defined as

$$\mathbf{ann}_{ij}(x)(k) := \begin{cases} 0 & \text{if } k = i, \\ x(i) + x(j) \mod(2) & \text{if } k = j, \\ x(k) & \text{otherwise,} \end{cases} \quad (1.22)$$

yields a system of annihilating random walks, that kill each other as soon as two particles land on the same site; see Figure 1.10.




Figure 1.9: Evolution of “rebellious” voter models with $\alpha = 0.8$ and $\alpha = 0.3$, respectively.

For each $i, j \in \mathbb{Z}^d$, we define an *exclusion map* $\text{excl}_{ij} : S^{\mathbb{Z}^d} \rightarrow S^{\mathbb{Z}^d}$ by

$$\text{excl}_{ij}(x)(k) := \begin{cases} x(j) & \text{if } k = i, \\ x(i) & \text{if } k = j, \\ x(k) & \text{otherwise.} \end{cases} \quad (1.23)$$

Applying excl_{ij} to a configuration x has the effect of interchanging the types




Figure 1.10: Systems of coalescing random walks (above) and annihilating random walks (below).

of j and j . The interacting particle system with state space $\{0, 1\}^{\mathbb{Z}^d}$ and generator

$$G_{\text{excl}}f(x) = \frac{1}{|\mathcal{N}_0|} \sum_{(i,j) \in \mathcal{E}^d} \{f(\text{excl}_{ij}(x)) - f(x)\} \quad (x \in \{0, 1\}^{\mathbb{Z}^d}) \quad (1.24)$$

is called the (symmetric) *exclusion process*. In the exclusion process, individual particles move according to random walks, that are independent as long as the particles are sufficiently far apart. Particles never meet, and the total number of particles is preserved.

The previous three maps (coalescing random walk map, annihilating ran-

dom walk map, and exclusion map) can be combined with, e.g., the branching map and death map from (1.6) and (1.7). In particular, adding coalescing random walk or exclusion dynamics to a contact process models displacement (migration) of organisms. Since in many organisms, you actually need two parents to produce offspring, several authors [Nob92, Dur92, Neu94, SS15a] have studied particle systems where the branching map is replaced by the *cooperative branching map*

$$\text{coop}_{ijk}(x)(l) := \begin{cases} 1 & \text{if } l = k, x(i) = 1, x(j) = 1, \\ x(l) & \text{otherwise.} \end{cases} \quad (1.25)$$

See Figure 1.11 for a one-dimensional interacting particle system involving cooperative branching and coalescing random walks.




Figure 1.11: A one-dimensional interacting particle system with cooperative branching and coalescing random walk dynamics.

We define a *killing map* by

$$\text{kill}_{ij}(x)(k) := \begin{cases} 0 & \text{if } k = j, x(i) = 1, x(j) = 1, \\ x(k) & \text{otherwise.} \end{cases} \quad (1.26)$$

In words, this says that if there are particles at i and j , then the particle at i kills the particle at j . Sudbury [Sud97, Sud99] has studied a “biased

annihilating branching process” with generator of the form

$$G_{\text{babp}}f(x) := \lambda \sum_{(i,j) \in \mathcal{E}^1} \{f(\text{bra}_{ij}(x)) - f(x)\} \\ + \sum_{(i,j) \in \mathcal{E}^1} \{f(\text{kill}_{ij}(x)) - f(x)\} \quad (x \in \{0,1\}^{\mathbb{Z}}). \quad (1.27)$$

In the physics literature, this model is known as the Fredrickson-Andersen one spin facilitated model, see formula (26) in [RS03] (with $f = 1$). It is part of the class of *kinetically constrained models*. In the mathematical literature on this subject, a slight variant of the model has been studied [BDT19]. Figure 1.12 shows a simulation of such a system when $\lambda = 0.2$. When λ is small, in the simulations, the process seems to behave similar to systems of branching and coalescing random walks.




Figure 1.12: A system with branching and killing.

1.8 Periodic behavior

An invariant law of an interacting particle system is a probability distribution on the space S^{Λ} of all possible configurations with the property that if the system at time zero is distributed according to this law, then at all later times it is also distributed according to this law. Invariant laws need not be unique. For example, Potts models above the critical point have q different

invariant laws, that are characterized by the color that occupies the majority of the sites.

For all the interacting particle systems and initial states we have considered so far, the system has the property that as time tends to infinity, the distribution of the system converges to an invariant law. This need not always be the case. Perhaps the simplest way in which this can fail is if the system has a *periodic law*, i.e., a law that has the property that if the system at time zero is distributed according to this law, then it returns to this law after a finite time $T > 0$ (the *period*), but the system has a different distribution at all intermediate times $0 < t < T$.

Very little is known rigorously about interacting particle systems with periodic laws. Jahnke and Külske [JK14a] have constructed a three dimensional interacting particle system that has a periodic law. A general result due to Mountford [Mou95] implies that one dimensional systems with finite range interactions cannot have periodic laws. Beyond this, very little is known for spatial models, although there are some studies of periodic behavior in the mean-field limit (see Chapter 3). In particular, it is not known whether periodic laws are possible in two dimensions. The construction in [JK14a] is rather abstract since they do not write down the dynamics of their system explicitly but only prove that such a system exists. Their system also does not have finite range interactions, although the strength of the interaction decays exponentially in the distance, which is almost as good.

Numerical simulations suggest that periodic behavior is not a rare phenomenon. Several interacting particle systems with explicit dynamics are known to exhibit periodic behavior in simulations. All known examples seem to work only in dimensions three and higher, however, which suggests that, perhaps, periodic behavior is not possible in two dimensions.

In the *mutually metastable Ising model*, based on the work of Collet, Formentin and Tovazzi [CFT16], the local state space is $S = \{-1, +1\}^2$ and the lattice is $\Lambda = \mathbb{Z}^3$. We denote an element of S^Λ as $x = (x(i))_{i \in \Lambda}$ where $x(i) = (x^1(i), x^2(i))$ with $x^1(i), x^2(i) \in \{-1, +1\}$ ($i \in \mathbb{Z}^3$). We let

$$M_i^1(x) := \frac{1}{|\mathcal{N}_i|} \sum_{j \in \mathcal{N}_i} x^1(j) \quad \text{and} \quad M_i^2(x) := \frac{1}{|\mathcal{N}_i|} \sum_{j \in \mathcal{N}_i} x^2(j)$$

denote the average values of $x^1(j)$ and $x^2(j)$ among the neighbors of a site i . The *mutually metastable Ising model* with parameters $\alpha, \beta \geq 0$ is the interacting particle system with state space S^Λ that evolves in such a way that

$$\begin{aligned} x^1(i) \text{ flips its value with rate} & \quad e^{-x^1(i)[\beta M_i^1(x) + \alpha x^2(i)]}, \\ x^2(i) \text{ flips its value with rate} & \quad e^{-x^2(i)[\beta M_i^2(x) - \alpha x^1(i)]}. \end{aligned}$$




Figure 1.13: Periodic behavior of the dissipative Ising model with $\alpha = 0.2$ and $\beta = 8$ on a cube with side length 40 and periodic boundary conditions. Colors indicate the state of $x^1(i)$. Good sites have bright colors and corrupted sites have dark colors. Shown is the transition from a predominantly minus state to a predominantly plus state.

When $\alpha = 0$, there is no interaction between the two spins at a given site. In this case $X_t^1 := (X_t^1(i))_{i \in \Lambda}$ and $X_t^2 := (X_t^2(i))_{i \in \Lambda}$ evolve as independent stochastic Ising models.¹⁰ In particular, the effect of the term $-\beta x^1(i)M_i^1(x)$ in the exponent is that the spin $x^1(i)$ does not change its value often if it agrees with most of its neighbors (so that $x^1(i)M_i^1(x) > 0$), and on the other hand changes its value with a high rate if it disagrees with most of its neighbors (so that $x^1(i)M_i^1(x) < 0$). Similarly, the spin $x^2(i)$ has a tendency to align with the majority of its neighbors.

When $\alpha > 0$, then in addition, the spin $x^1(i)$ tries to align with $x^2(i)$, but on the other hand, the spin $x^2(i)$ prefers to have a different value from $x^2(i)$. When most of the spins $x^1(i)$ have the value +1 and most of the spins $x^2(i)$ also have the value +1, then the spins $x^1(i)$ are perfectly satisfied, because they agree with most of their neighbors and also with the other spins $x^2(i)$.

¹⁰The parameter β here is defined differently from the parameter β in Section 1.4. Denoting the latter by β' , one has $\frac{1}{2}\beta' = \frac{1}{|\mathcal{N}_i|}\beta$. Also, the dynamics here are different from the Glauber dynamics of Section 1.4.




Figure 1.14: Three snapshots of a stationary dissipative Ising model with $\alpha = 0.2$ and $\beta = 8$ on a square with side length 100 and periodic boundary conditions. Although the system locally behaves in a periodic way sites do not coordinate their action over longer distances.

But the spins $x^2(i)$ are not completely satisfied. From their point of view, it would be ideal if most of the spins $x^1(i)$ have the value $+1$ but most of the spins $x^2(i)$ have the value -1 . Numerical simulations suggest that the result of this is, for suitable choices of the parameters, that the systems starts to cycle, in a coordinated way, through the states

$$(+1, +1) \mapsto (+1, -1) \mapsto (-1, -1) \mapsto (-1, +1) \mapsto (+1, +1) \mapsto \dots$$

However, this sort of coordinated behavior seems to happen only in dimensions three and more. In two dimensions, even though individual sites still tend to pass through the states $(+1, +1) \mapsto (+1, -1) \mapsto (-1, -1) \mapsto (-1, +1) \mapsto$ in this order, the sites at larger distances from each other no longer coordinate their behavior.

In the *dissipative Ising model*, based on a model described in [DFR13, Tov19, CDFT20], the local state space is again $S = \{-1, +1\}^2$. There are again two model parameters $\alpha, \beta \geq 0$ but the dynamics are different. Letting $M_i(x) := (M_i^1(x) + M_i^2(x))/2$ denote the average value of $x^1(j)$ and $x^2(j)$ for all sites j neighboring the site i , the system evolves in such a way that

$$\begin{aligned} x^1(i) \text{ flips its the value with rate } & e^{-\beta x^1(i) M_i(x)}, \\ x^2(i) \text{ flips its the value with rate } & \alpha 1_{\{x^1(i) = x^2(i)\}}. \end{aligned}$$

We call sites where $x^1(i) = x^2(i)$ *good* sites with spin $x^1(i)$ and sites where $x^1(i) \neq x^2(i)$ *corrupted* sites with spin $x^1(i)$. Note that corrupted sites have no influence on $M_i(x)$. Therefore, we can describe the dynamics informally as follows: sites try to align with their neighbors, but they only see the good

neighbors. After flipping their spin, sites are initially good, but they become corrupt with rate α and stay so until the next time when they flip their spin.

In simulations of the three dimensional model, for a suitable choice of the parameters, we see periodic behavior, where the system cycles through the states

$$(+1, +1) \mapsto (+1, -1) \mapsto (-1, -1) \mapsto (-1, +1) \mapsto (+1, +1) \mapsto \dots$$

in a coordinated way. In two dimensions, we still see this behavior locally, but sites at larger distances from each other no longer coordinate their behavior.

Chapter 2

Continuous-time Markov chains

2.1 Poisson point sets

Let S be a σ -compact¹ metrizable space. We will mainly be interested in the case that $S = \mathbb{R} \times \Lambda$ where Λ is a countable set. We let \mathcal{S} denote the Borel- σ -field on S . A *locally finite measure* on (S, \mathcal{S}) is a measure μ such that $\mu(C) < \infty$ for all compact $C \subset S$.

Let $(\Omega, \mathcal{F}, \mathbb{P})$ be our underlying probability space. A random measure on S is a function $\xi : \Omega \times \mathcal{S} \rightarrow [0, \infty]$ such that for fixed $\omega \in \Omega$, the function $\xi(\omega, \cdot)$ is a locally finite measure on (S, \mathcal{S}) , and for fixed $A \in \mathcal{S}$, the function $\xi(\cdot, A)$ is measurable. By [Kal97, Lemma 1.37], we can think of ξ as a random variable with values in the space of locally finite measures on (S, \mathcal{S}) , equipped with the σ -field generated by the maps $\mu \mapsto \mu(A)$ with $A \in \mathcal{S}$. Then the integral $\int f d\xi$ defines a $[0, \infty]$ -valued random variable for all measurable $f : S \rightarrow [0, \infty]$. There exists a unique measure, denoted by $\mathbb{E}[\xi]$, such that

$$\int f d\mathbb{E}[\xi] = \mathbb{E}\left[\int f d\xi\right]$$

for all measurable $f : S \rightarrow [0, \infty]$. The measure $\mathbb{E}[\xi]$ is called the *intensity* of ξ .

The following result follows from [Kal97, Lemma 10.1 and Prop. 10.4].² Below, $\hat{\mathcal{S}} := \{A \in \mathcal{S} : \overline{A} \text{ is compact}\}$ denotes the set of measurable subsets of S whose closure is compact.

¹This means that there exists a countable collection of compact sets $S_i \subset S$ such that $\bigcup_i S_i = S$.

²In fact, [Kal97, Prop. 10.4] shows that it is possible to construct Poisson point measures on arbitrary measurable spaces, assuming only that the intensity measure is σ -finite, but we will not need this generality.

Proposition 2.1 (Poisson point measures) *Let μ be a locally finite measure on (S, \mathcal{S}) . Then there exists a random measure ξ , unique in distribution, such that for any disjoint $A_1, \dots, A_n \in \hat{\mathcal{S}}$, the random variables $\xi(A_1), \dots, \xi(A_n)$ are independent and $\xi(A_i)$ is Poisson distributed with mean $\mu(A_i)$.*

We call a random measure ξ as in (2.1) a *Poisson point measure* with *intensity* μ . Indeed, one can check that $\mathbb{E}[\xi] = \mu$. We note that $\xi(A) \in \mathbb{N}$ for all $A \in \hat{\mathcal{S}}$. Such measures are called (locally finite) *counting measures*. Each locally finite counting measure ν on S is of the form

$$\nu = \sum_{x \in \text{supp}(\nu)} n_x \delta_x,$$

where $\text{supp}(\nu)$, the support of ν , is a locally finite subset of S , the n_x are positive integers, and δ_x denotes the delta-measure at x . We say that ν is *simple* if $n_x = 1$ for all $x \in \text{supp}(\nu)$. Recall that a measure μ has an *atom* at x is $\mu(\{x\}) > 0$. A measure μ is called *atomless* if it has no atoms, i.e., $\mu(\{x\}) = 0$ for all $x \in S$. The already mentioned [Kal97, Prop. 10.4] tells us the following.

Lemma 2.2 (Simple Poisson point measures) *Let ξ be a Poisson point measure with locally finite intensity μ . Then ξ is a.s. simple if and only if μ is atomless.*

If μ is atomless, then a Poisson point measure ξ with intensity μ is characterized by its support $\omega := \text{supp}(\xi)$. We call ω a *Poisson point set* with intensity μ . Intuitively, ω is a set such that $\mathbb{P}[\omega \cap dx \neq \emptyset] = \mu(dx)$, independently for each infinitesimal subset $dx \subset S$.

For any counting measure ν on S and measurable function $f : S \rightarrow [0, 1]$ we introduce the notation

$$f^\nu := \prod_{i=1}^n f(x_i) \quad \text{where} \quad \nu = \sum_i \delta_{x_i}.$$

Here, by definition, $f^0 := 1$, where 0 denotes the counting measure that is identically zero. Alternatively, our definition says that

$$f^\nu = e^{\int (\log f) d\nu},$$

where $\log 0 := -\infty$ and $e^{-\infty} := 0$. It is easy to see that $f^\nu f^{\nu'} = f^{\nu+\nu'}$.

Lemma 2.3 (Laplace functionals) *Let μ be a locally finite measure on (S, \mathcal{S}) and let ξ be a Poisson point measure with intensity μ . Then*

$$\mathbb{E}[(1 - f)^\xi] = e^{-\int f d\mu} \quad (2.1)$$

for each measurable $f : S \rightarrow [0, 1]$. Conversely, if ξ is a random counting measure and (2.1) holds for all continuous, compactly supported f , then ξ is a Poisson point measure with intensity μ .

Proof The fact that Poisson point measures satisfy (2.1) is proved in [Kal97, Lemma 10.2], which is written in terms of $-\log f$, rather than f . The fact that (2.1) determines the law of ξ uniquely follows from [Kal97, Lemma 10.1]. ■

Formula (2.1) can be interpreted in terms of thinning. Consider a counting measure $\nu = \sum_i \delta_{x_i}$, let $f : S \rightarrow [0, 1]$ be measurable, and let χ_i be independent *Bernoulli random variables* (i.e., random variables with values in $\{0, 1\}$) with $\mathbb{P}[\chi_i = 1] = f(x_i)$. Then the random counting measure

$$\nu' := \sum_i \chi_i \delta_{x_i}$$

is called an *f-thinning* of the counting measure ν . Note that

$$\mathbb{P}[\nu' = 0] = \prod_i \mathbb{P}[\chi_i = 0] = (1 - f)^\nu.$$

In view of this, the left-hand side of (2.1) can be interpreted as the probability that after thinning the random counting measure ξ with f , no points remain. By [Kal97, Lemma 10.1], knowing this probability for each continuous, compactly supported f uniquely determines the law of a random counting measure.

Using Lemma 2.3, it is easy to prove that if ξ_1 and ξ_2 are independent Poisson point measures with intensities μ_1 and μ_2 , then $\xi_1 + \xi_2$ is a Poisson point measure with intensity $\mu_1 + \mu_2$. We also mention [Kal97, Lemma 10.17], which says the following.

Lemma 2.4 (Poisson points on the halfline) *Let $(\tau_k)_{k \geq 0}$ be real random variables such that $\tau_0 = 0$ and $\sigma_k := \tau_k - \tau_{k-1} > 0$ ($k \geq 1$). Then $\omega := \{\tau_k : k \geq 1\}$ is a Poisson point set on $[0, \infty)$ with intensity $c\ell$, where ℓ denotes the Lebesgue measure, if and only if the random variables $(\sigma_k)_{k \geq 1}$ are i.i.d. exponentially distributed with mean c^{-1} .*

2.2 Transition probabilities and generators

Let S be any finite set. A (real) matrix indexed by S is a collection of real constants $A = (A(x, y))_{x, y \in S}$. We calculate with such matrices in the same way as with normal finite matrices. Thus, the product AB of two matrices is defined as

$$(AB)(x, z) := \sum_{y \in S} A(x, y)B(y, z) \quad (x, z \in S).$$

We let 1 denote the identity matrix $1(x, y) = 1_{\{x=y\}}$ and define A^n in the obvious way, with $A^0 := 1$. If $f : S \rightarrow \mathbb{R}$ is a function, then we also define

$$Af(x) := \sum_{y \in S} A(x, y)f(y) \quad \text{and} \quad fA(y) := \sum_{x \in S} f(x)A(x, y). \quad (2.2)$$

A *probability kernel* on S is a matrix $K = (K(x, y))_{x, y \in S}$ such that $K(x, y) \geq 0$ ($x, y \in S$) and $\sum_{y \in S} K(x, y) = 1$ ($x \in S$). Clearly, the composition of two probability kernels yields a third probability kernel. A probability kernel is *deterministic* if it is of the form

$$K_m(x, y) := \begin{cases} 1 & \text{if } x = m(y), \\ 0 & \text{otherwise,} \end{cases}$$

for some function $m : S \rightarrow S$. It is easy to see that the space of all probability kernels on a finite set S is convex, and the deterministic probability kernels are exactly the extremal elements of this set. It follows that each probability kernel can be written as a convex combination of deterministic probability kernels. Another way to say this is that for each probability kernel K on S , it is possible to find a random map $M : S \rightarrow S$ such that³

$$K(x, y) = \mathbb{P}[M(x) = y] \quad (x, y \in S). \quad (2.3)$$

A formula of this form is called a *random mapping representation* of the probability kernel K .

A *Markov semigroup* is a collection of probability kernels $(P_t)_{t \geq 0}$ such that

$$\lim_{t \downarrow 0} P_t = P_0 = 1 \quad \text{and} \quad P_s P_t = P_{s+t} \quad (s, t \geq 0).$$

Each such Markov semigroup is of the form

$$P_t = e^{tG} := \sum_{n=0}^{\infty} \frac{1}{n!} (tG)^n,$$

³Indeed, this formula says nothing else than $K = \sum_m \mathbb{P}[M = m]K_m$, where the sum runs over all maps $m : S \rightarrow S$, and K_m is the deterministic kernel defined by the map m .

where the *generator* G is a matrix of the form

$$G(x, y) \geq 0 \quad (x \neq y) \quad \text{and} \quad \sum_y G(x, y) = 0. \quad (2.4)$$

By definition, we say that a function f that is defined on an interval $I \subset \mathbb{R}$ is *piecewise constant* if each compact subinterval of I can be divided into finitely many subintervals, so that f is constant on each subinterval. By definition, a *Markov process* with semigroup $(P_t)_{t \geq 0}$ is a stochastic process $X = (X_t)_{t \geq 0}$ with values in S and piecewise constant, right-continuous sample paths, such that

$$\mathbb{P}[X_u \in \cdot \mid (X_s)_{0 \leq s \leq t}] = P_{u-t}(X_t, \cdot) \quad \text{a.s.} \quad (0 \leq t \leq u). \quad (2.5)$$

Here, in the left-hand side, we condition on the σ -field generated by the random variables $(X_s)_{0 \leq s \leq t}$. One can prove that formula (2.5) is equivalent to the statement that

$$\begin{aligned} & \mathbb{P}[X_0 = x_0, \dots, X_{t_n} = x_n] \\ &= \mathbb{P}[X_0 = x_0] P_{t_1-t_0}(x_0, x_1) \cdots P_{t_n-t_{n-1}}(x_{n-1}, x_n) \quad (0 < t_1 < \dots < t_n). \end{aligned} \quad (2.6)$$

From this last formula, we see that for each initial law $\mathbb{P}[X_0 = \cdot] = \mu$, there is a unique Markov process with semigroup $(P_t)_{t \geq 0}$ and this initial law. Moreover, recalling our notation (2.2), we see that

$$\mu P_t(x) = \mathbb{P}[X_t = x] \quad (x \in S)$$

is the law of the process at time t . It is custom to let \mathbb{P}^x denote the law of the Markov process with deterministic initial state $X_0 = x$ a.s. We note that

$$\mathbb{P}^x[X_t = y] = P_t(x, y) = 1_{\{x=y\}} + tG(x, y) + O(t^2) \quad \text{as } t \downarrow 0.$$

For $x \neq y$, we call $G(x, y)$ the *rate* of jumps from x to y . Intuitively, if the process is in x , then in the next infinitesimal time interval of length dt it has a probability $G(x, y)dt$ to jump to y , independently for all $y \neq x$.

Let X be the process started in x and let $\tau := \inf\{t \geq 0 : X_t \neq x\}$. Then one can show that τ is exponentially distributed with mean r^{-1} , where $r := \sum_{y \neq x} G(x, y)$ is the total rate of all jumps from x . Moreover,

$$\mathbb{P}^x[X_\tau = y] = \frac{G(x, y)}{\sum_{z \neq x} G(x, z)} \quad (y \in S, y \neq x). \quad (2.7)$$

Conditional on $X_\tau = y$, the time of the next jump is again exponentially distributed, and this leads to a construction of $(X_t)_{t \geq 0}$ based on an *embedded Markov chain* with transition kernel $K(x, y)$ given by the right-hand side of (2.7), and exponential holding times. For us, a somewhat different construction based on maps that are applied at Poissonian times will be more useful.

2.3 Poisson construction of Markov processes

Let S be a finite set. Let \mathcal{K} be a finite set whose elements are probability kernels on S and let $(r_K)_{K \in \mathcal{K}}$ be nonnegative constants. Then it is straightforward to check that setting

$$Gf := \sum_{K \in \mathcal{K}} r_K \{Kf - f\}$$

defines a Markov generator. The following exercise says that conversely, each Markov generator can be written in this form, where we can even choose the set \mathcal{K} so that it has only one element.

Exercise 2.5 *Let S be a finite set. Show that each Markov generator G on S can be written in the form $Gf = r\{Kf - f\}$, where $r \geq 0$ is a constant and K is a probability kernel on S . Hint: first add a multiple of the identity matrix to G to make all diagonal entries nonnegative and then normalize.*

If all kernels in the set \mathcal{K} are deterministic, then our expression for G takes the form

$$Gf(x) = \sum_{m \in \mathcal{G}} r_m \{f(m(x)) - f(x)\}, \quad (2.8)$$

where \mathcal{G} is a finite set whose elements are maps $m : S \rightarrow S$ and $(r_m)_{m \in \mathcal{G}}$ are nonnegative constants. We call (2.8) a *random mapping representation* of the generator G .

Exercise 2.6 *Let S be a finite set. Show that each Markov generator G on S has a random mapping representation.*

Let $(P_t)_{t \geq 0}$ be the Markov semigroup generated by G . We will see that the random mapping representation (2.8) can be used to give a random mapping representation for the probability kernels $(P_t)_{t \geq 0}$. We equip the space $\mathcal{G} \times \mathbb{R}$ with the measure

$$\rho(\{m\} \times A) := r_m \ell(A) \quad (m \in \mathcal{G}, A \in \mathcal{B}(\mathbb{R})), \quad (2.9)$$

where $\mathcal{B}(\mathbb{R})$ denotes the Borel- σ -field on \mathbb{R} and ℓ denotes the Lebesgue measure. Let ω be a Poisson point set with intensity ρ . We claim that for each $t \in \mathbb{R}$, there exists at most one $m \in \mathcal{G}$ such that $(m, t) \in \omega$. To see this, we note that for each $m \in \mathcal{G}$, the set

$$\xi_m := \sum_{t: (m, t) \in \omega} \delta_t$$

is a Poisson point measure on \mathbb{R} with intensity $r_m \ell$. Since the sets $\mathbb{R} \times \{m\}$ ($m \in \mathcal{G}$) are disjoint, the random measures ξ_m ($m \in \mathcal{G}$) are independent, and hence for each $m \neq m'$, the measure $\xi_m + \xi_{m'}$ is a Poisson point measure on \mathbb{R} with intensity $(r_m + r_{m'})\ell$. Since the Lebesgue measure is atomless, by Lemma 2.2, this Poisson point measure is simple, so there are no times $t \in \mathbb{R}$ for which both $(m, t) \in \omega$ and $(m', t) \in \omega$. In view of this, we can unambiguously define a random function $\mathbb{R} \ni t \mapsto \mathbf{m}_t^\omega$ by setting

$$\mathbf{m}_t^\omega := \begin{cases} m & \text{if } (m, t) \in \omega, \\ 1 & \text{otherwise,} \end{cases}$$

where we write 1 to denote the identity map. For $s \in \mathbb{R}$ and $x \in S$, we will be interested in piecewise constant, right-continuous functions $[s, \infty) \ni t \mapsto X_t \in S$ that solve the equation

$$X_s = x \quad \text{and} \quad X_t = \mathbf{m}_t^\omega(X_{t-}) \quad (t > s), \quad (2.10)$$

where $X_{t-} := \lim_{s \uparrow t} X_s$ denotes the value of the function $t \mapsto X_t$ just before time t . Let us write

$$\omega_{s,u} := \{(m, t) \in \omega : t \in (s, u]\} \quad (s \leq u). \quad (2.11)$$

Since $\rho(\omega_{s,u}) = (u - s) \sum_{m \in \mathcal{G}} r_m < \infty$, the set $\omega_{s,u}$ contains only finitely many points, which we can order as

$$\omega_{s,u} = \{(m_1, t_1), \dots, (m_n, t_n)\} \quad \text{with} \quad t_1 < \dots < t_n. \quad (2.12)$$

The unique solution of (2.10) is then given by

$$\begin{aligned} X_t &= x && \text{for } t \in [s, t_1), \\ X_t &= m_1(x) && \text{for } t \in [t_1, t_2), \\ X_t &= m_2 \circ m_1(x) && \text{for } t \in [t_2, t_2), \quad \text{etc.} \end{aligned}$$

To formalize this, we define a collection of random maps $(\mathbf{X}_{s,u})_{s \leq u}$ by

$$\mathbf{X}_{s,u} := m_n \circ \dots \circ m_1,$$

where m_1, \dots, m_n are as in (2.12). Here, by definition, the composition of no maps is the identity map, i.e., $\mathbf{X}_{s,u}$ is the identity map if $\omega_{s,u} = \emptyset$. It is not hard to see that

$$\lim_{t \downarrow s} \mathbf{X}_{s,t} = \mathbf{X}_{s,s} = 1 \quad \text{and} \quad \mathbf{X}_{t,u} \circ \mathbf{X}_{s,t} = \mathbf{X}_{s,u} \quad (s \leq t \leq u), \quad (2.13)$$

i.e., the maps $(\mathbf{X}_{s,t})_{s \leq t}$ form a *stochastic flow*. Also, $\mathbf{X}_{s,t}$ is right-continuous in both s and t . Finally, $(\mathbf{X}_{s,t})_{s \leq t}$ has *independent increments* in the sense that

$$\mathbf{X}_{t_0, t_1}, \dots, \mathbf{X}_{t_{n-1}, t_n} \quad \text{are independent} \quad \forall t_0 < \dots < t_n.$$

The unique solution of (2.10) is now given by

$$X_t = \mathbf{X}_{s,t}(x) \quad (t \geq s).$$

The following proposition says that $(X_t)_{t \geq s}$ is in fact a Markov process. Formula (2.15) below moreover shows that the random map $\mathbf{X}_{s,t}$ is in fact a random mapping representation of the kernel P_{t-s} .

Proposition 2.7 (Poisson construction of Markov processes)

Define a stochastic flow $(\mathbf{X}_{s,t})_{s \leq t}$ as above in terms of a Poisson point set ω . Let X_0 be an S -valued random variable, independent of ω . Then

$$X_t := \mathbf{X}_{0,t}(X_0) \quad (t \geq 0) \tag{2.14}$$

defines a Markov process $X = (X_t)_{t \geq 0}$ with generator (2.8).

Proof The process $X = (X_t)_{t \geq 0}$, defined in (2.14), has piecewise constant, right-continuous sample paths. Define

$$P_t(x, y) := \mathbb{P}[\mathbf{X}_{s, s+t}(x) = y] \quad (t \geq 0), \tag{2.15}$$

where the definition does not depend on the choice of $s \in \mathbb{R}$ since the law of the Poisson process ω is invariant under translations in the time direction. Using the fact that $(\mathbf{X}_{s,t})_{s \leq t}$ has independent increments and X_0 is independent of ω , we see that the finite-dimensional distributions of X satisfy (2.6).

It follows from (2.13) that the probability kernels $(P_t)_{t \geq 0}$ defined in (2.15) form a Markov semigroup. To see that its generator G is given by (2.8), we observe that by the properties of Poisson processes,

$$\mathbb{P}[|\omega_{0,t}| \geq 2] = O(t^2) \quad \text{as } t \downarrow 0,$$

while

$$\mathbb{P}[\omega_{0,t} = \{(m, s)\} \text{ for some } s \in (0, t)] = r_m t + O(t^2) \quad \text{as } t \downarrow 0.$$

Using this, it follows that for any $f : S \rightarrow \mathbb{R}$, as $t \downarrow 0$,

$$P_t f(x) = \mathbb{E}[f(\mathbf{X}_{0,t}(x))] = f(x) + t \sum_{m \in \mathcal{G}} r_m \{f(m(x)) - f(x)\} + O(t^2).$$

Since $P_t f = f + tGf + O(t^2)$, this proves that G is given by (2.8). ■

2.4 Examples of Poisson representations

Random mapping representations of generators are generally not unique. Consider the following example. We choose the state space $S := \{0, 1\}$ and the generator G defined by⁴

$$G(0, 1) := 2 \quad \text{and} \quad G(1, 0) := 1,$$

which corresponds to a Markov process that jumps

$$0 \mapsto 1 \quad \text{with rate } 2 \quad \text{and} \quad 1 \mapsto 0 \quad \text{with rate } 1.$$

We define maps **down**, **up**, and **swap**, mapping the state space $S = \{0, 1\}$ into itself, by

$$\left. \begin{array}{l} \text{down}(x) := 0, \\ \text{up}(x) := 1, \\ \text{swap}(x) := 1 - x \end{array} \right\} \quad (x \in S).$$




Figure 2.1: Two stochastic flows representing the same Markov process.

It is straightforward to check that the generator G can be represented in terms of the set of maps $\mathcal{G} := \{\text{down}, \text{up}\}$ as

$$Gf(x) = r_{\text{down}}\{f(\text{down}(x)) - f(x)\} + r_{\text{up}}\{f(\text{up}(x)) - f(x)\}, \quad (2.16)$$

⁴By (2.4), if G is a Markov generator, then $G(x, x) = -\sum_{y: y \neq x} G(x, y)$, so in order to specify a Markov generator, it suffices to give its off-diagonal elements.

where

$$r_{\text{down}} := 1 \quad \text{and} \quad r_{\text{up}} := 2.$$

But the same generator G can also be represented in terms of the set of maps $\mathcal{G}' := \{\text{swap}, \text{up}\}$ as

$$Gf(x) = r'_{\text{swap}}\{f(\text{down}(x)) - f(x)\} + r'_{\text{up}}\{f(\text{up}(x)) - f(x)\}, \quad (2.17)$$

where

$$r'_{\text{swap}} := 1 \quad \text{and} \quad r'_{\text{up}} := 1.$$

The random mapping representations (2.16) and (2.17) lead to different ways to construct the same Markov process. In the first construction, we start with a Poisson point set $\omega \subset \mathcal{G} \times \mathbb{R}$, which then defines a stochastic flow $(\mathbf{X}_{s,t})_{s \leq t}$, while in the second construction, we start with a Poisson point set $\omega' \subset \mathcal{G}' \times \mathbb{R}$, which defines a different stochastic flow $(\mathbf{X}'_{s,t})_{s \leq t}$, that nevertheless can be used to construct (in distribution) the same Markov process.

The situation is illustrated in Figure 2.1. Note that in the second representation, both the maps **swap** and **up** make the process jump $1 \mapsto 0$ if its previous state is 1. Therefore, the total rate of jumps $1 \mapsto 0$ is

$$r'_{\text{swap}} + r'_{\text{up}} = 2,$$

just as in the first representation. Note that the picture in Figure 2.1 is a bit misleading since it suggests the processes arising from the two constructions are almost surely equal, while in reality they are only equal in distribution.

2.5 Countable state space

Continuous-time Markov processes with countable state space (also known as continuous-time Markov chains) can in many ways be treated in the same way as those with a finite state space, but there are some complications, which mainly stem from the fact that such processes may *explode*. A good general reference for this material is [Lig10, Chapter 2]. In later chapters, we will need a few facts about the Poisson construction of continuous-time Markov chains, which we conveniently collect here. The reader is advised to only glance over this section at a first reading and only return to it when the need arises.

Let S be a countable set, let \mathcal{G} be a countable set whose elements are maps $m : S \rightarrow S$, and let $(r_m)_{m \in \mathcal{G}}$ be nonnegative constants. Then as we did for finite state space, one can construct a Poisson point subset ω of $\mathcal{G} \times \mathbb{R}$ with intensity measure ρ as defined in (2.9). Then as in the finite case, for

each time $t \in \mathbb{R}$, there is at most one $m \in \mathcal{G}$ such that $(m, t) \in \omega$, so we can again unambiguously define a random function $\mathbb{R} \ni t \mapsto \mathbf{m}_t^\omega$ by setting $\mathbf{m}_t^\omega := m$ if $(m, t) \in \omega$ and $:= 1$ otherwise. It is now natural to ask if the evolution equation (2.10) has a unique piecewise constant, right-continuous solution. The following lemma gives a partial answer to this question. Below, we let $S \cup \{\infty\}$ denote the *one-point compactification* of S . By definition, a sequence $x_n \in S$ converges to ∞ if for each finite set $S' \subset S$, there exists an m such that $x_n \notin S'$ for all $n \geq m$.

Proposition 2.8 (Process up to explosion time) *Assume that*

$$\sum_{m: m(x) \neq x} r_m < \infty \quad \text{for all } x \in S. \quad (2.18)$$

Then, almost surely, for each $x \in S$, there exists a random time $0 < \tau \leq \infty$ and a piecewise constant, right-continuous function $[0, \tau) \ni t \mapsto X_t \in S$ that are uniquely determined by the following conditions:

- (i) $X_s = x$ and $X_t = \mathbf{m}_t^\omega(X_{t-})$ ($0 < t < \tau$),
- (ii) if $\tau < \infty$, then $\lim_{t \rightarrow \tau} X_t = \infty$.

Proof Note that condition (2.18) says that the total rate of all maps that take the process out of a given state x is finite for each $x \in S$. This allows us to inductively define times $0 = t_0 < t_1 < \dots$ and points $x = x_0, x_1, \dots$ by setting

$$t_{n+1} := \inf\{t > t_n : \mathbf{m}_t^\omega(x_n) \neq x_n\} \quad \text{and} \quad x_{n+1} := \mathbf{m}_{t_{n+1}}^\omega(x_n) \quad (n \geq 0).$$

Setting $\tau := \lim_{n \rightarrow \infty} t_n$ and $X_t := x_n$ for $t \in [t_n, t_{n+1})$, we see that $[0, \tau) \ni t \mapsto X_t \in S$ is a piecewise constant, right-continuous function that satisfies (i). To prove also (ii), imagine X_t does not converge to ∞ as $t \rightarrow \infty$. By the definition of the one-point compactification, this implies that there exist a finite set $S' \subset S$ and times $s_n \rightarrow \tau$ such that $X_{s_n} \in S'$. This, in turn, implies that the function X_t has infinitely often jumped out of a point $x \in S'$, which is only possible if the set

$$\bigcup_{x \in S'} \{t \in [0, \tau) : (t, m) \in \omega \text{ for some } m \text{ with } m(x) \neq x\}$$

is infinite. By our assumption (2.18), this is only possible if $\tau = \infty$.

To see that the conditions (i) and (ii) determine the time τ and function $[0, \tau) \ni t \mapsto X_t$ uniquely, assume that τ' and $t \mapsto X'_t$ also satisfy (i) and (ii). Let t_0, t_1, \dots and $\tau := \lim_{n \rightarrow \infty} t_n$ be defined as before. Then by induction,

for each $n \geq 1$, we have $\tau \geq t_n$ and $X'_t = X_t$ for all $t \in [0, t_n]$. It follows that $\tau' \geq \tau$ and $X'_t = X_t$ for all $0 \leq t < \tau$. If $\tau = \infty$ we are done. If $\tau < \infty$, then condition (ii) shows that it is not possible to extend the function $t \mapsto X'_t$ beyond time τ in a piecewise constant way, which proves that $\tau' \leq \tau$. ■

The time τ from Proposition 2.8 is called the *explosion time*. If $\tau < \infty$, then it is natural to extend the function $t \mapsto X_t$ by setting

$$X_t := \infty \quad (t \geq \tau).$$

With this definition, one can use the unique solutions of the evolution equation to define a stochastic flow $(\mathbf{X}_{s,u})_{s \leq u}$ as in the finite case, where now $\mathbf{X}_{s,u}$ is a random map mapping the space $S \cup \{\infty\}$ into itself, with $\mathbf{X}_{s,u}(\infty) := \infty$ for all $s \leq u$. One can check that the stochastic flow $(\mathbf{X}_{s,u})_{s \leq u}$ has independent increments and defines a Markov process with state space $S \cup \{\infty\}$ as in Proposition 2.7. If $\tau = \infty$ a.s., then we say that this Markov process is *nonexplosive*. In the opposite case, it is *explosive*.

In general, there is no single good way to check whether a given continuous-time Markov chain is explosive, but there are several useful methods that are often applicable. A discussion of this falls out of the scope of these lecture notes.

Chapter 3

The mean-field limit

3.1 Processes on the complete graph

In Chapter 1, we have made acquaintances with a number of interacting particle systems. While some properties of these systems sometimes turn out easy to prove, other seemingly elementary questions can sometimes be remarkably difficult. A few examples of such hard problems have been mentioned in Chapter 1. In view of this, interacting particle systems are being studied by a range of different methods, from straightforward numerical simulations as we have seen in Chapter 1, to nonrigorous renormalization group techniques and rigorous mathematical methods. All these approaches complement each other. In addition, when a given problem appears too hard, one often looks for simpler models that (one hopes) still catch the essence, or at least some essential features of the behavior that one is interested in.

A standard way to turn a difficult model into an (often) much easier model is to take the *mean-field limit*, which we explain in the present chapter. Basically, this means that one replaces the graph structure of the underlying lattice that one is really interested in (in practice often \mathbb{Z}^d) by the structure of the complete graph with N vertices, and then takes the limit $N \rightarrow \infty$. As we will see, many properties of “real” interacting particle systems are already reflected in these mean-field models. In particular, phase transitions can often already be observed and even the values of critical exponents of high-dimensional models are correctly predicted by the mean-field model. In view of this, studying the mean-field limit is a wise first step in the study of any more complicated model that one may encounter.

Of course, not all phenomena can be captured by replacing the graph structure that one is really interested in by the complete graph. Comparing the real model with the mean-field model, one can learn which elements of

the observed behavior are a consequence of the specific spatial structure of the lattice, and which are not. Also for this reason, studying the mean-field limit should be part of a complete study of any interacting particle system.

3.2 The mean-field limit of the Ising model

In this section we study the mean-field Ising model, also known as the *Curie-Weiss model*, with Glauber dynamics.

We recall from formulas (1.11) and (1.12) in Chapter 1 that the Ising model is an interacting particle system with local state space $S = \{-1, +1\}$, where each site i updates its spin value $x(i) \in \{-1, +1\}$ at rate one. When a spin value is updated, the probability that the new value is $+1$ resp. -1 is proportional to $e^{\beta N_{x,i}(+1)}$ resp. $e^{\beta N_{x,i}(-1)}$, where $N_{x,i}(\sigma) := \sum_{j \in \mathcal{N}_i} 1_{\{x(j)=\sigma\}}$ denotes the number of neighboring sites that have the spin value σ .

For the aim of taking the mean-field model, it will be convenient to formulate the model slightly differently. We let

$$\bar{N}_{x,i} := \frac{1}{|\mathcal{N}_i|} \sum_{j \in \mathcal{N}_i} 1_{\{x(j)=\sigma\}}$$

denote the fraction of neighbors that have the spin value σ , and consider the model where (compare (1.12))

$$\text{site } i \text{ flips to the value } \sigma \text{ with rate } \frac{e^{\beta \bar{N}_{x,i}(\sigma)}}{\sum_{\tau \in S} e^{\beta \bar{N}_{x,i}(\tau)}}. \quad (3.1)$$

Assuming that $|\mathcal{N}_i|$ is just a constant that does not depend on $i \in \Lambda$ (as is the case, e.g., for the model on \mathbb{Z}^d), this is just a reparametrization of the original model where the parameter β is replaced by $\beta/|\mathcal{N}_i|$.

We now wish to construct the mean-field model, i.e., the model on a complete graph Λ_N with $|\Lambda_N| = N$ vertices (sites), where each site is a neighbor of each other site. For mathematical simplicity, we even count a site as a neighbor of itself, i.e., we set

$$\mathcal{N}_i := \Lambda_N \quad \text{and} \quad |\mathcal{N}_i| = N.$$

A consequence of this choice is that the *average magnetization*

$$\bar{X}_t := \frac{1}{N} \sum_{i \in \Lambda_N} X_t(i) \quad (t \geq 0)$$

forms a Markov process $\bar{X} = (\bar{X}_t)_{t \geq 0}$. Indeed, \bar{X}_t takes values in the space

$$\left\{ -1, -1 + \frac{2}{N}, \dots, 1 - \frac{2}{N}, 1 \right\},$$

and jumps

$$\begin{aligned} \bar{x} \mapsto \bar{x} + \frac{2}{N} & \quad \text{with rate} \quad N_x(-1) \frac{e^{\beta N_x(+1)/N}}{e^{\beta N_x(-1)/N} + e^{\beta N_x(+1)/N}}, \\ \bar{x} \mapsto \bar{x} - \frac{2}{N} & \quad \text{with rate} \quad N_x(+1) \frac{e^{\beta N_x(-1)/N}}{e^{\beta N_x(-1)/N} + e^{\beta N_x(+1)/N}}, \end{aligned}$$

where $N_x(\sigma) := N_{x,i}(\sigma) = \sum_{j \in \Lambda_n} 1_{\{x(j)=\sigma\}}$ does not depend on $i \in \Lambda_N$. We observe that

$$N_x(+1)/N = (1 + \bar{x})/2 \quad \text{and} \quad N_x(-1)/N = (1 - \bar{x})/2.$$

In view of this, we can rewrite the jump rates of \bar{X} as

$$\begin{aligned} \bar{x} \mapsto \bar{x} + \frac{2}{N} & \quad \text{with rate} \quad N(1 - \bar{x})/2 \frac{e^{\beta(1+\bar{x})/2}}{e^{\beta(1-\bar{x})/2} + e^{\beta(1+\bar{x})/2}}, \\ \bar{x} \mapsto \bar{x} - \frac{2}{N} & \quad \text{with rate} \quad N(1 + \bar{x})/2 \frac{e^{\beta(1-\bar{x})/2}}{e^{\beta(1-\bar{x})/2} + e^{\beta(1+\bar{x})/2}}. \end{aligned}$$

In particular, since these rates are a function of \bar{x} only (and do not depend on other functions of $x = (x(i))_{i \in \Lambda_N}$), we see that $\bar{X} = (\bar{X}_t)_{t \geq 0}$, on its own, is a Markov process. (This argument will be made rigorous in Section 3.4 below.) Cancelling a common factor $e^{\beta/2}$ in the nominator and denominator of the rates, we can simplify our formulas a bit to

$$\begin{aligned} \bar{x} \mapsto \bar{x} + \frac{2}{N} & \quad \text{with rate} \quad r_+(\bar{x}) := N(1 - \bar{x})/2 \frac{e^{\beta\bar{x}/2}}{e^{-\beta\bar{x}/2} + e^{\beta\bar{x}/2}}, \\ \bar{x} \mapsto \bar{x} - \frac{2}{N} & \quad \text{with rate} \quad r_-(\bar{x}) := N(1 + \bar{x})/2 \frac{e^{-\beta\bar{x}/2}}{e^{-\beta\bar{x}/2} + e^{\beta\bar{x}/2}}. \end{aligned} \tag{3.2}$$

In Figure 3.1 we can see simulations of the Markov process in (3.2) on a lattice with $N = 10, 100, 1000$, and $10,000$ sites, respectively. It appears that in the limit $N \rightarrow \infty$, the process \bar{X}_t is given by a smooth, deterministic function.

It is not hard to guess what this function is. Indeed, denoting the generator of the process in (3.2) by $\bar{G}_{N,\beta}$, we see that the *local drift* of the process \bar{X} is given by

$$\mathbb{E}^{\bar{x}}[\bar{X}_t] = \bar{x} + t g_{\beta}(\bar{x}) + O(t^2) \quad \text{where} \quad g_{\beta}(\bar{x}) := \bar{G}_{N,\beta} f(\bar{x}) \quad \text{with} \quad f(\bar{x}) := \bar{x}.$$




Figure 3.1: The mean-field Ising model on lattice with $N = 10, 100, 1000$, and $10,000$ sites, respectively. In these simulations, the parameter is $\beta = 3$, and the initial state is $\bar{X}_0 = 0.1$, except in the first picture, where $\bar{X}_0 = 0.2$.

We calculate

$$\begin{aligned}
 g_\beta(\bar{x}) &= r_+(\bar{x}) \cdot \frac{2}{N} - r_-(\bar{x}) \cdot \frac{2}{N} = \frac{(1 - \bar{x})e^{\beta\bar{x}/2} - (1 + \bar{x})e^{-\beta\bar{x}/2}}{e^{\beta\bar{x}/2} + e^{-\beta\bar{x}/2}} \\
 &= \frac{e^{\beta\bar{x}/2} - e^{-\beta\bar{x}/2}}{e^{\beta\bar{x}/2} + e^{-\beta\bar{x}/2}} - \bar{x} = \tanh\left(\frac{1}{2}\beta\bar{x}\right) - \bar{x}.
 \end{aligned} \tag{3.3}$$

Note that the constant N cancels out of this formula. In view of this, by some law of large numbers (that will be made rigorous in Theorem 3.2 below), we expect $(\bar{X}_t)_{t \geq 0}$ to converge in distribution, as $N \rightarrow \infty$, to a solution of the differential equation

$$\frac{\partial}{\partial t} \bar{X}_t = g_\beta(\bar{X}_t) \quad (t \geq 0). \tag{3.4}$$

3.3 Analysis of the mean-field model

Assuming the correctness of (3.4) for the moment, we can study the behavior of the mean-field Ising model \bar{X} in the limit that we first send $N \rightarrow \infty$, and then $t \rightarrow \infty$. A simple analysis of the function g_β (see Figure 3.2) reveals that the differential equation (3.4) has a single fixed point for $\beta \leq 2$, and three fixed points for $\beta > 2$. Here, with a *fixed point* of the differential equation,

we mean a point z such that $\bar{x}_0 = z$ implies $\bar{x}_t = z$ for all $t \geq 0$, i.e., this is a point such that $g_\beta(z) = 0$.




Figure 3.2: The drift function g_β for $\beta = 1.8, 2, 2.1$, and 2.3 , respectively. For $\beta > 2$, the fixed point $\bar{x} = 0$ becomes unstable and two new fixed points appear.

Indeed, using the facts that \tanh is an odd function that is concave on $[0, \infty)$ and satisfies $\frac{\partial}{\partial x} \tanh(x)|_{x=0} = 1$, we see that:

- For $\beta \leq 2$, the equation $g_\beta(x) = 0$ has the unique solution $x = 0$.
- For $\beta > 2$, the equation $g_\beta(x) = 0$ has three solutions $x_- < 0 < x_+$.

For $\beta \leq 2$, solutions to the differential equation (3.4) converge to the unique fixed point $x = 0$ as time tends to zero. On the other hand, for $\beta > 2$, the fixed point $x = 0$ becomes unstable. Solutions \bar{X} to the differential equation (3.4) starting in $\bar{X}_0 > 0$ converge to x_+ , while solutions starting in $\bar{X}_0 < 0$ converge to x_- .

In Figure 3.3, we have plotted the three fixed points $x_- < 0 < x_+$ as a function of β , and indicated their domains of attraction. The function

$$x_{\text{upp}}(\beta) := \begin{cases} 0 & \text{if } \beta \leq 2, \\ \text{the unique pos. sol. of } \tanh(\frac{1}{2}\beta x) = x & \text{if } \beta > 2 \end{cases} \quad (3.5)$$




Figure 3.3: Fixed points of the mean-field Ising model as a function of β , with their domains of attraction. The upper fixed point as a function of β is indicated with a bold line.

plays a similar role as the spontaneous magnetization $m_*(\beta)$ for the Ising model on \mathbb{Z}^d (see formula (1.15)). More precisely, for mean-field processes started in initial states $\bar{X}_0 > 0$, the quantity x_{upp} describes the double limit

$$\lim_{t \rightarrow \infty} \lim_{N \rightarrow \infty} \bar{X}_t = x_{\text{upp}}. \quad (3.6)$$

We see from (3.5) that the mean-field Ising model (as formulated in (3.1)) exhibits a second-order (i.e., continuous) phase transition at the critical point $\beta_c = 2$. Since

$$x_{\text{upp}}(\beta) \propto (\beta - \beta_c)^{1/2} \quad \text{as } \beta \downarrow \beta_c,$$

the *mean-field critical exponent* associated with the magnetization¹ is $c = 1/2$, which is the same as for the Ising model on \mathbb{Z}^d in dimensions $d \geq 4$ (see Section 1.5). Understanding why the mean-field model correctly predicts the critical exponent in sufficiently high dimensions goes beyond the scope of the present chapter.

To conclude the present section, we note that the two limits in (3.6) cannot be interchanged. Indeed, for each fixed N , the Markov process \bar{X} is irreducible, and hence, by standard theory, has a unique equilibrium law that is the long-time of the law at time t , started from an arbitrary initial state. In view of the symmetry of the problem, the magnetization in equilibrium must be zero, so regardless of the initial state, we have, for each fixed N ,

$$\lim_{t \rightarrow \infty} \mathbb{E}[\bar{X}_t] = 0.$$

¹In general, for a given second-order phase transition, there are several quantities of interest that all show power-law behavior near the critical point, and hence there are also several critical exponents associated with a given phase transition.

The reason why this can be true while at the same time (3.6) also holds is that the speed of convergence to equilibrium of the Markov process \bar{X} becomes very slow as $N \rightarrow \infty$.




Figure 3.4: Metastable behavior of a mean-field Ising model with $N = 50$ and $\beta = 3$. Note the different time scale compared to Figure 3.1.

In Figure 3.4, we have plotted the time evolution of a mean-field Ising model \bar{X} on a lattice with $N = 50$ sites, for a value of β above the critical point (concretely $\beta = 3$, which lies above $\beta_c = 2$). Although the average of \bar{X} in the long run is 0, we see that the process spends most of its time around the values x_{upp} and $-x_{\text{upp}}$, with rare transitions between the two. This sort of behavior is called *metastable behavior*.

The value $N = 50$ was near the highest possible value for which I could still numerically observe this sort of behavior. For $N = 100$ the transitions between the two metastable states x_{upp} and $-x_{\text{upp}}$ become so rare that my program was no longer able to see them within a reasonable runtime. With the help of *large deviations theory*, one can show that the time that the system spends in one metastable state is approximately exponentially distributed (with a large mean), and calculate the asymptotics of the mean waiting time as $N \rightarrow \infty$. It turns out that the mean time one has to wait for a transition grows exponentially fast in N .

3.4 Functions of Markov processes

In the present section we formulate a proposition and a theorem that we have already implicitly used. Both are concerned with functions of Markov processes. Let $X = (X_t)_{t \geq 0}$ be a Markov process with finite state space S , generator G , and semigroup $(P_t)_{t \geq 0}$. Let T be another finite set and let $f : S \rightarrow T$ be a function. For each $x \in S$ and $y' \in T$ such that $f(x) \neq y'$, let

$$\mathcal{H}(x, y') := \sum_{x' \in S: f(x')=y'} G(x, x') \quad (3.7)$$

be the total rate at which $f(X_t)$ jumps to the state y' , when the present state is $X_t = x$. The next proposition says that if these rates are a function of $f(x)$ only, then the process $Y = (Y_t)_{t \geq 0}$ defined by

$$Y_t := f(X_t) \quad (t \geq 0) \quad (3.8)$$

is itself a Markov process.

Proposition 3.1 (Autonomous Markov process) *Assume that the rates in (3.7) are of the form*

$$\mathcal{H}(x, y') = H(f(x), y') \quad (x \in S, y' \in T, f(x) \neq y') \quad (3.9)$$

where H is a Markov generator of some process in T . Then the process Y defined in (3.8) is a Markov process with generator H . Conversely, if for each initial law of the process X , it is true that Y is a Markov process with generator H , then (3.9) must hold.

Proof of Proposition 3.1 Let us define $\mathcal{H}(x, y')$ as in (3.7) also when $f(x) = y'$. We start by noting that if (3.9) holds for all $x \in S$ and $y' \in T$ such that $f(x) \neq y'$, then it also holds when $f(x) = y'$. To see this, we write

$$\begin{aligned} H(f(x), f(x)) &= - \sum_{y': y' \neq f(x)} H(f(x), y') = - \sum_{y': y' \neq f(x)} \sum_{x': f(x')=y'} G(x, x') \\ &= - \sum_{x': f(x') \neq f(x)} G(x, x') = \sum_{x': f(x')=f(x)} G(x, x'), \end{aligned}$$

where we have used that since H and G are Markov generators, one has $\sum_{y' \in T} H(f(x), y') = 0$ and $\sum_{x' \in S} G(x, x') = 0$. We have thus shown that (3.9) is equivalent to

$$H(f(x), y') = \sum_{x': f(x')=y'} G(x, x') \quad (x \in S, y' \in T). \quad (3.10)$$

We claim that this is equivalent to

$$Q_t(f(x), y') = \sum_{x': f(x')=y'} P_t(x, x') \quad (t \geq 0, x \in S, y' \in T), \quad (3.11)$$

where $(Q_t)_{t \geq 0}$ is the semigroup generated by H . To prove this, we start by observing that for any function $g : T \rightarrow \mathbb{R}$,

$$\begin{aligned} G(g \circ f)(x) &= \sum_{x'} G(x, x') g(f(x')) = \sum_{y'} \sum_{x': f(x')=y'} G(x, x') g(y'), \\ (Hg) \circ f(x) &= \sum_{y'} H(f(x), y') g(y'). \end{aligned}$$

The right-hand sides of these equations are equal for all $g : T \rightarrow \mathbb{R}$ if and only if (3.10) holds, so (3.10) is equivalent to the statement that

$$G(g \circ f) = (Hg) \circ f \quad (t \geq 0, g : T \rightarrow \mathbb{R}). \quad (3.12)$$

By exactly the same argument with G replaced by P_t and H replaced by Q_t , we see that (3.11) is equivalent to

$$P_t(g \circ f) = (Q_t g) \circ f \quad (t \geq 0, g : T \rightarrow \mathbb{R}). \quad (3.13)$$

To see that (3.12) and (3.13) are equivalent, we write

$$P_t = e^{Gt} = \sum_{n=0}^{\infty} \frac{1}{n!} t^n G^n \quad \text{and} \quad Q_t = e^{Ht} = \sum_{n=0}^{\infty} \frac{1}{n!} t^n H^n. \quad (3.14)$$

We observe that (3.12) implies

$$G^2(g \circ f) = G((Hg) \circ f) = (H^2 g) \circ f,$$

and similarly, by induction, $G^n(g \circ f) = (H^n g) \circ f$ for all $n \geq 0$, which by (3.14) implies (3.13). Conversely, if (3.13) holds for all $t \geq 0$, then it must hold up to first order in t as $t \downarrow 0$, which implies (3.12). This completes the proof that (3.9) is equivalent to (3.11).

If (3.11) holds, then by (2.6), the finite dimensional distributions of Y are given by

$$\begin{aligned} &\mathbb{P}[Y_0 = y_0, \dots, Y_{t_n} = y_n] \\ &= \sum_{x_0: f(x_0)=y_0} \cdots \sum_{x_n: f(x_n)=y_n} \mathbb{P}[X_0 = x_0] P_{t_1-t_0}(x_0, x_1) \cdots P_{t_n-t_{n-1}}(x_{n-1}, x_n) \\ &= \mathbb{P}[Y_0 = y_0] Q_{t_1-t_0}(y_0, y_1) \cdots Q_{t_n-t_{n-1}}(y_{n-1}, y_n) \end{aligned} \quad (3.15)$$

$(0 = t_0 < \dots < t_n)$. Again by (2.6), this implies that Y is a Markov process with generator H .

Conversely, if Y is a Markov process with generator H for each initial state of X , then for each $x_0 \in S$, (3.15) must hold when $X_0 = x_0$ a.s. and for $n = 1$, from which we see that (3.11) and hence (3.9) hold. ■

Summarizing, Proposition 3.1 says that if $Y_t = f(X_t)$ is a function of a Markov process, and the jump rates of Y are a function of the present state of Y only (and do not otherwise depend on the state of X), then Y is itself a Markov process. In such a situation, we will say that Y is an *autonomous* Markov process. We have already implicitly used Proposition 3.1 in Section 3.2, when we claimed that the process \bar{X} is a Markov process with jump rates as in (3.2).

Remark For the final statement of the proposition, it is essential that Y is a Markov process for *each* initial law X . There exist interesting examples of functions of Markov processes that are not autonomous Markov processes, but nonetheless are Markov processes for some *special* initial laws of the original Markov process. This is closely related to the concept of intertwining of Markov processes that will briefly be mentioned in Section 6.8 below.

Our next aim is to make the claim rigorous that for large N , the process \bar{X} can be approximated by solutions to the differential equation (3.4). We will apply a theorem from [DN08]. Although the proof is not very complicated, it is a bit lengthy and would detract from our main objects of interest here, so we only show how the theorem below can be deduced from a theorem in [DN08]. That paper also treats the multi-dimensional case and gives explicit estimates on probabilities of the form (3.19) below.

For each $N \geq 1$, let $X^N = (X_t^N)_{t \geq 0}$ be a Markov process with finite state space S_N , generator G_N , and semigroup $(P_t^N)_{t \geq 0}$, and let $f_N : S_N \rightarrow \mathbb{R}$ be functions. We will be interested in conditions under which the processes $(f_N(X_t^N))_{t \geq 0}$ approximate the solution $(y_t)_{t \geq 0}$ of a differential equation, in the limit $N \rightarrow \infty$. Note that we do not require that $f_N(X_t^N)$ is an autonomous Markov process. To ease notation, we will sometimes drop the super- and subscripts N when no confusion arises.

We define two functions $\alpha = \alpha_N$ and $\beta = \beta_N$ that describe the quadratic variation and drift, respectively, of the process $f(X_t)$. More precisely, these functions are given by

$$\begin{aligned}\alpha(x) &:= \sum_{x' \in S} G(x, x') (f(x') - f(x))^2, \\ \beta(x) &:= \sum_{x' \in S} G(x, x') (f(x') - f(x)).\end{aligned}$$

The idea is that if α tends to zero and β approximates a nice, Lipschitz continuous function of $f(X_t)$, then $f(X_t)$ should in the limit be given by the solution of a differential equation.

We assume that the functions f_N all take values in a closed interval $I \subset \mathbb{R}$ with left and right boundaries $I_- := \inf I$ and $I_+ := \sup I$, which may be finite or infinite. We also assume that there exists a globally Lipschitz function $b : I \rightarrow \mathbb{R}$ such that

$$\sup_{x \in S_N} |\beta_N(x) - b(f_N(x))| \xrightarrow{N \rightarrow \infty} 0, \quad (3.16)$$

i.e., the drift function β is uniformly approximated by $b \circ f_N$. Assuming also that

$$b(I_-) \geq 0 \quad \text{if } I_- > -\infty \quad \text{and} \quad b(I_+) \leq 0 \quad \text{if } I_+ < \infty, \quad (3.17)$$

the differential equation

$$\frac{\partial}{\partial t} y_t = b(y_t) \quad (t \geq 0)$$

has a unique I -valued solution $(y_t)_{t \geq 0}$ for each initial state $y_0 \in I$. The following theorem gives sufficient conditions for the I -valued processes $(f_N(X_t^N))_{t \geq 0}$ to approximate a solution of the differential equation.

Theorem 3.2 (Limiting differential equation) *Assume that $f_N(X_0^N)$ converges in probability to y_0 and that as well as (3.16), one moreover has*

$$\sup_{x \in S_N} \alpha_N(x) \xrightarrow{N \rightarrow \infty} 0. \quad (3.18)$$

Then, for each $T < \infty$ and $\varepsilon > 0$,

$$\mathbb{P}[|f_N(X_t^N) - y_t| \leq \varepsilon \quad \forall t \in [0, T]] \xrightarrow{N \rightarrow \infty} 1. \quad (3.19)$$

Proof We apply [DN08, Thm 4.1]. Fix $T < \infty$ and $\varepsilon > 0$ and also fix $y_0 \in I$. Let L denote the Lipschitz constant of b . The assumptions of [DN08, Thm 4.1] allow for the case that f_N does not in general take values in I , but only under the additional condition that $f_N(x)$ is not further than ε from a possible value the solution of the differential equation. In our case, these more general assumptions are automatically satisfied. Set $\delta := \frac{1}{3}\varepsilon e^{-LT}$. We consider the events

$$\Omega_0 := \{|f(X_0) - y_0| \leq \delta\} \quad \text{and} \quad \Omega_1 := \left\{ \int_0^T |\beta(X_t) - b(f(X_t))| dt \leq \delta \right\}.$$

For $K > 0$, we also define

$$\Omega_{K,2} := \left\{ \int_0^T \alpha(X_t) dt \leq KT \right\}.$$

Then [DN08, Thm 4.1] tells us that

$$\mathbb{P} \left[\sup_{t \in [0, T]} |f(X_t) - y_t| > \varepsilon \right] \leq 4KT\delta^{-2} + \mathbb{P}(\Omega_0^c \cup \Omega_1^c \cup \Omega_{K,2}^c). \quad (3.20)$$

Our assumption that $f_N(X_0^N) \rightarrow y_0$ in probability implies that $\mathbb{P}(\Omega_0^c) \rightarrow 0$ as $N \rightarrow \infty$. Set

$$A_N := \sup_{x \in S_N} \alpha_N(x) \quad \text{and} \quad B_N := \sup_{x \in S_N} |\beta_N(x) - b(f_N(x))|$$

Then $A_N \rightarrow 0$ by (3.18) and $B_N \rightarrow 0$ by (3.16). Since

$$\int_0^T |\beta(X_t) - b(f(X_t))| dt \leq B_N T \leq \delta$$

for N sufficiently large, we see that $\mathbb{P}(\Omega_1^c) = 0$ for N sufficiently large. Also, since

$$\int_0^T \alpha(X_t) dt \leq A_N T,$$

we see that $\mathbb{P}(\Omega_{A_N,2}^c) = 0$ for all N . Inserting $K = A_N$ in (3.20), we see that the right-hand side tends to zero as $N \rightarrow \infty$. ■

Using Theorem 3.2, we can make the approximation of the mean-field Ising model by the differential equation (3.4) rigorous. Let $X^N = (X_t^N)_{t \geq 0}$ denote the Markov process with state space $\{-1, +1\}^{\Lambda_N}$, where Λ_N is a set containing N elements and the jump rates of X^N are given in (3.1). By Proposition 3.1, the process $\bar{X}_t^N := \frac{1}{N} \sum_{i \in \Lambda_N} X_t(i)$ is itself a Markov process with jump rates as in (3.2). We can either apply Theorem 3.2 directly to the Markov processes X^N and the functions $f_N(x) := \frac{1}{N} \sum_{i \in \Lambda_N} x(i)$, or we can apply Theorem 3.2 to the Markov processes \bar{X}^N and choose for f_N the identity function $f_N(\bar{x}) = \bar{x}$. In either case, the assumption (3.16) is already verified in (3.3). To check also (3.18), we calculate

$$\alpha_N(x) = r_+(\bar{x}) \left(\frac{2}{N} \right)^2 + r_-(\bar{x}) \left(\frac{2}{N} \right)^2 = \frac{2}{N} \left(1 + \bar{x} \frac{e^{-\beta\bar{x}/2} - e^{\beta\bar{x}/2}}{e^{-\beta\bar{x}/2} + e^{\beta\bar{x}/2}} \right),$$

which clearly tends uniformly to zero as $N \rightarrow \infty$.

3.5 The mean-field contact process

Recall the definition of the generator of the contact process from (1.8). We slightly reformulate this as

$$G_{\text{cont}}f(x) := \lambda \sum_{i \in \mathbb{Z}^d} \frac{1}{|\mathcal{N}_i|} \sum_{j \in \mathcal{N}_i} \{f(\text{bra}_{ij}(x)) - f(x)\} \\ + \sum_{i \in \mathbb{Z}^d} \{f(\text{death}_i(x)) - f(x)\} \quad (x \in \{0, 1\}^\Lambda), \quad (3.21)$$

where as customary we have set the death rate to $\delta = 1$, and we have also reparametrized the infection rate so that λ denotes the total rate of all outgoing infections from a given site, instead of the infection rate per neighbor.

We will be interested in the contact process on the complete graph, which means that we take for $\Lambda = \Lambda_N$ a set with N elements, which we equip with the structure of a complete graph with (undirected) edge set $E = E_N := \{\{i, j\} : i, j \in \Lambda_N\}$ and corresponding set of oriented edges $\mathcal{E} = \mathcal{E}_N$. We will be interested in the fraction of infected sites

$$\bar{X}_t = \bar{X}_t^N := \frac{1}{N} \sum_{i \in \Lambda_N} X_t(i) \quad (t \geq 0),$$

which jumps with the following rates

$$\begin{aligned} \bar{x} &\mapsto \bar{x} + \frac{1}{N} && \text{with rate} && r_+(\bar{x}) := \lambda N \bar{x} (1 - \bar{x}), \\ \bar{x} &\mapsto \bar{x} - \frac{1}{N} && \text{with rate} && r_-(\bar{x}) := N \bar{x}. \end{aligned} \quad (3.22)$$

Here $N(1 - \bar{x})$ is the number of healthy sites, each of which gets infected with rate $\lambda \bar{x}$, and $N \bar{x}$ is the number of infected sites, each of which recovers with rate one. Note that since these rates are a function of \bar{x} only, by Proposition 3.1, the process $(\bar{X}_t)_{t \geq 0}$ is an autonomous Markov chain.

We wish to apply Theorem 3.2 to conclude that \bar{X} can, for large N be approximated by the solution of a differential equation. To this aim, we calculate the drift β and quadratic variation function α .

$$\begin{aligned} \alpha_N(x) &= r_+(\bar{x}) \frac{1}{N^2} + r_-(\bar{x}) \frac{1}{N^2} = \frac{1}{N} (\lambda \bar{x} (1 - \bar{x}) + \bar{x}), \\ \beta_N(x) &= r_+(\bar{x}) \frac{1}{N} - r_-(\bar{x}) \frac{1}{N} = \lambda \bar{x} (1 - \bar{x}) - \bar{x}. \end{aligned}$$


By Theorem 3.2, it follows that in the mean-field limit $N \rightarrow \infty$, the fraction of infected sites can be approximated by solutions of the differential equation

$$\frac{\partial}{\partial t} \bar{X}_t = b_\lambda(\bar{X}_t) \quad (t \geq 0), \quad \text{where} \quad b_\lambda(\bar{x}) := \lambda \bar{x} (1 - \bar{x}) - \bar{x}. \quad (3.23)$$

The equation $b_\lambda(\bar{x}) = 0$ has the solutions

$$\bar{x} = 0 \quad \text{and} \quad \bar{x} = 1 - \lambda^{-1}.$$

The second solution lies inside the interval $[0, 1]$ of possible values of \bar{X}_t if and only if $\lambda \geq 1$. Plotting the function b_λ for $\lambda < 1$ and $\lambda > 1$ yields the following pictures.



We see from this that the fixed point $\bar{x} = 0$ is stable for $\lambda \leq 1$ but becomes unstable for $\lambda > 1$, in which case $\bar{x} = 1 - \lambda^{-1}$ is the only stable fixed point that attracts all solutions started in a nonzero initial state. The situation is summarized in Figure 3.5.




Figure 3.5: Mean-field analysis of the contact process.

Letting $\bar{x}_{\text{upp}}(\lambda) := 0 \vee (1 - \lambda^{-1})$ denote the stable fixed point, we see that the mean-field contact process exhibits a second-order phase transition at the critical point $\lambda_c = 1$. Since

$$\bar{x}_{\text{upp}}(\lambda) \propto (\lambda - \lambda_c) \quad \text{as } \lambda \downarrow \lambda_c,$$

the associated critical point is $c = 1$, in line with what we know for contact processes in dimensions $d \geq 4$ (see the discussion in Section 1.5).

3.6 The mean-field voter model

Recall the definition of the generator of the voter model from (1.5). For simplicity, we will only consider the two-type model and as the local state space we will choose $S = \{0, 1\}$. Specializing to the complete graph $\Lambda = \Lambda_N$ with N vertices, the generator becomes

$$G_{\text{vot}}f(x) = \frac{1}{|\Lambda|} \sum_{(i,j) \in \mathcal{E}} \{f((\text{vot}_{ij}(x))) - f(x)\} \quad (x \in \{0, 1\}^\Lambda). \quad (3.24)$$

Note that the factor $|\Lambda|^{-1}$ says that each site i updates its type with rate one, and at such an event chooses a new type from a uniformly chosen site j (allowing for the case $i = j$, which has no effect).

We are interested in the fraction of sites of type 1,

$$\bar{X}_t = \bar{X}_t^N := \frac{1}{N} \sum_{i \in \Lambda_N} X_t(i) \quad (t \geq 0),$$

which jumps as (compare (3.22))

$$\begin{aligned} \bar{x} &\mapsto \bar{x} + \frac{1}{N} && \text{with rate} && r_+(\bar{x}) := N\bar{x}(1 - \bar{x}), \\ \bar{x} &\mapsto \bar{x} - \frac{1}{N} && \text{with rate} && r_-(\bar{x}) := N\bar{x}(1 - \bar{x}). \end{aligned}$$

Note that $N(1 - \bar{x})$ is the number of sites of type 0, and that each such site adopts the type 1 with rate \bar{x} . The derivation of $r_-(\bar{x})$ is similar. We calculate the drift β and quadratic variation function α .

$$\begin{aligned} \alpha_N(x) &= r_+(\bar{x}) \frac{1}{N^2} + r_-(\bar{x}) \frac{1}{N^2} = \frac{2}{N} \bar{x}(1 - \bar{x}), \\ \beta_N(x) &= r_+(\bar{x}) \frac{1}{N} - r_-(\bar{x}) \frac{1}{N} = 0. \end{aligned}$$

Applying Theorem 3.2, we see that in the limit $N \rightarrow \infty$, the process $(\bar{X}_t)_{t \geq 0}$ is well approximated by solutions to the differential equation

$$\frac{\partial}{\partial t} \bar{X}_t = 0 \quad (t \geq 0),$$

i.e., \bar{X}_t is approximately constant as a function of t .

Of course, if we go to larger time scales, then \bar{X}_t will no longer be constant; compare Figure 3.4. In fact, we can determine the time scale at which \bar{X}_t fluctuates quite precisely. Scaling up time by a factor $|\Lambda| = N$ is the same as multiplying all rates by a factor $|\Lambda|$. If we repeat our previous calculations for the process with generator

$$G_{\text{vot}}f(x) = \sum_{(i,j) \in \mathcal{E}} \{f((\text{vot}_{ij}(x))) - f(x)\} \quad (x \in \{0, 1\}^\Lambda), \quad (3.25)$$




Figure 3.6: The fraction of type 1 individuals in the mean-field voter model from (3.25) on $N = 100$ sites. This process approximates the Wright-Fisher diffusion.

then the drift and quadratic variation are given by

$$\alpha_N(x) = 2\bar{x}(1 - \bar{x}),$$

$$\beta_N(x) = 0.$$

In this case, the quadratic variation does not go to zero, so Theorem 3.2 is no longer applicable. One can show, however, that in the limit $N \rightarrow \infty$ the new, sped-up process is well approximated by solutions to the (Itô) stochastic differential equation (SDE)

$$d\bar{X}_t = \sqrt{2\bar{X}_t(1 - \bar{X}_t)} dB_t \quad (t \geq 0),$$

where $2\bar{X}_t(1 - \bar{X}_t) = \alpha(X_t)$ is of course the quadratic variation function we have just calculated. Solutions to this SDE are *Wright-Fisher diffusions*, i.e., Markov processes with continuous sample paths and generator

$$Gf(\bar{x}) = \bar{x}(1 - \bar{x}) \frac{\partial^2}{\partial \bar{x}^2} f(\bar{x}). \quad (3.26)$$

These calculations can be made rigorous using methods from the theory of convergence of Markov processes; see, e.g., the book [EK86]. See Figure 3.6 for a simulation of the process \bar{X} when X has the generator in (3.25) and $N = 100$.

3.7 Exercises

Exercise 3.3 Do a mean-field analysis of the process with generator

$$Gf(x) = b|\Lambda|^{-2} \sum_{ii'j} \{f(\text{coop}_{ii'j}x) - f(x)\} \\ + \sum_i \{f(\text{death}_i x) - f(x)\},$$

where the maps $\text{coop}_{ii'j}$ and death_i are defined in (1.25) and (1.7), respectively. Do you observe a phase transition? Is it first- or second order? Hint: Figure 3.7.




Figure 3.7: Mean-field analysis of a model with cooperative branching and deaths.

Exercise 3.4 Same as above for the model with generator

$$Gf(x) = b|\Lambda|^{-2} \sum_{ii'j} \{f(\text{coop}_{ii'j}x) - f(x)\} \\ + |\Lambda|^{-1} \sum_{ij} \{f(\text{rw}_{ij}x) - f(x)\}.$$

Exercise 3.5 Derive an SDE in the limit $|\Lambda| \rightarrow \infty$ for the density of the mean-field voter model with small bias and death rates, with generator

$$Gf(x) = \sum_{ij \in \Lambda} \{f(\text{vot}_{ij}x) - f(x)\} \\ + s|\Lambda|^{-1} \sum_{ij \in \Lambda} \{f(\text{bra}_{ij}x) - f(x)\} \\ + d \sum_{i \in \Lambda} \{f(\text{death}_i x) - f(x)\}.$$

Hint: You should find expressions of the form

$$\begin{aligned}\mathbb{E}^{\bar{x}}[(\bar{X}_t - \bar{x})] &= b(\bar{x}) \cdot t + O(t^2), \\ \mathbb{E}^{\bar{x}}[(\bar{X}_t - \bar{x})^2] &= a(\bar{x}) \cdot t + O(t^2),\end{aligned}$$

which leads to a limiting generator of the form

$$Gf(\bar{x}) = \frac{1}{2}a(\bar{x})\frac{\partial^2}{\partial \bar{x}^2}f(\bar{x}) + b(\bar{x})\frac{\partial}{\partial \bar{x}}f(\bar{x}).$$

Exercise 3.6 Do a mean-field analysis of the following extension of the voter model, introduced in [NP99]. In this model, the site i flips

$$\begin{aligned}0 &\mapsto 1 \quad \text{with rate} \quad (f_0 + \alpha_{01}f_1)f_1, \\ 1 &\mapsto 0 \quad \text{with rate} \quad (f_1 + \alpha_{10}f_0)f_0,\end{aligned}$$

where $\alpha_{01}, \alpha_{10} > 0$ and $f_\tau = |\mathcal{N}_i|^{-1} \sum_{j \in \mathcal{N}_i} 1_{\{x(j)=\tau\}}$ is the relative frequency of type τ in the neighborhood of i .

Find all stable and unstable fixed points of the mean-field model in the regimes: I. $\alpha_{01}, \alpha_{10} < 1$, II. $\alpha_{01} < 1 < \alpha_{10}$, III. $\alpha_{10} < 1 < \alpha_{01}$, IV. $1 < \alpha_{01}, \alpha_{10}$.

Chapter 4

Construction and ergodicity

4.1 Introduction

As explained in Chapter 1, interacting particle systems are Markov processes with a state space of the form S^Λ where S is a finite set, called the *local state space*, and Λ is a countable set, called the *lattice*. The general form of a generator of an interacting particle system is

$$Gf = \sum_{K \in \mathcal{K}} r_K \{Kf - f\}, \quad (4.1)$$

where \mathcal{K} is a countable collection of local probability kernels on S^Λ and $(r_K)_{K \in \mathcal{K}}$ are nonnegative constants. The precise definition of a local probability kernel will be given in Section 4.3. Roughly speaking, these are kernels that change the values at finitely many sites in a way that does not depend too much on the values of sites that are far away. Usually, it is possible to take all kernels $K \in \mathcal{K}$ deterministic, in which case the generator takes the form

$$Gf(x) = \sum_{m \in \mathcal{G}} r_m \{f(m(x)) - f(x)\} \quad (x \in S^\Lambda), \quad (4.2)$$

where \mathcal{G} is a set whose elements are *local maps* $m : S^\Lambda \rightarrow S^\Lambda$ and $(r_m)_{m \in \mathcal{G}}$ is a collection of nonnegative rates. If Λ is finite, then S^Λ is also a finite set and we can use Proposition 2.7 to construct a Markov process $X = (X_t)_{t \geq 0}$ with generator G in terms of a Poisson process ω .

On the other hand, if Λ is countable but infinite, then the space S^Λ is not finite, and, in fact, not even countable. Indeed, as is well-known, $\{0, 1\}^\mathbb{N}$ has the same cardinality as the real numbers. As a result, the construction of interacting particle systems on infinite lattices is considerably more involved than in the finite case. Nevertheless, we will see that they can be constructed

using more or less the same approach as in Proposition 2.7. The only complication is that the total rate of all local maps is usually infinite, so that it is no longer possible to order the elements of the Poisson set ω according to the time when they occur. However, since each map is *local*, and since in finite time intervals only finitely many local maps can influence the local state at any given site i , we will see that under certain summability assumptions, the Poisson construction still yields a well-defined process.

In practice, one usually needs not only the Poisson construction of an interacting particle system, but also wishes to show that the process is uniquely characterized by its generator. One reason is that, as we have already seen in Section 2.4, sometimes the same process can be constructed using different Poisson constructions, and one wants to prove that these constructions are indeed equivalent.

To give a generator construction of interacting particle systems, we will apply the theory of Feller processes. We start by equipping S^Λ with the *product topology*, which says that a sequence $x_n \in S^\Lambda$ converges to a limit x if and only if

$$x_n(i) \xrightarrow[n \rightarrow \infty]{} x(i) \quad \forall i \in \Lambda.$$

Note that since S is finite, this says simply that for each $i \in \Lambda$, there is an N (which may depend on i) such that $x_n(i) = x(i)$ for all $n \geq N$. Since S is finite, it is in particular compact, so by Tychonoff's theorem, the space S^Λ is compact in the product topology. The product topology is metrizable. For example, if $(a_i)_{i \in \Lambda}$ are strictly positive constants such that $\sum_{i \in \Lambda} a_i < \infty$, then

$$d(x, y) := \sum_{i \in \Lambda} a_i 1_{\{x(i) \neq y(i)\}}$$

defines a metric that generates the product topology.

In Section 4.2, we will collect some general facts about Feller processes, which are a class of Markov processes with compact, metrizable state spaces, that are uniquely characterized by their generators. Since this is rather functional theoretic material, which is moreover well-known, we will state the main facts without proof, but give references to places where proofs can be found.

In Section 4.3, we then give the Poisson construction of interacting particle systems (including proofs). In Section 4.4, we show that our construction yields a Feller process and determine its generator.

Luckily, all this abstract theory gives us more than just the information that the systems we are interested in are well defined. In Section 4.5, we will see that as a side-result of our proofs, we can derive sufficient conditions for an interacting particle system to be ergodic, i.e., to have a unique invariant

law that is the long-time limit starting from any initial state. We will apply this to derive lower bounds on the critical points of the Ising model.

4.2 Feller processes

In Section 2.2, we gave a summary of the basic theory of continuous-time Markov processes with finite state space S . In the present section, we will see that with a bit of care, much of this theory can be generalized in a rather elegant way to Markov processes taking values in a compact metrizable state space. The basic assumption we will make is that the transition probabilities $(P_t)_{t \geq 0}$ are continuous, which means that we will be discussing *Feller processes*.

Let E be a compact metrizable space.¹ We use the notation

- $\mathcal{B}(E) :=$ the Borel- σ -field on E ,
- $B(E) :=$ the space of bounded, Borel-measurable functions $f : E \rightarrow \mathbb{R}$,
- $\mathcal{C}(E) :=$ the space of continuous functions $f : E \rightarrow \mathbb{R}$,
- $\mathcal{M}_1(E) :=$ the space of probability measures μ on E .

We equip $\mathcal{C}(E)$ with the supremumnorm

$$\|f\|_\infty := \sup_{x \in E} |f(x)| \quad (f \in \mathcal{C}(E)),$$

making $\mathcal{C}(E)$ into a Banach space. We equip $\mathcal{M}_1(E)$ with the topology of weak convergence, where by definition,² μ_n converges weakly to μ , denoted $\mu_n \Rightarrow \mu$, if $\int f d\mu_n \rightarrow \int f d\mu$ for all $f \in \mathcal{C}(E)$. With this topology, $\mathcal{M}_1(E)$ is a compact metrizable space.

A *probability kernel* on E is a function $K : E \times \mathcal{B}(E) \rightarrow \mathbb{R}$ such that

- (i) $K(x, \cdot)$ is a probability measure on E for each $x \in E$,
- (ii) $K(\cdot, A)$ is a real measurable function on E for each $A \in \mathcal{B}(E)$.

This is equivalent to the statement that $x \mapsto K(x, \cdot)$ is a measurable map from E to $\mathcal{M}_1(E)$ (where the latter is equipped with the topology of weak convergence and the associated Borel- σ -field). By definition, a probability kernel is *continuous* if the map $x \mapsto K(x, \cdot)$ is continuous (with respect

¹Such spaces are always separable and complete in any metric that generates the topology; in particular, they are Polish spaces.

²More precisely, the topology of weak convergence is the unique *metrizable* topology with this property. Since in metrizable spaces, convergent subsequences uniquely characterize the topology, such a definition is unambiguous.

to the topologies with which we have equipped these spaces). A probability kernel is *deterministic* if it is of the form $K(x, \cdot) = \delta_{m(x)}$ for some measurable map $m : E \rightarrow E$, where $\delta_{m(x)}$ denotes the delta-measure at $m(x)$. It is easy to see that a deterministic kernel is continuous if and only if m is a continuous map. A *random mapping representation* of a probability kernel K is a random measurable map³ $M : E \rightarrow E$ such that $K(x, \cdot) = \mathbb{P}[M(x) \in \cdot] \ (x \in E)$.⁴

If $K(x, dy)$ is a probability kernel on a Polish space E , then setting

$$Kf(x) := \int_E K(x, dy)f(y) \quad (x \in E, f \in B(E))$$

defines a linear operator $K : B(E) \rightarrow B(E)$. We define the *composition* of two probability kernels K, L as

$$(KL)(x, A) := \int_E K(x, dy)L(y, A) \quad (x \in E, A \in \mathcal{B}(E)).$$

Then KL is again a probability kernel on E and the linear operator $(KL) : B(E) \rightarrow B(E)$ associated with this kernel is the composition of the linear operators K and L . It follows from the definition of weak convergence that a kernel K is continuous if and only if its associated linear operator maps the space $\mathcal{C}(E)$ into itself. If μ is a probability measure and K is a probability kernel, then

$$(\mu K)(A) := \int \mu(dx)K(x, A) \quad (A \in \mathcal{B}(E))$$

defines another probability measure μK . Introducing the notation $\mu f := \int f d\mu$, one has $(\mu K)f = \mu(Kf)$ for all $f \in B(E)$.

By definition, a *continuous transition probability* on E is a collection $(P_t)_{t \geq 0}$ of probability kernels on E , such that

- (i) $(x, t) \mapsto P_t(x, \cdot)$ is a continuous map from $E \times [0, \infty)$ into $\mathcal{M}_1(E)$,
- (ii) $P_0 = 1$ and $P_s P_t = P_{s+t} \ (s, t \geq 0)$.

In particular, (i) implies that each P_t is a continuous probability kernel, so

³More formally, this means that $M : \Omega \times E \rightarrow E$ is measurable with respect to the product- σ -field $\mathcal{F} \otimes \mathcal{B}(E)$, where $(\Omega, \mathcal{F}, \mathbb{P})$ is the underlying probability space.

⁴For infinite spaces, it is not so clear if every probability kernel has a random mapping representation. One could also ask if every continuous probability kernel has a representation in terms of continuous maps. Although these questions are interesting, we will neglect them here.

each P_t maps the space $\mathcal{C}(E)$ into itself. One has

- (i) $\lim_{t \rightarrow 0} P_t f = P_0 f = f \quad (f \in \mathcal{C}(E)),$
- (ii) $P_s P_t f = P_{s+t} f \quad (s, t \geq 0),$
- (iii) $f \geq 0$ implies $P_t f \geq 0,$
- (iv) $P_t 1 = 1,$

and conversely, each collection of linear operators $P_t : \mathcal{C}(E) \rightarrow \mathcal{C}(E)$ with these properties corresponds to a unique continuous transition probability on E . Such a collection of linear operators $P_t : \mathcal{C}(E) \rightarrow \mathcal{C}(E)$ is called a *Feller semigroup*. We note that in (i), the limit is (of course) with respect to the topology we have chosen on $\mathcal{C}(E)$, i.e., with respect to the supremum norm.

By definition, a function $w : [0, \infty) \rightarrow E$ is *cadlag* if it is right-continuous with left limits,⁵ i.e.,

- (i) $\lim_{t \downarrow s} w_t = w_s \quad (s \geq 0),$
- (ii) $\lim_{t \uparrow s} w_t =: w_{s-}$ exists $(s > 0).$

Let $(P_t)_{t \geq 0}$ be a Feller semigroup. By definition a *Feller process* with semigroup $(P_t)_{t \geq 0}$ is a stochastic process $X = (X_t)_{t \geq 0}$ with cadlag sample paths⁶ such that

$$\mathbb{P}[X_u \in \cdot \mid (X_s)_{0 \leq s \leq t}] = P_{u-t}(X_t, \cdot) \quad \text{a.s.} \quad (0 \leq t \leq u). \quad (4.3)$$

Here we condition on the σ -field generated by the random variables $(X_s)_{0 \leq s \leq t}$. Formula (4.3) is equivalent to the statement that the finite dimensional distributions of X are given by

$$\begin{aligned} \mathbb{P}[X_0 \in dx_0, \dots, X_{t_n} \in dx_n] \\ = \mathbb{P}[X_0 \in dx_0] P_{t_1-t_0}(x_0, dx_1) \cdots P_{t_n-t_{n-1}}(x_{n-1}, dx_n) \end{aligned} \quad (4.4)$$

$(0 < t_1 < \cdots < t_n)$. Formula (4.4) is symbolic notation, which means that

$$\begin{aligned} \mathbb{E}[f(X_0, \dots, X_{t_n})] \\ = \int \mathbb{P}[X_0 \in dx_0] \int P_{t_1-t_0}(x_0, dx_1) \cdots \int P_{t_n-t_{n-1}}(x_{n-1}, dx_n) f(x_0, \dots, x_n) \end{aligned}$$

⁵The word cadlag is an abbreviation of the French *continue à droite, limite à gauche*.

⁶It is possible to equip the space $\mathcal{D}_E[0, \infty)$ of cadlag functions $w : [0, \infty) \rightarrow E$ with a (rather natural) topology, called the *Skorohod topology*, such that $\mathcal{D}_E[0, \infty)$ is a Polish space and the Borel- σ -field on $\mathcal{D}_E[0, \infty)$ is generated by the coordinate projections $w \mapsto w_t$ ($t \geq 0$). As a result, we can view a stochastic process $X = (X_t)_{t \geq 0}$ with cadlag sample paths as a single random variable X taking values in the space $\mathcal{D}_E[0, \infty)$. The law of such a random variable is then uniquely determined by the finite dimensional distributions of $(X_t)_{t \geq 0}$.

for all $f \in B(E^{n+1})$. By (4.4), the law of a Feller process X is uniquely determined by its initial law $\mathbb{P}[X_0 \in \cdot]$ and its transition probabilities $(P_t)_{t \geq 0}$. Existence is less obvious than uniqueness, but the next theorem says that this holds in full generality.

Theorem 4.1 (Construction of Feller processes) *Let E be a compact metrizable space, let μ be a probability measure on E , and let $(P_t)_{t \geq 0}$ be a Feller semigroup. Then there exists a Feller process $X = (X_t)_{t \geq 0}$ with initial law $\mathbb{P}[X_0 \in \cdot] = \mu$, and such a process is unique in distribution.*

Just as in the case for finite state space, we would like to characterize a Feller semigroup by its generator. This is somewhat more complicated than in the finite setting since in general, it is not possible to make sense of the exponential formula $P_t = e^{tG} := \sum_{n=0}^{\infty} \frac{1}{n!} (tG)^n$. This is related to the fact that if G is the generator of a Feller semigroup, then in general it is not possible to define Gf for all $f \in \mathcal{C}(E)$, as we now explain.

Let \mathcal{V} be a Banach space. (In our case, the only Banach spaces that we will need are spaces of the form $\mathcal{C}(E)$, equipped with the supremum norm.) By definition, a *linear operator* on \mathcal{V} is a pair $(A, \mathcal{D}(A))$ where $\mathcal{D}(A)$ is a linear subspace of \mathcal{V} , called the *domain* and A is a linear map $A : \mathcal{D}(A) \rightarrow \mathcal{V}$. Even though a linear operator is really a pair $(A, \mathcal{D}(A))$, one often writes sentences such as “let A be a linear operator” without explicitly mentioning the domain. This is similar to phrases like: “let \mathcal{V} be a Banach space” (without mentioning the norm) or “let M be a measurable space” (without mentioning the σ -field).

We say that a linear operator A (with domain $\mathcal{D}(A)$) on a Banach space \mathcal{V} is *closed* if and only if its graph $\{(f, Af) : f \in \mathcal{D}(A)\}$ is a closed subset of $\mathcal{V} \times \mathcal{V}$. By definition, a linear operator A (with domain $\mathcal{D}(A)$) on a Banach space \mathcal{V} is *closable* if the closure of its graph (as a subset of $\mathcal{V} \times \mathcal{V}$) is the graph of a linear operator \bar{A} with domain $\mathcal{D}(\bar{A})$. This operator is then called the *closure* of A . We mention the following theorem.

Theorem 4.2 (Closed graph theorem) *Let \mathcal{V} be a Banach space and let A be a linear operator that is everywhere defined, i.e., $\mathcal{D}(A) = \mathcal{V}$. Then the following statements are equivalent.*

- (i) A is continuous as a map from \mathcal{V} into itself.
- (ii) A is bounded, i.e., there exists a constant $C < \infty$ such that $\|Af\| \leq C\|f\|$ ($f \in \mathcal{V}$).
- (iii) A is closed.

Theorem 4.2 shows in particular that if A is an unbounded operator (i.e., there exists $0 \neq f_n \in \mathcal{D}(A)$ such that $\|Af_n\|/\|f_n\| \rightarrow \infty$) and A is closable, then its closure \overline{A} will not be everywhere defined. Closed (but possibly unbounded) linear operators are in a sense “the next good thing” after bounded operators.

As before, let E be a compact metrizable space and let $(P_t)_{t \geq 0}$ be a continuous transition probability (or equivalently Feller semigroup) on E . By definition, the *generator* of $(P_t)_{t \geq 0}$ is the linear operator

$$Gf := \lim_{t \rightarrow 0} t^{-1}(P_t f - f),$$

with domain

$$\mathcal{D}(G) := \{f \in \mathcal{C}(E) : \text{the limit } \lim_{t \rightarrow 0} t^{-1}(P_t f - f) \text{ exists}\}.$$

Here, when we say that the limit exists, we mean (of course) with respect to the topology on $\mathcal{C}(E)$, i.e., w.r.t. the supremum norm. The following lemma says that generators are closed, densely defined operators.

Lemma 4.3 (Elementary properties of generators) *Let G be the generator of a Feller semigroup $(P_t)_{t \geq 0}$. Then G is closed and $\mathcal{D}(G)$ is a dense subspace of $\mathcal{C}(E)$.*

Since G is closed, Theorem 4.2 tells us that G is everywhere defined (i.e., $\mathcal{D}(G) = \mathcal{C}(G)$) if and only if G is bounded. For bounded generators, it is not hard to show that the exponential formula $e^{tG} := \sum_{n=0}^{\infty} \frac{1}{n!} (tG)^n$ converges in the norm on $\mathcal{C}(G)$ and that the Feller semigroup with generator G is given by $P_t = e^{tG}$. On the other hand, if G is unbounded, then it is in general not possible to make sense of the exponential formula.⁷ In the context of interacting particle systems, it is not hard to show that a generator of the form (4.2) is bounded if $\sum_{m \in \mathcal{G}} r_m < \infty$. For the particle systems we will be interested in, this sum will usually be infinite and the generator will be unbounded.

Since we cannot use the exponential formula $P_t = e^{tG}$, we need another way to characterize $(P_t)_{t \geq 0}$ in terms of G . Let A be a linear operator on $\mathcal{C}(E)$. By definition, we say that a function $[0, \infty) \ni t \mapsto u_t \in \mathcal{C}(E)$ solves the *Cauchy equation*

$$\frac{\partial}{\partial t} u_t = Au_t \quad (t \geq 0) \tag{4.5}$$

⁷In order for $\sum_{n=0}^{\infty} \frac{1}{n!} t^n G^n f$ to make sense, we need that $G^n f$ is well-defined for all $n \geq 0$. For $n = 1$ this already requires that $f \in \mathcal{D}(G)$ but for higher n we need even more since it is in general not true that G maps $\mathcal{D}(G)$ into itself. Thus, it is not even clear for which class of functions we can make sense of each term in the expansion separately.

if $u_t \in \mathcal{D}(A)$ for all $t \geq 0$, the maps $t \mapsto u_t$ and $t \mapsto Au_t$ are continuous (w.r.t. the topology on $\mathcal{C}(E)$), the limit $\frac{\partial}{\partial t}u_t := \lim_{s \rightarrow 0} s^{-1}(u_{t+s} - u_s)$ exists (w.r.t. the topology on $\mathcal{C}(E)$) for all $t \geq 0$, and (4.5) holds. The following proposition shows that a Feller semigroup is uniquely characterized by its generator.

Proposition 4.4 (Cauchy problem) *Let G be the generator of a Feller semigroup $(P_t)_{t \geq 0}$. Then, for each $f \in \mathcal{D}(G)$, the Cauchy equation $\frac{\partial}{\partial t}u_t = Gu_t$ ($t \geq 0$) has a unique solution $(u_t)_{t \geq 0}$ with initial state $u_0 = f$. Denoting this solution by $U_t f := u_t$ defines for each $t \geq 0$ a linear operator U_t with domain $\mathcal{D}(G)$, of which $P_t = \overline{U_t}$ is the closure.*

We need a way to check that (the closure of) a given operator is the generator of a Feller semigroup. For a given linear operator A , constant $\lambda > 0$, and $f \in \mathcal{C}(E)$, we say that a function $p \in \mathcal{C}(E)$ solves the *Laplace equation*

$$(\lambda - A)p = f \tag{4.6}$$

if $p \in \mathcal{D}(A)$ and (4.6) holds. The following lemma shows how solutions to Laplace equations typically arise.

Lemma 4.5 (Laplace equation) *Let G be the generator of a Feller semigroup $(P_t)_{t \geq 0}$ on $\mathcal{C}(E)$, let $\lambda > 0$ and $f \in \mathcal{C}(E)$. Then the Laplace equation $(\lambda - G)p = f$ has a unique solution, that is given by*

$$p = \int_0^\infty P_t f e^{-\lambda t} dt.$$

We say that an operator A on $\mathcal{C}(E)$ with domain $\mathcal{D}(A)$ satisfies the *positive maximum principle* if, whenever a function $f \in \mathcal{D}(A)$ assumes its maximum over E in a point $x \in E$ and $f(x) \geq 0$, we have $Af(x) \leq 0$. The following proposition gives necessary and sufficient conditions for a linear operator G to be the generator of a Feller semigroup.

Theorem 4.6 (Generators of Feller semigroups) *A linear operator G on $\mathcal{C}(E)$ is the generator of a Feller semigroup $(P_t)_{t \geq 0}$ if and only if*

- (i) $1 \in \mathcal{D}(G)$ and $G1 = 0$.
- (ii) G satisfies the positive maximum principle.
- (iii) $\mathcal{D}(G)$ is dense in $\mathcal{C}(E)$.
- (iv) For every $f \in \mathcal{C}(E)$ and $\lambda > 0$, the Laplace equation $(\lambda - G)p = f$ has a solution.

In practice, it is rarely possible to give an explicit description of the (full) domain of a Feller generator. Rather, one often starts with an operator that is defined on a smaller domain of “nice” functions and then takes its closure. In general, if G is a closed linear operator and $\mathcal{D}' \subset \mathcal{D}(G)$, then we let $G|_{\mathcal{D}'}$ denote the restriction of G to \mathcal{D}' , i.e., $G|_{\mathcal{D}'}$ is the linear operator with domain $\mathcal{D}(G|_{\mathcal{D}'}) := \overline{\mathcal{D}'}$ defined as $G|_{\mathcal{D}'}f := Gf$ for all $f \in \mathcal{D}'$. We say that \mathcal{D}' is a *core* for G if $\overline{G|_{\mathcal{D}'}} = A$.

Lemma 4.7 (Core of a generator) *Let G be the generator of a Feller semigroup and let $\mathcal{D}' \subset \mathcal{D}(G)$. Assume that \mathcal{D}' is dense in $\mathcal{C}(E)$. Then the following conditions are equivalent:*

- (i) \mathcal{D}' is a core for G ,
- (ii) the set $\{(\lambda - A)p : p \in \mathcal{D}'\}$ is dense in $\mathcal{C}(E)$ for some $\lambda > 0$,
- (iii) the set $\{(\lambda - A)p : p \in \mathcal{D}'\}$ is dense in $\mathcal{C}(E)$ for all $\lambda > 0$.

Note that by condition (ii) of Lemma 4.7, to check that a dense set $\mathcal{D}' \subset \mathcal{C}(E)$ is a core for G , it suffices to show that for some $\lambda > 0$, there exists a dense subspace $\mathcal{R} \subset \mathcal{C}(E)$ such that for every $f \in \mathcal{R}$, the Laplace equation $(\lambda - A)p = f$ has a solution $p \in \mathcal{D}'$. Using Lemma 4.7, one can prove the following alternative version of the Hille-Yosida theorem.

Theorem 4.8 (Hille-Yosida) *A linear operator A on $\mathcal{C}(E)$ with domain $\mathcal{D}(A)$ is closable and its closure $G := \overline{A}$ is the generator of a Feller semigroup if and only if*

- (i) There exist $f_n \in \mathcal{D}(A)$ such that $f_n \rightarrow 1$ and $Af_n \rightarrow 0$.
- (ii) A satisfies the positive maximum principle.
- (iii) $\mathcal{D}(A)$ is dense in $\mathcal{C}(E)$.
- (iv) For some (and hence for all) $\lambda \in (0, \infty)$, there exists a dense subspace $\mathcal{R} \subset \mathcal{C}(E)$ such that for every $f \in \mathcal{R}$, the Laplace equation $(\lambda - A)p = f$ has a solution p .

Conditions (i)–(iii) are usually easy to verify for a given operator A , but condition (iv) is the “hard” condition since this means that one has to prove existence of solutions to the Laplace equation $(\lambda - G)p = f$ for a dense set of functions f .

If K is a probability kernel on E and $r > 0$, then

$$Gf := r(Kf - f) \quad (f \in \mathcal{C}(E)) \quad (4.7)$$

defines a Feller generator that is everywhere defined (i.e., $\mathcal{D}(G) = \mathcal{C}(E)$) and hence, in view of Theorem 4.2, a bounded operator. For generators of this simple form, one can construct the corresponding semigroup by the exponential formula

$$P_t f = e^{tG} f := \sum_{n=0}^{\infty} \frac{1}{n!} (tG)^n f,$$

where the infinite sum converges in $\mathcal{C}(E)$. The corresponding Markov process has a simple description: with rate r , the process jumps from its current position x to a new position chosen according to the probability law $K(x \cdot)$.

As soon as Feller processes get more complicated in the sense that “the total rate of all things that can happen” is infinite (as will be the case for interacting particle systems), one needs the more complicated Hille-Yosida theory. To demonstrate the strength of Theorem 4.8, consider $E := [0, 1]$ and the linear operator A defined by $\mathcal{D}(A) := \mathcal{C}^2[0, 1]$ (the space of twice continuously differentiable functions on $[0, 1]$) and

$$Af(x) := x(1-x) \frac{\partial^2}{\partial x^2} f(x) \quad (x \in [0, 1]). \quad (4.8)$$

One can show that A satisfies the conditions of Theorem 4.8 and hence \overline{A} generates a Feller semigroup. The corresponding Markov process turns out to have continuous sample paths and is indeed the *Wright-Fisher diffusion* that we met before in formula (3.26).

Exercise 4.9 (Brownian motion) Let $(P_t)_{t \geq 0}$ denote the transition kernels of Brownian motion on \mathbb{R}^d . Let $E := \mathbb{R}^d \cup \{\infty\}$ denote the one-point compactification of \mathbb{R}^d and extend P_t ($t \geq 0$) to probability kernels on E by setting $P_t(\infty, \cdot) := \delta_\infty$. Show that $(P_t)_{t \geq 0}$ is a Feller semigroup.

Exercise 4.10 (Wright-Fisher diffusion) Show that the operator A defined in (4.8) satisfies the conditions of Theorem 4.8. Hint: show that if f is a polynomial of order n , then so is Af . Use this to show that the Cauchy equation $\frac{\partial}{\partial t} u_t = Au_t$ has a solution for each initial state $u_0 = f$ that is a polynomial. Then show that $p := \int_0^\infty u_t e^{-\lambda t} dt$ solves the Laplace equation $(\lambda - A)p = f$.

Some notes on the proofs

In the remainder of this section, we indicate where proofs of the stated theorems can be found. Readers who are more interested in interacting particle systems than in functional analysis may skip from here to the next section.

The fact that there is a one-to-one correspondence between continuous transition probabilities and collections $(P_t)_{t \geq 0}$ of linear operators satisfying the assumptions (i)–(iv) of a Feller semigroup follows from [Kal97, Prop. 17.14].

Theorem 4.1 (including a proof) can be found in [Kal97, Thm 17.15] and [EK86, Thm 4.2.7]. Theorem 4.2 (the closed graph theorem and characterization of continuous linear maps) can be found on many places (including Wikipedia).

Lemma 4.3 follows from [EK86, Corollary I.1.6]. The statements of this lemma can also easily be derived from the Hille-Yosida theorem (see below). Proposition 4.4 summarizes a number of well-known facts. The fact that $u_t := P_t f$ solves the Cauchy equation if $f \in \mathcal{D}(G)$ is proved in [EK86, Prop 1.1.5 (b)], [Kal97, Thm 17.6], and [Lig10, Thm 3.16 (b)]. To see that solutions to the Cauchy equation are unique, we use the following fact.

Lemma 4.11 (Positive maximum principle) *Let A be a linear operator on $\mathcal{C}(E)$ and let $u = (u_t)_{t \geq 0}$ be a solution to the Cauchy equation $\frac{\partial}{\partial t} u_t = Au_t$ ($t \geq 0$). Assume that A satisfies the positive maximum principle and $u_0 \geq 0$. Then $u_t \geq 0$ for all $t \geq 0$.*

Proof By linearity, we may equivalently show that $u_0 \leq 0$ implies $u_t \leq 0$. Assume that $u_t(x) > 0$ for some $x \in E$. By the compactness of E , the function $(x, t) \mapsto e^{-t} u_t(x)$ must assume its maximum over $E \times [0, t]$ in some point (y, s) . Our assumptions imply that $e^{-s} u_s(y) > 0$ and hence $s > 0$. But now, since A satisfies the positive maximum principle,

$$0 \leq \frac{\partial}{\partial s} (e^{-s} u_s(y)) = -e^{-s} u_s(y) + e^{-s} A u_s(y) \leq -e^{-s} u_s(y) < 0,$$

so we arrive at a contradiction. ■

By linearity, Lemma 4.11 implies that if u, v are two solutions to the same Cauchy equation and $u_0 \leq v_0$, then $u_t \leq v_t$ for all $t \geq 0$. In particular, since by Theorem 4.6, Feller generators satisfy the positive maximum principle, this implies uniqueness of solutions of the Cauchy equation in Proposition 4.4. Again by Theorem 4.6, the domain of a Feller semigroup is a dense subspace of $\mathcal{C}(E)$, so the final statement of Proposition 4.4 follows from the following simple lemma and the fact that $\|P_t f\|_\infty \leq \|f\|_\infty$.

Lemma 4.12 (Closure of bounded operators) *Let $(\mathcal{V}, \|\cdot\|)$ be a Banach space and let A be a linear operator on \mathcal{V} such that $\mathcal{D}(A)$ is dense and $\|Af\| \leq C\|f\|$ ($f \in \mathcal{D}(A)$) for some $C < \infty$. Then A is closable, $\mathcal{D}(\bar{A}) = \mathcal{V}$, and $\|\bar{A}f\| \leq C\|f\|$ ($f \in \mathcal{V}$).*

Proof (sketch) Since $\mathcal{D}(A)$ is dense, for each $f \in \mathcal{V}$ we can choose $\mathcal{D}(A) \ni f_n \rightarrow f$. Using the fact that A is bounded, it is easy to check that if $(f_n)_{n \geq 0}$ is a Cauchy sequence and $f_n \in \mathcal{D}(A)$ for all n , then $(Af_n)_{n \geq 0}$ is also a Cauchy sequence. By the completeness of \mathcal{V} , it follows that the limit $\bar{A}f := \lim_{n \rightarrow \infty} Af_n$ exists for all $f \in \mathcal{V}$. To see that this defines \bar{A} unambiguously, assume that $f_n \rightarrow f$ and $g_n \rightarrow f$ and observe that $\|Af_n - Ag_n\| \leq C\|f_n - g_n\| \rightarrow 0$. The fact that $\|\bar{A}f\| \leq C\|f\|$ ($f \in \mathcal{V}$) follows from the continuity of the norm. ■

Lemma 4.5 follows from [EK86, Prop 1.2.1]. Theorems 4.6 and 4.8 both go under the name of the Hille-Yosida theorem. Often, they are stated in a more general form without condition (i). In this generality, the operator G generates a semigroup of *subprobability kernels* $(P_t)_{t \geq 0}$, i.e., $P_t(x, \cdot)$ is a measure with total mass $P_t(x, E) \leq 1$. In this context, a Feller semigroup with $P_t(x, E) = 1$ for all t, x is called *conservative*. It is clear from Proposition 4.4 that condition (i) in Theorems 4.6 and 4.8 is necessary and sufficient for the Feller group to be conservative.

The versions of the Hille-Yosida theorem stated in [EK86, Kal97] are more general than Theorems 4.6 and 4.8 since they allow for the case that E is not compact but only locally compact. This is not really more general, however, since what these books basically do if E is not compact is the following. First, they construct the one-point compactification $\bar{E} = E \cup \{\infty\}$ of E . Next, they extend the transition probabilities to \bar{E} by putting $P_t(\infty, \cdot) := \delta_\infty$ for all $t \geq 0$. Having proved that they generate a conservative Feller semigroup on \bar{E} of this form, they then still need to prove that the associated Markov process does not explode in the sense that $\mathbb{P}^x[X_t \in E \ \forall t \geq 0] = 1$. In practical situations (such as when constructing Markov processes with state space \mathbb{R}^d) it is usually better to explicitly work with the one-point compactification of \mathbb{R}^d instead of trying to formulate theorems for locally compact spaces that try hide this compactification in the background.

Theorems 4.6 and 4.8 are special cases of more general theorems (also called Hille-Yosida theorem) for strongly continuous contraction semigroups taking values in a general Banach space. In this context, the positive maximum principle is replaced by the assumption that the operator under consideration is *dissipative*. In this more general setting, Theorems 4.6 and 4.8 correspond to [EK86, Thms 1.2.6 and 1.2.12]. Lemma 4.7 follows from [EK86, Lemma 1.2.11 and Prop 1.3.1]. In the more specific set-up of Feller semigroups, versions of Theorem 4.8 can be found in [EK86, Thm 4.2.2] and [Kal97, Thm 17.11]. There is also an account of Hille-Yosida theory for Feller semigroups in [Lig10, Chap 3], but this reference does not mention the positive maximum principle (using a dissipativity assumption instead).

Feller semigroups with bounded generators such as in (4.7) are treated in [EK86, Sect 4.2] and [Kal97, Prop 17.2]. The fact that the operator A in (4.8) satisfies the assumptions of Theorem 4.8 is proved in [EK86, Thm 8.2.8].

4.3 Poisson construction

We briefly recall the set-up introduced in Section 4.1. S is a finite set, called the *local state space*, and Λ is a countable set, called the *lattice*. We equip the product space S^Λ with the product topology, making it into a compact metrizable space. Elements of S^Λ are denoted $x = (x(i))_{i \in \Lambda}$.

By definition, a *local probability kernel* is a probability kernel K on S^Λ such that:

- (i) K is continuous,
- (ii) there exists a finite set $\Delta \subset \Lambda$ such that $K(x, \cdot)$ is concentrated on $\{y \in S^\Lambda : y(i) = x(i) \ \forall i \in \Lambda \setminus \Delta\}$ for all $x \in S^\Lambda$.

Condition (ii) says that if starting from x , we choose a new configuration according to $K(x, \cdot)$, then only the local states in the finite set Δ can change. An *interacting particle system* is a Feller process with state space of the form S^Λ and generator of the form

$$Gf = \sum_{K \in \mathcal{K}} r_K \{Kf - f\}, \quad (4.9)$$

where \mathcal{K} is a countable collection of local probability kernels on S^Λ and $(r_K)_{K \in \mathcal{K}}$ are nonnegative constants. As explained in the previous section, we cannot expect Gf to be defined for all $f \in \mathcal{C}(S^\Lambda)$, but instead define Gf first for a class of “nice” functions and then find the full generator by taking the closure. We wish to give sufficient conditions so that the closure of G is the generator of a Feller process.

We will use a Poisson construction in the spirit of Proposition 2.7. In the next section, it will then be shown that the process constructed in this way indeed has a generator of the form (4.10). Since we want to use a Poisson construction, we specialize to the case that the kernels $K \in \mathcal{K}$ are all deterministic. By definition, a *local map* is a function $m : S^\Lambda \rightarrow S^\Lambda$ such that:

- (i) m is continuous,
- (ii) there exists a finite set $\Delta \subset \Lambda$ such that $m(x)(i) = x(i)$ for all $i \in \Lambda \setminus \Delta$ and $x \in S^\Lambda$.

It is easy to see that a local probability kernel is deterministic if and only if it is of the form $K(x, \cdot) = \delta_{m(x)}$ for some local map m . We will therefore be interested in generators of the form

$$Gf(x) = \sum_{m \in \mathcal{G}} r_m \{f(m(x)) - f(x)\} \quad (x \in S^\Lambda), \quad (4.10)$$

where \mathcal{G} is a countable collection of local maps and $(r_m)_{m \in \mathcal{G}}$ are nonnegative rates.

For any map $m : S^\Lambda \rightarrow S^\Lambda$ and $i \in \Lambda$, we define $m[i] : S^\Lambda \rightarrow S$ by $m[i](x) := m(x)(i)$ ($x \in S^\Lambda$, $i \in \Lambda$). We let

$$\begin{aligned} \mathcal{D}(m) &:= \{i \in \Lambda : \exists x \in S^\Lambda \text{ s.t. } m(x)(i) \neq x(i)\}, \\ &= \{i \in \Lambda : m[i] \neq 1\} \end{aligned}$$

denote the set of lattice points i for which $m[i]$ is not the identity map. Then m is local if and only if $\mathcal{D}(m)$ is finite and m is continuous. It follows immediately from the definition of the product topology that m is continuous if and only if $m[i]$ is continuous for all $i \in \Lambda$.

Let S and T be finite sets, let Λ be a countable set, and let $f : S^\Lambda \rightarrow T$ be a function. Then we say that a point $j \in \Lambda$ is *f-relevant* if

$$\exists x, y \in S^\Lambda \text{ s.t. } f(x) \neq f(y) \text{ and } x(k) = y(k) \ \forall k \neq j,$$

i.e., changing the value of x in j may change the value of $f(x)$. We write

$$\mathcal{R}(f) := \{j \in \Lambda : j \text{ is } f\text{-relevant}\}.$$

The following lemma (which we have taken from [SS15b, Lemma 24]) says that a function $f : S^\Lambda \rightarrow T$ is continuous with respect to the product topology if and only if it depends on finitely many coordinates.

Lemma 4.13 (Continuous maps) *Let S and T be finite sets and let Λ be a countable set. Then a function $f : S^\Lambda \rightarrow T$ is continuous with respect to the product topology if and only if the following two conditions are satisfied:*

- (i) $\mathcal{R}(f)$ is finite,
- (ii) If $x, y \in S^\Lambda$ satisfy $x(j) = y(j)$ for all $j \in \mathcal{R}(f)$, then $f(x) = f(y)$.

Before we give the proof of Lemma 4.13, we first make some observations. The following exercise shows how continuity can fail if condition (i) of Lemma 4.13 does not hold.

Exercise 4.14 (A discontinuous map) Let $2\mathbb{N} := \{2n : n \in \mathbb{N}\}$ and $2\mathbb{N} + 1 := \{2n + 1 : n \in \mathbb{N}\}$. Define $f : \{0, 1\}^{\mathbb{N}} \rightarrow \{0, 1\}$ by

$$f(x) := \begin{cases} 1 & \text{if } \inf\{i \in \mathbb{N} : x(i) = 1\} \in 2\mathbb{N} \cup \{\infty\}, \\ 0 & \text{if } \inf\{i \in \mathbb{N} : x(i) = 1\} \in 2\mathbb{N} + 1. \end{cases} \quad (4.11)$$

Show that f satisfies condition (ii) of Lemma 4.13 but not condition (i). Show that f is not continuous.

The following exercise shows that contrary to what one might initially have guessed, condition (ii) of Lemma 4.13 is not automatically satisfied, even when condition (i) holds.

Exercise 4.15 (Another discontinuous map) Define $f : \{0, 1\}^{\mathbb{N}} \rightarrow \{0, 1\}$ by

$$f(x) := \begin{cases} 1 & \text{if } \{i \in \mathbb{N} : x(i) = 1\} \text{ is finite,} \\ 0 & \text{if } \{i \in \mathbb{N} : x(i) = 1\} \text{ is infinite.} \end{cases} \quad (4.12)$$

Show that f satisfies condition (i) of Lemma 4.13 but not condition (ii). Show that f is not continuous.

Proof of Lemma 4.13 Let $(\alpha_j)_{j \in \Lambda}$ be strictly positive constants such that $\sum_{j \in \Lambda} \alpha_j < \infty$. Then the metric

$$d(x, y) := \sum_{j \in \Lambda} \alpha_j 1_{\{x(j) \neq y(j)\}} \quad (x, y \in S^\Lambda) \quad (4.13)$$

generates the product topology on S^Λ . By Tychonoff's theorem, S^Λ is compact, so the function f is uniformly continuous. Since the target space T is finite, this means that there exists an $\varepsilon > 0$ such that $d(x, y) < \varepsilon$ implies $f(x) = f(y)$. Since $\sum_{j \in \Lambda} \alpha_j < \infty$, there exists some finite $\Lambda' \subset \Lambda$ such that $\sum_{j \in \Lambda \setminus \Lambda'} \alpha_j < \varepsilon$. It follows that

(ii)' If $x, y \in S^\Lambda$ satisfy $x(j) = y(j)$ for all $j \in \Lambda'$, then $f(x) = f(y)$.

We conclude from this that $\mathcal{R}(f) \subset \Lambda'$, proving (i). If this is a strict inclusion, then we can inductively remove those points from Λ' that are not elements of $\mathcal{R}(f)$ while preserving the property (ii)', until in a finite number of steps we see that (ii) holds.

Conversely, if (i) and (ii) hold and $x_k \rightarrow x$ pointwise, then by (i) there exists some n such that $x_k(j) = x(j)$ for all $j \in \mathcal{R}(f)$ and hence by (ii) $f(x_k) = f(x)$ for all $k \geq n$, proving that f is continuous. ■

We observe that a map $m : S^\Lambda \rightarrow S^\Lambda$ is local if and only if $\mathcal{D}(m)$ is finite and $m[i]$ satisfies conditions (i) and (ii) of Lemma 4.13 for each $i \in \mathcal{D}(m)$. The following exercise describes yet another way to look at local maps.

Exercise 4.16 Show that a map $m : S^\Lambda \rightarrow S^\Lambda$ is local if and only if there exists a finite $\Delta \subset \Lambda$ and a map $m' : S^\Delta \rightarrow S^\Delta$ such that

$$m(x)(k) = \begin{cases} m'((x(i))_{i \in \Delta})(k) & \text{if } k \in \Delta, \\ x(k) & \text{otherwise.} \end{cases}$$

Before we continue, it is good to see a number of examples.

- The voter map vot_{ij} defined in (1.4) satisfies

$$\mathcal{D}(\text{vot}_{ij}) = \{j\} \quad \text{and} \quad \mathcal{R}(\text{vot}_{ij}[j]) = \{i\},$$

since only the type at j changes, and it suffices to know the type at i to predict the new type of j .

- The branching map bra_{ij} defined in (1.6) satisfies

$$\mathcal{D}(\text{bra}_{ij}) = \{j\} \quad \text{and} \quad \mathcal{R}(\text{bra}_{ij}[j]) = \{i, j\},$$

since only the type at j changes, but we need to know both the type at i and j to predict the new type of j since $\text{bra}_{ij}(x)(j) = x(i) \vee x(j)$.

- The death map death_i defined in (1.7) satisfies

$$\mathcal{D}(\text{death}_i) = \{i\} \quad \text{and} \quad \mathcal{R}(\text{death}_i[i]) = \emptyset$$

since only the type at i changes, and the new type at i is 0 regardless of x .

- The coalescing random walk map rw_{ij} defined in (1.20) satisfies

$$\mathcal{D}(\text{rw}_{ij}) = \{i, j\}, \quad \mathcal{R}(\text{rw}_{ij}[i]) = \emptyset, \quad \text{and} \quad \mathcal{R}(\text{rw}_{ij}[j]) = \{i, j\},$$

since the types at both i and j can change, the new type at i is 0 regardless of the previous state, but to calculate $\text{rw}_{ij}(x)(j)$ we need to know both $x(i)$ and $x(j)$.

Exercise 4.17 Recall the exclusion map excl_{ij} defined in (1.23) and the cooperative branching map coop_{ij} defined in (1.25). For $m = \text{excl}_{ij}$ or $m = \text{coop}_{ij}$, determine $\mathcal{D}(m)$, and determine $\mathcal{R}(m[i])$ for all $i \in \mathcal{D}(m)$.

Let \mathcal{G} be a countable set whose elements are local maps $m : S^\Lambda \rightarrow S^\Lambda$, let $(r_m)_{m \in \mathcal{G}}$ be nonnegative constants, and (as in Proposition 2.7) let ω be a Poisson point set on $\mathcal{G} \times \mathbb{R}$ with intensity $r_m dt$. Since \mathcal{G} is countable, by the argument used in Section 2.3, it is easy to see that almost surely, the

time coordinates of all points $(m, t) \in \omega$ are all different. Therefore, as we did in the finite setting, we can still unambiguously define a random function $\mathbb{R} \ni t \mapsto \mathbf{m}_t^\omega$ by setting

$$\mathbf{m}_t^\omega := \begin{cases} m & \text{if } (m, t) \in \omega, \\ 1 & \text{otherwise,} \end{cases}$$

where we write 1 to denote the identity map. For $s \in \mathbb{R}$, and $x \in S^\Lambda$, we will be interested in cadlag functions $[s, \infty) \ni t \mapsto X_t \in S^\Lambda$ that solve the equation

$$X_s = x \quad \text{and} \quad X_t = \mathbf{m}_t^\omega(X_{t-}) \quad (t > s). \quad (4.14)$$

The following exercise shows that at least in the case when $\omega = \emptyset$, the equation (4.14) has a unique solution.

Exercise 4.18 *A topological space E is totally disconnected if for each $x_1, x_2 \in E$ with $x_1 \neq x_2$, there exist open sets $O_1 \ni x_1$ and $O_2 \ni x_2$ such that $O_1 \cap O_2 = \emptyset$ and $O_1 \cup O_2 = E$. Prove that S^Λ is totally disconnected. Prove that if E is a totally disconnected space, then each continuous function $f : [0, 1] \rightarrow E$ is constant.*

The difficulty with proving existence and uniqueness of solutions of (4.14) is that we will typically have that $\sum_{m \in \mathcal{G}} r_m = \infty$. As a result, $\{t : (m, t) \in \omega\}$ will be a dense subset of \mathbb{R} , so it will no longer be possible to order the elements of $\omega_{s,t}$ according to their times as we did in (2.12) to see that (4.14) has a unique solution. Nevertheless, since our maps m are local, we can hope that under suitable assumptions on the rates, only finitely many points of $\omega_{s,t}$ are needed to determine the local state $X_t(i)$ of our process at a given lattice point $i \in \Lambda$ and time $t \geq s$. This intuition is made precise in the following theorem, which we will prove below, and which will be one of the main results of this section.

Theorem 4.19 (Pathwise solution) *Let \mathcal{G} be a countable set whose elements are local maps $m : S^\Lambda \rightarrow S^\Lambda$, let $(r_m)_{m \in \mathcal{G}}$ be nonnegative constants satisfying*

$$\sup_{i \in \Lambda} \sum_{\substack{m \in \mathcal{G} \\ \mathcal{D}(m) \ni i}} r_m (|\mathcal{R}(m[i])| + 1) < \infty, \quad (4.15)$$

and let ω be a Poisson point set on $\mathcal{G} \times \mathbb{R}$ with intensity $r_m dt$. Then, almost surely, for each $s \in \mathbb{R}$ and $x \in S^\Lambda$, there exists a unique solution $X = (X_t)_{t \geq s}$ to (4.14).

Note that in Theorem 4.19, the almost sure statement holds for all $s \in \mathbb{R}$ and $x \in S^\Lambda$ simultaneously. This allows us to define random maps $(\mathbf{X}_{s,u})_{s \leq u}$ by

$$\mathbf{X}_{s,u}(x) := X_u \quad \text{where } (X_t)_{t \geq s} \text{ solves (4.14).} \quad (4.16)$$

It is easy to see that the random maps $(\mathbf{X}_{s,u})_{s \leq u}$ form a stochastic flow in the sense of (2.13) and have independent increments. The following theorem says that these random maps can be used to define a Markov process just as in Proposition 2.7, and the resulting process is in fact a Feller process.

Theorem 4.20 (Poisson construction of particle systems) *Assume (4.15). Let $(\mathbf{X}_{s,u})_{s \leq u}$ be the random maps defined in (4.16) and let X_0 be an S^Λ -valued random variable, independent of $(\mathbf{X}_{s,u})_{s \leq u}$. Then*

$$X_t := \mathbf{X}_{0,t}(X_0) \quad (t \geq 0) \quad (4.17)$$

defines a Feller process with semigroup $(P_t)_{t \geq 0}$ given by

$$P_t(x, \cdot) := \mathbb{P}[\mathbf{X}_{0,t}(x) \in \cdot] \quad (x \in S^\Lambda, t \geq 0). \quad (4.18)$$

Theorems 4.19 and 4.20 give a “pathwise” construction of interacting particle systems based on i.i.d. randomness. In this respect, they are broadly similar to other constructions of Markov processes from independent randomness such as, for example, the construction of diffusion processes as the pathwise unique solutions to stochastic differential equations driven by Brownian motions.

To get a first impression of why a result like Theorem 4.19 might be true, let us look at the contact process. In this case, the generator takes the form (1.8) and elements of ω are points of the form (\mathbf{bra}_{ij}, t) or (\mathbf{death}_i, t) , which indicate that the corresponding local map should be applied at time t . We will call elements $(m, t) \in \omega$ *incidents*.⁸ In Figure 4.1, we have drawn space horizontally and time vertically and visualized one random realization of ω in such a way that for each incident (m, t) we draw a symbol representing the map m at the time t and at the sites that are involved in the map. Such a picture is called a *graphical representation* (also called *graphical construction*) for an interacting particle system. In practice, various symbols (such as arrows, squares, stars etc.) are used to indicate different maps. Our aim is to find sufficient conditions under which such a graphical representation almost surely yields a well-defined process.

As a first step, we observe that for each $i \in \Lambda$, the set

$$\{t \in \mathbb{R} : \exists m \in \mathcal{G} \text{ s.t. } i \in \mathcal{D}(m), (m, t) \in \omega\}$$

⁸Another natural choice of terminology would be “event” but this word already has a quite different meaning in probability theory.




Figure 4.1: Graphical representation of a one-dimensional contact process, with paths of potential influence and the set $\zeta_0^t(\{4\})$ of lattice points at time zero whose value is relevant for $X_t(4)$.

is a Poisson point set with intensity $\sum_{m \in \mathcal{G}, \mathcal{D}(m) \ni i} r_m$. Therefore, provided that

$$K_0 := \sup_i \sum_{\substack{m \in \mathcal{G} \\ \mathcal{D}(m) \ni i}} r_m < \infty, \quad (4.19)$$

each finite time interval contains only finitely many incidents that have the potential to change the state of a given lattice point i . This does not automatically imply, however, that our process is well-defined, since incidents that happen at i might depend on incidents that happen at other sites at earlier times, and in this way a large and possibly infinite number of incidents and lattice points can potentially influence the state of a single lattice point at a given time.

For any $i, j \in \Lambda$ and $s < u$, by definition, a *path of potential influence* from (i, s) to (j, u) is a cadlag function $\gamma : (s, u] \rightarrow \Lambda$ such that $\gamma_{s-} = i$, $\gamma_u = j$, and

- (i) if $\gamma_{t-} \neq \gamma_t$ for some $t \in (s, u]$, then there exists some $m \in \mathcal{G}$ such that $(m, t) \in \omega$, $\gamma_t \in \mathcal{D}(m)$ and $\gamma_{t-} \in \mathcal{R}(m[\gamma_t])$,
 - (ii) for each $(m, t) \in \omega$ with $t \in (s, u]$ and $\gamma_t \in \mathcal{D}(m)$, one has $\gamma_{t-} \in \mathcal{R}(m[\gamma_t])$.
- (4.20)

We write $(i, s) \rightsquigarrow (j, u)$ if there is a path of potential influence from (i, s) to

(j, u) . Similarly, for any $A \subset \Lambda$, we write $(i, s) \rightsquigarrow A \times \{u\}$ if there is a path of potential influence from (i, s) to some point (j, u) with $j \in A$. For any finite set $A \subset \Lambda$ and $s < u$, we set

$$\zeta_s^u(A) := \{i \in \Lambda : (i, s) \rightsquigarrow A \times \{u\}\}, \quad (4.21)$$

and we let $\zeta_u^u(A) := A$. If we start the process at time zero, then $\zeta_0^t(A)$ will be the set of lattice points whose values at time zero are potentially relevant for the local state of the process in A at time t , in a way that will be made more precise in Lemma 4.26 below. See Figure 4.1 for a picture of $\zeta_u^u(A)$ and the collection of all paths of potential influence that end in $A \times \{t\}$. The following lemma will be the cornerstone of our Poisson construction of interacting particle systems.

Lemma 4.21 (Exponential bound) *Assume that the rates $(r_m)_{m \in \mathcal{G}}$ satisfy (4.19) and that*

$$K := \sup_{i \in \Lambda} \sum_{\substack{m \in \mathcal{G} \\ \mathcal{D}(m) \ni i}} r_m (|\mathcal{R}(m[i])| - 1) < \infty. \quad (4.22)$$

Then, for each finite $A \subset \Lambda$, one has

$$\mathbb{E}[|\zeta_s^u(A)|] \leq |A| e^{K(u-s)} \quad (s \leq u). \quad (4.23)$$

Proof To simplify notation, we fix A and u and write

$$\xi_t := \zeta_{u-t}^u(A) \quad (t \geq 0).$$

The idea of the proof is as follows: since $(\xi_t)_{t \geq 0}$ is constructed from Poisson incidents in the spirit of Propositions 2.7 and 2.8, it is in fact a Markov process. Using a simple generator calculation, one finds that $\frac{\partial}{\partial t} \mathbb{E}[|\xi_t|] \leq K \mathbb{E}[|\xi_t|]$, which implies that $\mathbb{E}[|\xi_t|] \leq \mathbb{E}[|\xi_0|] e^{Kt}$. To make this idea precise, we use a cut-off to ensure that the state space of our Markov process is finite and Proposition 2.1 is applicable, and we also modify ξ_t so that its sample paths are right-continuous, as is the habit for Markov processes.

We start with the cut-off. Let $\Lambda_n \subset \Lambda$ be finite sets such that $\Lambda_n \uparrow \Lambda$. For n large enough such that $A \subset \Lambda_n$, let us write

$$\xi_t^n := \{i \in \Lambda : (i, t) \rightsquigarrow_n A \times \{u\}\},$$

where $(i, s) \rightsquigarrow_n A \times \{u\}$ denotes the presence of a path of potential influence from (i, s) to $A \times \{u\}$ that stays in Λ_n . We observe that since $\Lambda_n \uparrow \Lambda$, we have

$$\xi_t^n \uparrow \xi_t \quad (t \geq 0).$$

The process $(\xi_t^n)_{t \geq 0}$ is left-continuous; let $\xi_{t+}^n := \lim_{s \downarrow t} \xi_s^n$ denote its right-continuous modification. We claim that $(\xi_{t+}^n)_{t \geq 0}$ is a Markov process. To see this, note that one can have $\xi_{t+}^n \neq \xi_t^n$ only when $(m, u - t) \in \omega$ for some $m \in \mathcal{G}$ and at such an instant, if the previous state is $\xi_t^n = A$, then the new state is $\xi_{t+}^n = A^m$, where we define

$$A^m := \Lambda_n \cap \bigcup_{i \in A} \mathcal{R}(m[i]).$$

Since $\mathcal{R}(m[i]) = \{i\}$ if $i \notin \mathcal{D}(m)$, it suffices to consider only those incidents for which $m \in \mathcal{G}_n := \{m \in \mathcal{G} : \mathcal{D}(m) \cap \Lambda_n \neq \emptyset\}$. It follows from (4.19) that the total rate $\sum_{m \in \mathcal{G}_n} r_m$ at which maps from \mathcal{G}_n are applied is finite. Proposition 2.7 now implies that the process $(\xi_{t+}^n)_{t \geq 0}$ is a Markov process taking values in the (finite) space of all subsets of Λ_n , with generator

$$G_n f(A) := \sum_{m \in \mathcal{G}_n} r_m (f(A^m) - f(A)).$$

Let $(P_t^n)_{t \geq 0}$ be the associated semigroup and let f be the function $f(A) := |A|$. Then

$$\begin{aligned} G_n f(A) &= \sum_{m \in \mathcal{G}_n} r_m (f(A^m) - f(A)) \\ &\leq \sum_{m \in \mathcal{G}_n} r_m \left(|A \setminus \mathcal{D}(m)| + \sum_{i \in A \cap \mathcal{D}(m)} |\mathcal{R}(m[i])| - |A| \right) \\ &= \sum_{m \in \mathcal{G}_n} r_m \left(\sum_{i \in A \cap \mathcal{D}(m)} (|\mathcal{R}(m[i])| - 1) \right) \\ &= \sum_{i \in A} \sum_{\substack{m \in \mathcal{G}_n \\ \mathcal{D}(m) \ni i}} r_m (|\mathcal{R}(m[i])| - 1) \leq K|A|. \end{aligned}$$

It follows that

$$\frac{\partial}{\partial t} (e^{-Kt} P_t^n f) = -K e^{-Kt} P_t^n f + e^{-Kt} P_t^n G_n f = e^{-Kt} P_t^n (G_n f - Kf) \leq 0$$

and therefore $e^{-Kt} P_t^n f \leq e^{-K0} P_0^n f = f$, which means that

$$\mathbb{E}[|\xi_t^n|] \leq |A| e^{Kt} \quad (t \geq 0). \quad (4.24)$$

Letting $n \uparrow \infty$ we arrive at (4.23). ■

Lemma 4.21 almost shows that the Markov process $(\zeta_{u-t}^u(A))_{t \geq 0}$ is non-explosive, but not quite, since it only gives a bound on the size of $\zeta_{u-t}^u(A)$ at deterministic times and also does not exclude the possibility that $\zeta_{u-t}^u(A)$ explodes in a different way than by growing to infinite size, for example, by containing a few sites very far away. The following lemma fixes this.

Lemma 4.22 (Finitely many relevant sites) *For each $A \subset \Lambda$ and $s \leq u$, let*

$$Z(A, s, u) := \bigcup \{ \zeta_{s'}^{u'}(A) : s \leq s' \leq u' \leq u \}.$$

Assume that the rates $(r_m)_{m \in \mathcal{G}}$ satisfy (4.19) and (4.22). Then, almost surely, $|Z(A, s, u)| < \infty$ for all $s \leq u$ and finite $A \subset \Lambda$.

Proof Let

$$\xi_s^u(A) := \{i \in \Lambda : (i, s) \rightsquigarrow' A \times \{u\}\},$$

where \rightsquigarrow' is defined in a similar way as \rightsquigarrow , except that we drop condition (ii) from the definition of a path of potential influence in (4.20). Clearly $\xi_s^u(A) \supset \zeta_s^u(A)$. For fixed u , the process $(\xi_{u-t}^u(A))_{t \geq 0}$ is a Markov process, just as was the case for $(\zeta_{u-t}^u(A))_{t \geq 0}$, but unlike the latter, the process $(\xi_{u-t}^u(A))_{t \geq 0}$ has the property that once a site lies inside $\xi_{u-t}^u(A)$, it cannot be removed, so $\xi_{s'}^u(A) \subset \xi_s^u(A)$ for all $s \leq s' \leq u$. In particular, this implies that $\xi_s^u(A) \supset \xi_u^u(A) = A$ for all $s \leq u$ and hence

$$\xi_s^{u'}(A) \subset \xi_s^u(\xi_{u'}^u(A)) = \xi_s^u(A) \quad (s \leq u' \leq u).$$

Combining this with our earlier claims, it follows that

$$Z(A, s, u) \subset \bigcup \{ \xi_{s'}^{u'}(A) : s \leq s' \leq u' \leq u \} \subset \xi_s^u(A) \subset \xi_{[s]}^{[u]}(A) \quad (s \leq u),$$

where $[u]$ and $[s]$ denote the numbers s and u rounded up and down to the nearest integer, respectively. Since moreover $\xi_s^u(A) = \bigcup_{i \in A} \xi_s^u(\{i\})$, it therefore suffices to prove that almost surely, $\xi_s^u(\{i\})$ is finite for all $s, u \in \mathbb{Z}$ with $s \leq u$ and for all $i \in \Lambda$. Since \mathbb{Z} and Λ are countable, it suffices to prove that $\xi_s^u(\{i\})$ is finite for fixed s, u , and i .

By exactly the same argument used to prove Lemma 4.21, we obtain

$$\mathbb{E}[|\xi_s^{A,u}|] \leq |A|e^{K_1(u-s)} \quad (s \leq u), \quad (4.25)$$

where

$$K_1 := \sup_{i \in \Lambda} \sum_{\substack{m \in \mathcal{G} \\ \mathcal{D}(m) \ni i}} r_m |\mathcal{R}(m[i])|, \quad (4.26)$$

where

$$K_1 = \sup_{i \in \Lambda} \left(\sum_{\substack{m \in \mathcal{G} \\ \mathcal{D}(m) \ni i}} r_m (|\mathcal{R}(m[i])| - 1) + \sum_{\substack{m \in \mathcal{G} \\ \mathcal{D}(m) \ni i}} r_m \right) \leq K + K_0 \quad (4.27)$$

is finite by (4.19) and (4.22). \blacksquare

For any $s \leq u$ and finite set $\tilde{\omega} \subset \omega_{s,t}$, we can order the elements according to the time when they occur:

$$\tilde{\omega} = \{(m_1, t_1), \dots, (m_n, t_n)\} \quad \text{with} \quad t_1 < \dots < t_n.$$

Then, setting

$$\mathbf{X}_{s,u}^{\tilde{\omega}}(x) := m_n \circ \dots \circ m_1(x) \quad (x \in S^\Lambda, s \leq u)$$

defines a map $\mathbf{X}_{s,u}^{\tilde{\omega}} : S^\Lambda \rightarrow S^\Lambda$. Our aim is to show that the pointwise limit

$$\lim_{\tilde{\omega}_n \uparrow \omega_{s,t}} \mathbf{X}_{s,u}^{\tilde{\omega}_n}(x)$$

exists and the limit does not depend on the choice of the finite sets $\tilde{\omega}_n \uparrow \omega_{s,t}$. For each $s \leq u$ and $i \in \Lambda$, we let

$$\omega_s^u(i) := \{(m, t) \in \omega_{s,u} : \mathcal{D}(m) \cap \zeta_t^u(\{i\}) \neq \emptyset\}$$

denote the set of Poisson incidents between times s and u that are relevant to determine the value of $\mathbf{X}_{s,u}(x)(i)$. Note that $\omega_s^u(i) \supset \omega_{s'}^u(i)$ for all $s \leq s' \leq u$. The following lemma says that only finitely many incidents are relevant for the evolution of an interacting particle system in a finite piece of space-time.

Lemma 4.23 (Finitely many relevant incidents) *Assume that the rates $(r_m)_{m \in \mathcal{G}}$ satisfy (4.19) and (4.22). Then, almost surely, the set $\bigcup_{t \in (s,u]} \omega_s^t(i)$ is finite for all $s \leq u$ and $i \in \Lambda$.*

Proof If $(m, t) \in \bigcup_{r \in (s,u]} \omega_s^r(i)$, then $t \in (s, u]$ and $\mathcal{D}(m) \cap Z(\{i\}, s, u) \neq \emptyset$. It follows from Lemma 4.22 that the set $Z(\{i\}, s, u)$ is finite while the assumption (4.19) implies that $\{(m, t) \in \omega_{s,u} : \mathcal{D}(m) \cap Z \neq \emptyset\}$ is finite for each $s \leq u$ and finite $Z \subset \Lambda$. \blacksquare

The following lemma is a precise formulation of the notion that $\omega_s^u(i)$ contains all incidents that are relevant to determine the value of $\mathbf{X}_{s,u}(x)(i)$.

Lemma 4.24 (Only the relevant incidents matter) *Assume that the rates $(r_m)_{m \in \mathcal{G}}$ satisfy (4.19) and (4.22). Then almost surely, for each $i \in \Lambda$, $s \leq u$, and $x \in S^\Lambda$, one has*

$$\mathbf{X}_{s,u}^{\tilde{\omega}}(x)(i) = \mathbf{X}_{s,u}^{\omega_s^{i,u}}(x)(i) \quad \text{for all } \tilde{\omega} \supset \omega_s^{i,u}.$$

Proof Immediate from our definitions. ■

We are now ready to prove the first main result of this section.

Proof of Theorem 4.19 The condition (4.15) clearly implies (4.19) and (4.22). Lemmas 4.23 and 4.24 now imply that almost surely, for each $s \leq u$, the pointwise limit

$$\mathbf{X}_{s,u}(x) := \lim_{\tilde{\omega}_n \uparrow \omega_{s,t}} \mathbf{X}_{s,u}^{\tilde{\omega}_n}(x),$$

exists and the limit does not depend on the choice of the finite sets $\tilde{\omega}_n \uparrow \omega_{s,t}$. We will show (4.14) has a unique solution, which is given by $X_t = \mathbf{X}_{s,t}(x)$ ($t \geq s$).

It suffices to show that almost surely, for each $s \leq u$ and $x \in S^\Lambda$, there exists a unique cadlag function $[s, u] \ni t \mapsto X_t$ such that

$$X_s = x \quad \text{and} \quad X_t = \mathbf{m}_t^\omega(X_{t-}) \quad (t \in (s, u]). \quad (4.28)$$

Lemma 4.23 tells us that for each $i \in \Lambda$, we can choose a finite set $\tilde{\omega}_i \subset \omega_{s,u}$ that contains $\bigcup_{t \in [s, u]} \omega_s^t(i)$. We use this to define

$$X_t(i) := \mathbf{X}_{s,t}(x)(i) = \mathbf{X}_{s,t}^{\tilde{\omega}_i}(x)(i) \quad (s \leq t \leq u, i \in \Lambda). \quad (4.29)$$

The function $t \mapsto \mathbf{X}_{s,t}^{\tilde{\omega}_i}(x)(i)$ is clearly piecewise constant, right-continuous, and changes only at times t when $i \in \mathcal{D}(m)$ for some $(m, t) \in \omega_{s,u}$, in which case $X_t = m(X_{t-})$. This shows that $(X_t)_{t \in [s, u]}$ solves (4.28).

To see that solutions to (4.28) are unique, imagine that $(X'_t)_{t \in [s, u]}$ also solves (4.28). Fix $i \in \Lambda$ and $r \in (s, u]$. Since $(X_t)_{t \in [s, u]}$ and $(X'_t)_{t \in [s, u]}$ both solve (4.28), they must be equal on the set $\zeta_s^r(\{i\})$ during the time interval $[s, t)$, where t is the first time t when $(m, t) \in \omega_s^r(i)$ for some $m \in \mathcal{G}$. But then they must be equal on the set $\zeta_t^r(\{i\})$ at time t as well. Continuing in this way, using the fact that $\omega_s^r(i)$ is finite, we see that $X_r(i) = X'_r(i)$. Since this holds for general i and r , the claim follows. ■

The proof of Theorem 4.19 yields a useful consequence.

Corollary 4.25 (Poisson construction of the stochastic flow) *The random maps $(\mathbf{X}_{s,u})_{s \leq u}$ defined in (4.16) are given by the pointwise limit*

$$\mathbf{X}_{s,u}(x) := \lim_{\tilde{\omega}_n \uparrow \omega_{s,t}} \mathbf{X}_{s,u}^{\tilde{\omega}_n}(x), \quad (4.30)$$

where the limit does not depend on the choice of the finite sets $\tilde{\omega}_n \uparrow \omega_{s,t}$.

The following useful lemma also follows easily from our constructions so far. Recall Lemma 4.13 which linked continuity of a map $m : S^\Lambda \rightarrow S^\Lambda$ with respect to the product topology to the sets $\mathcal{R}(m[i])$.

Lemma 4.26 (Continuity of the stochastic flow) *The random maps $(\mathbf{X}_{s,u})_{s \leq u}$ defined in (4.16) are continuous with respect to the product topology and satisfy*

$$\mathcal{R}(\mathbf{X}_{s,u}[i]) \subset \zeta_s^u(\{i\}) \quad (i \in \Lambda, s \leq u). \quad (4.31)$$

Proof Our construction implies that if $x, y \in S^\Lambda$ satisfy $x(j) = y(j)$ for all $j \in \zeta_s^u(\{i\})$, then $\mathbf{X}_{s,u}(x)(i) = \mathbf{X}_{s,u}(y)(i)$. This implies (4.31). Lemma 4.22 shows that almost surely, the sets $\zeta_s^u(\{i\})$ are finite for all $s \leq u$ and $i \in \Lambda$. Now Lemma 4.13 tells us that the maps $(\mathbf{X}_{s,u})_{s \leq u}$ are continuous with respect to the product topology. ■

To finish this section we need one more proof.

Proof of Theorem 4.20 It is straightforward to check that $(\mathbf{X}_{s,t})_{s \leq t}$ is a stochastic flow with independent increments. The proof that $(X_s)_{0 \leq s \leq t}$ is a Markov process with semigroup $(P_t)_{t \geq 0}$ now follows in exactly the same way as in the proof of Proposition 2.7, with (4.4) taking the place of (2.6).

The fact that $P_s P_t = P_{s+t}$ follows from the fact that $(\mathbf{X}_{s,t})_{s \leq t}$ is a stochastic flow. Thus, to see that $(P_t)_{t \geq 0}$ is a Feller semigroup, it suffices to show that $(x, t) \mapsto P_t(x, \cdot)$ is a continuous map from $S^\Lambda \times [0, \infty)$ to $\mathcal{M}_1(S^\Lambda)$. In order to do this, it is convenient to use negative times. (Note that we have defined ω to be a Poisson point process on $\mathcal{G} \times \mathbb{R}$, even though for (4.17) we only need points $(m, t) \in \omega$ with $t > 0$.) Since the law of ω is invariant under translations in the time direction, we have (compare (4.18))

$$P_t(x, \cdot) := \mathbb{P}[\mathbf{X}_{-t,0}(x) \in \cdot] \quad (x \in S^\Lambda, t \geq 0).$$

Therefore, in order to prove that $P_{t_n}(x_n, \cdot)$ converges weakly to $P_t(x, \cdot)$ as we let $(x_n, t_n) \rightarrow (x, t)$, it suffices to prove that

$$\mathbf{X}_{0,t_n}(x_n) \xrightarrow[n \rightarrow \infty]{} \mathbf{X}_{0,t}(x) \quad \text{a.s.}$$

as $(x_n, t_n) \rightarrow (x, t)$. Since we equip S^Λ with the product topology, we need to show that

$$\mathbf{X}_{0,t_n}(x_n)(i) \xrightarrow[n \rightarrow \infty]{} \mathbf{X}_{0,t}(x)(i) \quad \text{a.s.}$$

for each $i \in \Lambda$. Let $I := \{s \in \mathbb{R} : \exists (m, s) \in \omega \text{ s.t. } i \in \mathcal{D}(m)\}$ and let $t_- := \sup\{s : s \in I, s \leq t\}$ and $t_+ := \inf\{s : s \in I, s \geq t\}$. Since t is a deterministic time, $t_- < t < t_+$ a.s. Then $\mathbf{X}_{0,t_n}(x_n)(i) = \mathbf{X}_{0,t}(x)(i)$ for all n large enough such that $t_- < t_n < t_+$ and $x_n = x$ on the finite set $\mathcal{R}(\mathbf{X}_{0,t}[i])$, proving the desired a.s. convergence. ■

4.4 Generator construction

Although Theorem 4.20 gives us an explicit way how to construct the Feller semigroup associated with an interacting particle system, it does not tell us very much about its generator. To fill this gap, we need a bit more theory. For any continuous function $f : S^\Lambda \rightarrow \mathbb{R}$ and $i \in \Lambda$, we define

$$\delta f(i) := \sup \{ |f(x) - f(y)| : x, y \in S^\Lambda, x(j) = y(j) \ \forall j \neq i \}.$$

Note that $\delta f(i)$ measures how much $f(x)$ can change if we change x only in the point i . We call δf the *variation* of f .⁹

Lemma 4.27 (Variation of a function) *Let $f \in \mathcal{C}(S^\Lambda)$. Then*

$$|f(x) - f(y)| \leq \sum_{i: x(i) \neq y(i)} \delta f(i) \quad (f \in \mathcal{C}(S^\Lambda), x, y \in S^\Lambda). \quad (4.32)$$

Proof Let n be the number of sites i where x and y differ. Enumerate these sites as $\{i : x(i) \neq y(i)\} = \{i_1, \dots, i_n\}$ or $= \{i_1, i_2, \dots\}$ depending on whether n is finite or not. For $0 \leq k < n + 1$, set

$$z_k(i) := \begin{cases} y(i) & \text{if } i \in \{i_1, \dots, i_k\}, \\ x(i) & \text{otherwise.} \end{cases}$$

If n is finite, then

$$|f(x) - f(y)| \leq \sum_{k=1}^n |f(z_k) - f(z_{k-1})| \leq \sum_{k=1}^n \delta f(i_k)$$

and we are done. If n is infinite, then the same argument gives

$$|f(x) - f(z_m)| \leq \sum_{k=1}^m \delta f(i_k) \quad (m \geq 1).$$

Since $z_m \rightarrow y$ pointwise and f is continuous, (4.32) now follows by letting $m \rightarrow \infty$. ■

We define spaces of functions by

$$\mathcal{C}_{\text{sum}} = \mathcal{C}_{\text{sum}}(S^\Lambda) := \{f \in \mathcal{C}(S^\Lambda) : \sum_i \delta f(i) < \infty\},$$

$$\mathcal{C}_{\text{fin}} = \mathcal{C}_{\text{fin}}(S^\Lambda) := \{f \in \mathcal{C}(S^\Lambda) : \delta f(i) = 0 \text{ for all but finitely many } i\}.$$

⁹This definition is similar to, but different from the more usual definition of the (total) variation of a function of one real variable. With functions of one real variable, the total variation is the maximal sum of all changes in the value of the function as one gradually increases the real variable. For functions on S^Λ , the idea is similar but instead of increasing a real variable we will gradually change a configuration x by modifying its coordinates one by one.

We say that functions in \mathcal{C}_{sum} are of *summable variation*. The next exercise shows that functions in \mathcal{C}_{fin} depend on finitely many coordinates only.

Exercise 4.28 *Let us say that a function $f : S^\Lambda \rightarrow \mathbb{R}$ depends on finitely many coordinates if there exists a finite set $A \subset \Lambda$ and a function $f' : S^A \rightarrow \mathbb{R}$ such that*

$$f((x(i))_{i \in \Lambda}) = f'((x(i))_{i \in A}) \quad (x \in S^\Lambda).$$

Show that each function that depends on finitely many coordinates is continuous, that

$$\mathcal{C}_{\text{fin}}(S^\Lambda) = \{f \in \mathcal{C}(S^\Lambda) : f \text{ depends on finitely many coordinates}\},$$

and that $\mathcal{C}_{\text{fin}}(S^\Lambda)$ is a dense linear subspace of the Banach space $\mathcal{C}(S^\Lambda)$ of all continuous real functions on S^Λ , equipped with the supremum norm.

Lemma 4.29 (Domain of pregenerator) *Assume that the rates $(r_m)_{m \in \mathcal{G}}$ satisfy (4.19). Then, for each $f \in \mathcal{C}_{\text{sum}}(S^\Lambda)$,*

$$\sum_{m \in \mathcal{G}} r_m |f(m(x)) - f(x)| \leq K_0 \sum_{i \in \Lambda} \delta f(i),$$

where K_0 is the constant from (4.19). In particular, for each $f \in \mathcal{C}_{\text{sum}}(S^\Lambda)$, the right-hand side of (4.10) is absolutely summable and Gf is well-defined.

Proof This follows by writing

$$\begin{aligned} \sum_{m \in \mathcal{G}} r_m |f(m(x)) - f(x)| &\leq \sum_{m \in \mathcal{G}} r_m \sum_{i \in \mathcal{D}(m)} \delta f(i) \\ &= \sum_{i \in \Lambda} \delta f(i) \sum_{\substack{m \in \mathcal{G} \\ \mathcal{D}(m) \ni i}} r_m \leq K_0 \sum_{i \in \Lambda} \delta f(i). \end{aligned}$$

■

The following theorem is the main result of the present section.

Theorem 4.30 (Generator construction of particle systems) *Assume that the rates $(r_m)_{m \in \mathcal{G}}$ satisfy (4.15), let $(P_t)_{t \geq 0}$ be the Feller semigroup defined in (4.18) and let G be the linear operator with domain $\mathcal{D}(G) := \mathcal{C}_{\text{sum}}$ defined by (4.10). Then G is closable and its closure \overline{G} is the generator of $(P_t)_{t \geq 0}$. Moreover, \mathcal{C}_{fin} is a core for G .*

To prepare for the proof of Theorem 4.30 we need a few lemmas.

Lemma 4.31 (Generator on local functions) *Under the assumptions of Theorem 4.30, one has $\lim_{t \downarrow 0} t^{-1}(P_t f - f) = Gf$ for all $f \in \mathcal{C}_{\text{fin}}$, where the limit exists in the topology on $\mathcal{C}(S^\Lambda)$.*

Proof Since $f \in \mathcal{C}_{\text{fin}}$, by Exercise 4.28, there exists some finite $A \subset \Lambda$ such that f depends only on the coordinates in A . Let $\mathcal{G}^A := \{m \in \mathcal{G} : \mathcal{D}(m) \cap A \neq \emptyset\}$ denote the set of maps $m \in \mathcal{G}$ that can potentially change the state in A and for $B \supset A$, let $\mathcal{G}^{A,B} := \{m \in \mathcal{G}^A : \mathcal{R}(m[i]) \subset B \ \forall i \in A\}$ denote those maps who only need information from B to update the state in A . For $s \leq t$, we write $\omega_{s,t}^A$ and $\omega_{s,t}^{A,B}$ to denote the sets of Poisson points $(m, s) \in \omega_{s,t}$ with $m \in \mathcal{G}^A$ resp. $m \in \mathcal{G}^{A,B}$. If $\omega_{0,t}^A = \emptyset$, then $f(\mathbf{X}_{0,t}(x)) = f(x)$. Also, if $\omega_{0,t}^{A,B}$ contains a single element (m, s) while $\omega_{0,t}^B$ contains no further elements, then $f(\mathbf{X}_{0,t}(x)) = f(m(x))$. Therefore

$$\begin{aligned} P_t f(x) &= \mathbb{E}[f(\mathbf{X}_{0,t}(x))] = f(x) \mathbb{P}[\omega_{0,t}^A = \emptyset] \\ &\quad + \sum_{m \in \mathcal{G}^{A,B}} f(m(x)) \mathbb{P}[\omega_{0,t}^{A,B} = \omega_{0,t}^B = \{(m, s)\} \text{ for some } 0 < s \leq t] \\ &\quad + \mathbb{E}[f(\mathbf{X}_{0,t}(x)) 1_{\{\omega_{0,t}^A \setminus \omega_{0,t}^{A,B} \neq \emptyset \text{ or } |\omega_{0,t}^B| \geq 2\}}]. \end{aligned}$$

Here $\mathbb{P}[\omega_{0,t}^A = \emptyset] = 1 - e^{-Rt}$ with $R := \sum_{m \in \mathcal{G}^A} r_m$, so

$$f(x) \mathbb{P}[\omega_{0,t}^A = \emptyset] = f(x) - \sum_{m \in \mathcal{G}^A} r_m f(x) + O(t^2) f(x),$$

where $O(t^2)$ is a function such that $\limsup_{t \rightarrow \infty} t^{-2} |O(t^2)| < \infty$. Similarly,

$$\begin{aligned} &\sum_{m \in \mathcal{G}^{A,B}} f(m(x)) \mathbb{P}[\omega_{0,t}^{A,B} = \omega_{0,t}^B = \{(m, s)\} \text{ for some } 0 < s \leq t] \\ &= t \sum_{m \in \mathcal{G}^{A,B}} r_m f(m(x)) + O_B(x, t^2) = t \sum_{m \in \mathcal{G}^A} r_m f(m(x)) + t \varepsilon_B(x) + O_B(x, t^2), \end{aligned}$$

where the error terms satisfy $\limsup_{t \rightarrow \infty} t^{-2} \sup_{x \in S} |O_B(x, t^2)| < \infty$ for each fixed $B \supset A$, and $\lim_{B \uparrow \Lambda} \sup_{x \in S} |\varepsilon_B(x)| = 0$. Similarly

$$\mathbb{P}[\omega_{0,t}^A \setminus \omega_{0,t}^{A,B} \neq \emptyset \text{ or } |\omega_{0,t}^B| \geq 2] = t \varepsilon_B + O_B(t^2),$$

where the term of order t comes from the event that $\omega_{0,t}^A$ contains exactly one element, which is not in $\omega_{0,t}^{A,B}$. Combining our last three formulas, we obtain

$$P_t f(x) - f(x) = t \sum_{m \in \mathcal{G}^A} r_m \{f(m(x)) - f(x)\} + t \varepsilon_B(x) + O_B(x, t^2).$$

It follows that for each fixed $B \supset A$,

$$\limsup_{t \rightarrow \infty} \|t^{-1} (P_t f(x) - f(x)) - G f(x)\| \leq \|\varepsilon_B\|.$$

Since $\|\varepsilon_B\| \rightarrow 0$ as $B \uparrow \Lambda$, the claim of the lemma follows. \blacksquare

Lemma 4.32 (Approximation by local functions) *Assume that the rates $(r_m)_{m \in \mathcal{G}}$ satisfy (4.19). Then for all $f \in \mathcal{C}_{\text{sum}}$ there exist $f_n \in \mathcal{C}_{\text{fin}}$ such that $\|f_n - f\| \rightarrow 0$ and $\|Gf_n - Gf\| \rightarrow 0$.*

Proof Choose finite $\Lambda_n \uparrow \Lambda$, set $\Gamma_n := \Lambda \setminus \Lambda_n$, fix $z \in S^\Lambda$, and for each $x \in S^\Lambda$ define $x_n \rightarrow x$ by

$$x_n(i) := \begin{cases} x(i) & \text{if } i \in \Lambda_n, \\ z(i) & \text{if } i \in \Gamma_n. \end{cases}$$

Fix $f \in \mathcal{C}_{\text{sum}}$ and define $f_n(x) := f(x_n)$ ($x \in S^\Lambda$). Then f_n depends only on the coordinates in Λ_n , hence $f_n \in \mathcal{C}_{\text{fin}}$. Formula (4.32) tells us that for any $x \in S^\Lambda$,

$$|f(x_n) - f(x)| \leq \sum_{i \in \Gamma_n} \delta f(i) \quad (x \in S^\Lambda, n \geq 1)$$

Since $f \in \mathcal{C}_{\text{sum}}$, it follows that

$$\|f_n - f\| \leq \sum_{i \in \Gamma_n} \delta f(i) \xrightarrow{n \rightarrow \infty} 0.$$

Moreover, we observe that

$$\begin{aligned} & |Gf_n(x) - Gf(x)| \\ &= \left| \sum_{m \in \mathcal{G}} r_m (f_n(m(x)) - f_n(x)) - \sum_{m \in \mathcal{G}} r_m (f(m(x)) - f(x)) \right| \\ &\leq \sum_{m \in \mathcal{G}} r_m |f(m(x)_n) - f(x_n) - f(m(x)) + f(x)|. \end{aligned} \quad (4.33)$$

On the one hand, we have

$$\begin{aligned} & |f(m(x)_n) - f(x_n) - f(m(x)) + f(x)| \\ &\leq |f(m(x)_n) - f(x_n)| + |f(m(x)) - f(x)| \leq 2 \sum_{i \in \mathcal{D}(m)} \delta f(i), \end{aligned}$$

while on the other hand, we can estimate the same quantity as

$$\leq |f(m(x)_n) - f(m(x))| + |f(x_n) - f(x)| \leq 2 \sum_{i \in \Gamma_n} \delta f(i).$$

Let $A \subset \Lambda$ be finite. Inserting either of our two estimates into (4.33), depending on whether $\mathcal{D}(m) \cap A \neq \emptyset$ or not, we find that

$$\begin{aligned} \|Gf_n - Gf\| &\leq 2 \sum_{\substack{m \in \mathcal{G} \\ \mathcal{D}(m) \cap A \neq \emptyset}} r_m \sum_{i \in \Gamma_n} \delta f(i) + 2 \sum_{\substack{m \in \mathcal{G} \\ \mathcal{D}(m) \cap A = \emptyset}} r_m \sum_{i \in \mathcal{D}(m)} \delta f(i) \\ &\leq 2K_0|A| \sum_{i \in \Gamma_n} \delta f(i) + 2 \sum_{i \in \Lambda} \delta f(i) \sum_{\substack{m \in \mathcal{G} \\ \mathcal{D}(m) \cap A = \emptyset \\ \mathcal{D}(m) \ni i}} r_m. \end{aligned}$$

It follows that

$$\limsup_{n \rightarrow \infty} \|Gf_n - Gf\| \leq 2 \sum_{i \in \Lambda \setminus A} \delta f(i) \sum_{\substack{m \in \mathcal{G} \\ \mathcal{D}(m) \ni i}} r_m \leq 2K_0 \sum_{i \in \Lambda \setminus A} \delta f(i).$$

Since A is arbitrary, letting $A \uparrow \Lambda$, we see that $\limsup_n \|Gf_n - Gf\| = 0$. ■

Lemma 4.33 (Functions of summable variation) *Under the assumptions of Theorem 4.30, one has*

$$\sum_{i \in \Lambda} \delta P_t f(i) \leq e^{Kt} \sum_{i \in \Lambda} \delta f(i) \quad (t \geq 0, f \in \mathcal{C}_{\text{sum}}(S^\Lambda)),$$

where K is the constant from (4.22). In particular, for each $t \geq 0$, P_t maps $\mathcal{C}_{\text{sum}}(S^\Lambda)$ into itself.

Proof For each $i \in \Lambda$ and $x, y \in S^\Lambda$ such that $x(j) = y(j)$ for all $j \neq i$, we have using (4.32)

$$\begin{aligned} |P_t f(x) - P_t f(y)| &= |\mathbb{E}[f(\mathbf{X}_{0,t}(x))] - \mathbb{E}[f(\mathbf{X}_{0,t}(y))]| \\ &\leq \mathbb{E}[|f(\mathbf{X}_{0,t}(x)) - f(\mathbf{X}_{0,t}(y))|] \\ &\leq \mathbb{E}[\sum_{j: \mathbf{X}_{0,t}(x)(j) \neq \mathbf{X}_{0,t}(y)(j)} \delta f(j)] \\ &= \sum_j \mathbb{P}[\mathbf{X}_{0,t}(x)(j) \neq \mathbf{X}_{0,t}(y)(j)] \delta f(j) \\ &\leq \sum_j \mathbb{P}[(i, 0) \rightsquigarrow (j, t)] \delta f(j). \end{aligned}$$

By Lemma 4.21, it follows that

$$\begin{aligned} \sum_i \delta P_t f(i) &\leq \sum_{ij} \mathbb{P}[(i, 0) \rightsquigarrow (j, t)] \delta f(j) \\ &= \sum_j \mathbb{E}[|\zeta_0^t(\{j\})|] \delta f(j) \leq e^{Kt} \sum_j \delta f(j). \end{aligned}$$

■

Proof of Theorem 4.30 Let H be the full generator of $(P_t)_{t \geq 0}$ and let $\mathcal{D}(H)$ denote its domain. Then Lemma 4.31 shows that $\mathcal{C}_{\text{fin}} \subset \mathcal{D}(H)$ and $Gf = Hf$ for all $f \in \mathcal{C}_{\text{fin}}$. By Lemma 4.32, it follows that $\mathcal{C}_{\text{sum}} \subset \mathcal{D}(H)$ and $Gf = Hf$ for all $f \in \mathcal{C}_{\text{sum}}$. To complete the proof, it suffices to show that \mathcal{C}_{fin} , and hence also the larger \mathcal{C}_{sum} , is a core for H .

We first prove that \mathcal{C}_{sum} is a core for H . We will apply Lemma 4.7. We will show that for each $r > K$, where K is the constant from (4.22), and

for each $f \in \mathcal{C}_{\text{sum}}(S^\Lambda)$, there exists a $p_r \in \mathcal{C}_{\text{sum}}(S^\Lambda)$ that solves the Laplace equation $(r - G)p_r = f$. Since $\mathcal{C}_{\text{sum}}(S^\Lambda)$ is dense in $\mathcal{C}(S^\Lambda)$ by Exercise 4.28, the claim then follows from Lemma 4.7 (ii).

Fix $r > K$ and $f \in \mathcal{C}_{\text{sum}}(S^\Lambda)$. We need to find a $p_r \in \mathcal{C}_{\text{sum}}(S^\Lambda)$ that solves the Laplace equation $(r - G)p_r = f$. In the light of Lemma 4.5 a natural candidate for such a function is

$$p_r := \int_0^\infty e^{-rt} P_t f \, dt$$

and we will show that this p_r indeed satisfies $p_r \in \mathcal{C}_{\text{sum}}(S^\Lambda)$ and $(r - G)p_r = f$. It follows from Theorem 4.8 that $p_r \in \mathcal{D}(H)$ and $(r - H)p_r = f$. Thus, it suffices to show that $p_r \in \mathcal{C}_{\text{sum}}$. To see this, note that if $x(j) = y(j)$ for all $j \neq i$, then

$$\begin{aligned} |p_r(x) - p_r(y)| &= \left| \int_0^\infty e^{-rt} P_t f(x) \, dt - \int_0^\infty e^{-rt} P_t f(y) \, dt \right| \\ &\leq \int_0^\infty e^{-rt} |P_t f(x) - P_t f(y)| \, dt \leq \int_0^\infty e^{-rt} \delta P_t f(i) \, dt, \end{aligned}$$

and therefore, by Lemma 4.33,

$$\sum_i \delta p(i) \leq \int_0^\infty e^{-rt} \sum_i \delta P_t f(i) \, dt \leq \left(\sum_i \delta f(i) \right) \int_0^\infty e^{-rt} e^{Kt} \, dt < \infty,$$

which proves that $p_r \in \mathcal{C}_{\text{sum}}$. This completes the proof that \mathcal{C}_{sum} is a core for H , i.e., the closure of $G|_{\mathcal{C}_{\text{sum}}}$ is H . By Lemma 4.32, the closure of $G|_{\mathcal{C}_{\text{fin}}}$ contains $G|_{\mathcal{C}_{\text{sum}}}$, so we see that \mathcal{C}_{fin} is also a core for H . ■

We conclude this section with the following lemma, that is sometimes useful.

Lemma 4.34 (Differentiation of semigroup) *Assume that the rates $(r_m)_{m \in \mathcal{G}}$ satisfy (4.15), let $(P_t)_{t \geq 0}$ be the Feller semigroup defined in (4.18) and let G be the linear operator with domain $\mathcal{D}(G) := \mathcal{C}_{\text{sum}}(S^\Lambda)$ defined by (4.10). Then, for each $f \in \mathcal{C}_{\text{sum}}(S^\Lambda)$, $t \mapsto P_t f$ is a continuously differentiable function from $[0, \infty)$ to $\mathcal{C}(S^\Lambda)$ satisfying $P_0 f = f$, $P_t f \in \mathcal{C}_{\text{sum}}(S^\Lambda)$, and $\frac{\partial}{\partial t} P_t f = G P_t f$ for each $t \geq 0$.*

Proof This is a direct consequence of Proposition 4.4, Lemma 4.33, and Theorem 4.30. A direct proof based on our definition of $(P_t)_{t \geq 0}$ (not using Hille-Yosida theory) is also possible, but quite long and technical. ■

Some bibliographical remarks

Theorem 4.30 is similar to Liggett's [Lig85, Theorem I.3.9], but there are also some differences. Liggett does not write his generators in terms of local maps, but in terms of local probability kernels, similar to (4.9). This way of writing the generator is more general and sometimes (for example, for stochastic Ising models) more natural than our approach using local maps. It is worth noting that Liggett's construction, like ours, depends on a clever way of writing the generator that is in general not unique.

Unlike our Theorem 4.20, Liggett does not give an explicit construction of his interacting particle systems using Poisson point sets, but instead gives a direct proof that the closure of G generates a Feller semigroup $(P_t)_{t \geq 0}$, and then invokes the abstract result Theorem 4.1 about Feller processes to prove the existence of a corresponding Markov process with cadlag sample paths. Later in his book, he does use explicit Poisson constructions for some systems, such as the contact process. He does not actually prove that these Poisson constructions yield the same process as the generator construction, but apparently finds this self-evident. (Equivalence of the two constructions follows from our Theorem 4.30 but alternatively can also be proved by approximation with finite systems, using approximation results such as [Lig85, Cor. I.3.14].)

Liggett's [Lig85, Theorem I.3.9] allows for the case that the local state space S is a (not necessarily finite) compact metrizable space. This is occasionally convenient. For example, this allows one to construct voter models with infinitely many types, where at time zero, the types $(X_0(i))_{i \in \Lambda}$ are i.i.d. and uniformly distributed on $S = [0, 1]$. For simplicity, we have restricted ourselves to finite local state spaces.

4.5 Ergodicity

Luckily, our efforts in the previous chapter are not wasted on knowing only that the systems we are interested in exist, but actually allow us to prove something interesting about these systems as well.

If X is a Markov process with state space E and transition probabilities $(P_t)_{t \geq 0}$, then by definition, an *invariant law* of X is a probability measure ν on E such that

$$\nu P_t = \nu \quad (t \geq 0).$$

This says that if we start the process in the initial law $\mathbb{P}[X_0 \in \cdot] = \nu$, then $\mathbb{P}[X_t \in \cdot] = \nu$ for all $t \geq 0$. As a consequence, one can construct a

stationary¹⁰ process $(X_t)_{t \in \mathbb{R}}$ such that (compare (4.3))

$$\mathbb{P}[X_u \in \cdot \mid (X_s)_{-\infty < s \leq t}] = P_{u-t}(X_t, \cdot) \quad \text{a.s.} \quad (t \leq u), \quad (4.34)$$

and $\mathbb{P}[X_t \in \cdot] = \nu$ for all $t \in \mathbb{R}$. Conversely, the existence of such a stationary Markov process implies that the law at any time $\nu := \mathbb{P}[X_t \in \cdot]$ must be an invariant law. For this reason, invariant laws are sometimes also called *stationary laws*.

Theorem 4.35 (Ergodicity) *Let X be an interacting particle system with state space of the form S^Λ and generator G of the form (4.10), and assume that the rates $(r_m)_{m \in \mathcal{G}}$ satisfy (4.15).*

(a) *Assume that the constant K from (4.22) satisfies $K < 0$. Then the process ζ defined in (4.21) satisfies*

$$\lim_{s \rightarrow -\infty} \zeta_s^u(\{i\}) = \emptyset \quad \text{a.s.} \quad (i \in \Lambda, u \in \mathbb{R}). \quad (4.35)$$

(b) *Assume that the process ζ defined in (4.21) satisfies (4.35). Then the interacting particle system X has a unique invariant law ν , and*

$$\mathbb{P}^x[X_t \in \cdot] \xrightarrow[t \rightarrow \infty]{} \nu \quad (x \in S^\Lambda). \quad (4.36)$$

Proof Part (a) is immediate from Lemma 4.21. If (4.35) holds, then by Lemma 4.26, the a.s. limit

$$X_t(i) := \lim_{s \rightarrow -\infty} \mathbf{X}_{s,t}(z)(i) \quad (i \in \Lambda, t \in \mathbb{R}) \quad (4.37)$$

exists and does not depend on the choice of a point $z \in S^\Lambda$, since the set $\zeta_s^t(\{i\})$ of lattice points whose value at time s is relevant for $\mathbf{X}_{s,t}(z)(i)$ is empty for s sufficiently small. As a result, (4.37) unambiguously defines a stationary process $X = (X_t)_{t \in \mathbb{R}}$. We claim that X is Markov with respect to the transition probabilities $(P_t)_{t \geq 0}$ in the sense of (4.34). Indeed, for almost every trajectory $(x_s)_{-\infty < s \leq t}$ with respect to the law of $(X_s)_{-\infty < s \leq t}$, we have

$$\begin{aligned} & \mathbb{P}[X_u \in \cdot \mid (X_s)_{-\infty < s \leq t} = (x_s)_{-\infty < s \leq t}] \\ &= \mathbb{P}\left[\lim_{s \rightarrow -\infty} \mathbf{X}_{t,u} \circ \mathbf{X}_{s,t}(z) \in \cdot \mid (X_s)_{-\infty < s \leq t} = (x_s)_{-\infty < s \leq t}\right] \\ &\stackrel{1}{=} \mathbb{P}[\mathbf{X}_{t,u}(X_t) \in \cdot \mid (X_s)_{-\infty < s \leq t} = (x_s)_{-\infty < s \leq t}] \\ &= \mathbb{P}[\mathbf{X}_{t,u}(x_t) \in \cdot \mid (X_s)_{-\infty < s \leq t} = (x_s)_{-\infty < s \leq t}] \\ &\stackrel{2}{=} \mathbb{P}[\mathbf{X}_{t,u}(x_t) \in \cdot] = P_{u-t}(x_t, \cdot), \end{aligned}$$

¹⁰Recall that a process $(X_t)_{t \in \mathbb{R}}$ is stationary if for each $s \in \mathbb{R}$, it is equal in distribution to $(X'_t)_{t \in \mathbb{R}}$ defined as $X'_t := X_{t-s}$ ($t \in \mathbb{R}$).

where in step 1 we have used the continuity of the map $\mathbf{X}_{t,u}$ (which is proved in Lemma 4.26) and in step 2 we have used that the random variables $\mathbf{X}_{t,u}$ and $(X_s)_{-\infty < s \leq t}$ are independent, since they are functions of the restriction of the Poisson set ω to the disjoint sets $\mathcal{G} \times (t, u]$ and $\mathcal{G} \times (-\infty, t]$, respectively. By stationarity,

$$\nu := \mathbb{P}[X_t \in \cdot] \quad (t \in \mathbb{R})$$

does not depend on $t \in \mathbb{R}$, and since X is Markov this defines an invariant law ν . Since

$$\mathbb{P}^x[X_t \in \cdot] = \mathbb{P}[\mathbf{X}_{-t,0}(x)(i) \in \cdot]$$

and since by (4.37), we have

$$\mathbf{X}_{-t,0}(x) \xrightarrow[t \rightarrow \infty]{} X_0 \quad \text{a.s.} \quad (x \in S^\Lambda)$$

with respect to the topology of pointwise convergence, we conclude that (4.36) holds. \blacksquare

Remark The condition (4.35) in fact implies that there exists a unique cadlag process $(X_t)_{t \in \mathbb{R}}$ such that (compare (4.14))

$$X_t = \mathbf{m}_t^\omega(X_{t-}) \quad (t \in \mathbb{R}). \quad (4.38)$$

Indeed, by the definition of the maps $(\mathbf{X}_{s,u})_{s \leq u}$ in (4.16), the equation (4.38) is equivalent to $X_t = \mathbf{X}_{s,t}(X_s)$ ($s \leq t$), which by (4.37) implies that $(X_t)_{t \in \mathbb{R}}$ must be the stationary process constructed in the proof of Theorem 4.35.

We note that (4.36) says that if we start the process in an arbitrary initial state x , then the law at time t converges weakly¹¹ as $t \rightarrow \infty$ to the invariant law ν . This property is often described by saying that the interacting particle system is *ergodic*. Indeed, this implies that the corresponding stationary process $(X_t)_{t \in \mathbb{R}}$ is ergodic in the usual sense of that word, i.e., the σ -field of events that are invariant under translations in time is trivial. The converse conclusion cannot be drawn, however, so the traditional way of describing (4.36) as “ergodicity” is a bit of a bad habit.

We have split Theorem 4.35 into a part (a) and (b) since the condition (4.35) is sometimes satisfied even when the constant K from (4.22) is positive. Indeed, we will later see that for the contact process, the condition (4.35) is sharp but the condition $K < 0$ is not. In Exercise 5.12 below, we will calculate the constant K for the contact process and deduce that this process is ergodic for small values of the infection rate.

¹¹Here weak convergence is of course w.r.t. our topology on S^Λ , i.e., w.r.t. the product topology.

Theorem 4.35 is similar, but not identical to [Lig85, Thm I.4.1]. For Theorem 4.35 (a) and (b) to be applicable, one needs to be able to express the generator in terms of local maps such that the constant K from (4.22) is negative. For [Lig85, Thm I.4.1], one needs to express the generator in a convenient way in terms of local transition kernels. For certain problems, the latter approach is more natural and [Lig85, Thm I.4.1] yields sharper estimates for the regime where ergodicity holds.

4.6 Application to the Ising model

The Ising model with Glauber dynamics has been introduced in Section 1.4. So far, we have not shown how to represent the generator of this interacting particle system in terms of local maps. In the present section, we will fill this gap. As an application of the theory developed so far, we will then show that the Ising model with Glauber dynamics is well-defined for all values of its parameter, and ergodic for β sufficiently small. Our construction will also prepare for the next chapter, where we discuss monotone interacting particle systems, by showing that the Ising model with Glauber dynamics can be represented in monotone maps.

We recall from Section 1.4 that the Ising model with Glauber dynamics on a graph (Λ, E) is the interacting particle system with state space $\{-1, +1\}^\Lambda$ and dynamics such that

$$\text{site } i \text{ flips to the value } \sigma \text{ with rate } r_i^\sigma(x) := \frac{e^{\beta N_{x,i}(\sigma)}}{e^{\beta N_{x,i}(+1)} + e^{\beta N_{x,i}(-1)}},$$

where

$$N_{x,i}(\sigma) := \sum_{j \in \mathcal{N}_i} 1_{\{x(j) = \sigma\}} \quad (\sigma \in \{-1, +1\})$$

denotes the number of neighbors of i that have the spin value σ . For each $i \in \Lambda$, let K_i^β denote the local probability kernel on $\{-1, +1\}^\Lambda$ defined as

$$K_i^\beta(x, y) := \begin{cases} r_i^\sigma(x) & \text{if } y = m_i^\sigma(x) \quad (\sigma \in \{-1, +1\}), \\ 0 & \text{otherwise,} \end{cases}$$

where $m_i^\sigma(x)$ is defined in (1.14). Then the generator (1.13) of the Ising model takes the form

$$G_{\text{Ising}} f = \sum_{i \in \Lambda} \{K_i^\beta f - f\}, \quad (4.39)$$

which is an expression of the form (4.1) but not of the form (4.2). To find an expression for G_{Ising} in terms of local maps as in (4.2), it suffices to find a random mapping representation for the kernels K_i^β . This needs some preparations. Let

$$M_{x,i} := N_{x,i}(+) - N_{x,i}(-) = \sum_{j \in \mathcal{N}_i} x(j)$$

denote the *local magnetization* in the neighborhood \mathcal{N}_i of i . Since $N_{x,i}(+) + N_{x,i}(-) = |\mathcal{N}_i|$, we can rewrite the rate of flipping to the spin value $+1$ as

$$\begin{aligned} r_i^+(x) &= \frac{e^{\beta N_{x,i}(+)}}{e^{\beta N_{x,i}(+)} + e^{\beta N_{x,i}(-)}} = \frac{e^{\beta(|\mathcal{N}_i| + M_{x,i})/2}}{e^{\beta(|\mathcal{N}_i| + M_{x,i})/2} + e^{\beta(|\mathcal{N}_i| - M_{x,i})/2}} \\ &= \frac{e^{\frac{1}{2}\beta M_{x,i}}}{e^{\frac{1}{2}\beta M_{x,i}} + e^{-\frac{1}{2}\beta M_{x,i}}} = \frac{1}{2} \left(1 + \frac{e^{\frac{1}{2}\beta M_{x,i}} - e^{-\frac{1}{2}\beta M_{x,i}}}{e^{\frac{1}{2}\beta M_{x,i}} + e^{-\frac{1}{2}\beta M_{x,i}}} \right) \\ &= \frac{1}{2} (1 + \tanh(\frac{1}{2}\beta M_{x,i})). \end{aligned}$$

Similarly, the rate of flipping to -1 is $r_i^-(x) = \frac{1}{2}(1 - \tanh(\frac{1}{2}\beta M_{x,i}))$.

For (mainly notational) simplicity, let us assume that each site i has the same number of neighbors in the graph (Λ, E) , so that the size of the neighborhood

$$N := |\mathcal{N}_i| \quad (i \in \Lambda)$$

does not depend on $i \in \Lambda$. Then $M_{x,i}$ takes values in $\{-N, -N+2, \dots, N\}$. We observe that the function $z \mapsto \frac{1}{2}(1 + \tanh(\frac{1}{2}\beta z))$ is increasing (see Figure 4.2). Inspired by this, for $L = -N, -N+2, \dots, N$, we define local maps $m_{i,L}^\pm$ by

$$\begin{aligned} m_{i,L}^+(x)(j) &:= \begin{cases} +1 & \text{if } j = i \text{ and } M_{x,i} \geq L, \\ x(j) & \text{otherwise.} \end{cases} \\ m_{i,L}^-(x)(j) &:= \begin{cases} -1 & \text{if } j = i \text{ and } M_{x,i} \leq L, \\ x(j) & \text{otherwise,} \end{cases} \end{aligned} \tag{4.40}$$

and we try a generator of the form

$$G_{\text{Ising}} f(x) = \sum_{i \in \Lambda} \sum_{\sigma \in \{-, +\}} \sum_{L \in \{-N, -N+2, \dots, N\}} r_{i,L}^\sigma \{f(m_{i,L}^\sigma(x)) - f(x)\}, \tag{4.41}$$

where $r_{i,L}^\sigma \geq 0$ are probabilities that need to be chosen in such a way that

$$K_i^\beta(x, y) = \sum_{\sigma \in \{-, +\}} \sum_{L \in \{-N, -N+2, \dots, N\}} r_{i,L}^\sigma 1\{m_{i,L}^\sigma(x) = y\} \tag{4.42}$$

is a random mapping representation of the kernel K_i^β . Let us fix x^-, x^+ such that $x^-(i) = -1$, $x^+(i) = +1$, and $x^-(j) = x^+(j)$ for all $j \neq i$. Setting $x = x^-$ and $y = x^+$ in (4.42) yields the equation

$$r_i^+(x) = \sum_{L=-N}^{M_{x,i}} r_{i,L}^+$$

Similarly, setting $x = x^+$ and $y = x^-$ in (4.42) yields the equation

$$r_i^-(x) = \sum_{L=M_{x,i}}^N r_{i,L}^-.$$

From this, we see that setting

$$\begin{aligned} r_{i,L}^+ &:= \begin{cases} \frac{1}{2}(1 + \tanh(-\frac{1}{2}\beta N)) & \text{if } L = -N, \\ \frac{1}{2}\tanh(\frac{1}{2}\beta L) - \frac{1}{2}\tanh(\frac{1}{2}\beta(L-2)) & \text{otherwise,} \end{cases} \\ r_{i,L}^- &:= \begin{cases} \frac{1}{2}(1 - \tanh(\frac{1}{2}\beta N)) & \text{if } L = N, \\ \frac{1}{2}\tanh(\frac{1}{2}\beta L) - \frac{1}{2}\tanh(\frac{1}{2}\beta(L+2)) & \text{otherwise,} \end{cases} \end{aligned} \quad (4.43)$$

has the effect that the generator in (4.41) equals the one in (4.39).




Figure 4.2: Definition of the rates $r_{i,L}^+$ from (4.43). In this example $N = 6$ and $\beta = 0.4$.

Theorem 4.36 (Existence and ergodicity of the Ising model) *Consider an Ising model with Glauber dynamics on a countable graph Λ in which each lattice point i has exactly $|\mathcal{N}_i| = N$ neighbors, i.e., the Markov process X with state space $\{-1, +1\}^\Lambda$ and generator G_{Ising} given by (4.41). Then, for each $\beta \geq 0$, the closure of G_{Ising} generates a Feller semigroup. Moreover, for each β such that*

$$e^{\beta N} < \frac{N}{N-1}, \quad (4.44)$$

the Markov process with generator $\overline{G}_{\text{Ising}}$ has a unique invariant law ν , and the process started in an arbitrary initial state x satisfies

$$\mathbb{P}^x[X_t \in \cdot] \xrightarrow[t \rightarrow \infty]{} \nu \quad (x \in \{-1, +1\}^\Lambda).$$

Proof We use the representation (4.41). We observe that

$$\mathcal{D}(m_{i,L}^\pm) = \{i\}$$

is the set of lattice points whose spin value can be changed by the map $m_{i,L}^\pm$. The set of lattice points that are $m_{i,L}^\pm$ -relevant for i is given by

$$\mathcal{R}(m_{i,L}^\sigma[i]) = \begin{cases} \emptyset & \text{if } \sigma = +, L = -N \quad \text{or} \quad \sigma = -, L = N, \\ \mathcal{N}_i & \text{otherwise.} \end{cases}$$

Here we have used that $-N \leq M_{x,i} \leq N$ holds always, so $m_{-N}^+(x)(i) = +1$ and $m_N^-(x)(i) = -1$ regardless of what x is. On the other hand, in all other cases, the value of each lattice point $j \in \mathcal{N}_i$ can potentially make a difference for the outcome $m_{i,L}^\pm(x)(i)$.

By Theorem 4.30, to conclude that the closure of G_{Ising} generates a Feller semigroup, it suffices to check (4.15), which in our case says that

$$\sup_{i \in \Lambda} \sum_{\sigma \in \{-, +\}} \sum_{L \in \{-N, -N+2, \dots, N\}} r_{i,L}^\sigma (|\mathcal{R}(m_{i,L}^\sigma[i])| + 1)$$

should be finite. Since $\sum_L r_{i,L}^\sigma \leq \frac{1}{2}(1 + \tanh(\frac{1}{2}\beta N)) \leq 1$ and $|\mathcal{R}(m_{i,L}^\sigma[i])| \leq |\mathcal{N}_i| = N$, this expression is $\leq 2(N+1) < \infty$ regardless of the value of β .

To prove ergodicity for β small enough, we apply Theorem 4.35. We calculate the constant K from (4.22). By the symmetry between minus and plus spins,

$$\begin{aligned} K &= 2 \sum_{L \in \{-N, -N+2, \dots, N\}} r_{i,L}^+ (|\mathcal{R}(m_{i,L}^+[i])| - 1) \\ &= -2r_{i,-N}^+ + 2 \sum_{L \in \{-N+2, \dots, N\}} r_{i,L}^+ (N-1) \\ &= -\left(1 + \tanh(-\tfrac{1}{2}\beta N)\right) + \left(\tanh(\tfrac{1}{2}\beta N) - \tanh(-\tfrac{1}{2}\beta N)\right)(N-1), \end{aligned}$$

which is negative if and only if

$$\begin{aligned}
1 + \tanh(-\tfrac{1}{2}\beta N) &> (\tanh(\tfrac{1}{2}\beta N) - \tanh(-\tfrac{1}{2}\beta N))(N-1) \\
\Leftrightarrow 1 + \frac{e^{-\frac{1}{2}\beta N} - e^{\frac{1}{2}\beta N}}{e^{\frac{1}{2}\beta N} + e^{-\frac{1}{2}\beta N}} &> \left(\frac{e^{\frac{1}{2}\beta N} - e^{-\frac{1}{2}\beta N}}{e^{\frac{1}{2}\beta N} + e^{-\frac{1}{2}\beta N}} - \frac{e^{-\frac{1}{2}\beta N} - e^{\frac{1}{2}\beta N}}{e^{\frac{1}{2}\beta N} + e^{-\frac{1}{2}\beta N}} \right) (N-1) \\
\Leftrightarrow 2e^{-\frac{1}{2}\beta N} &> 2(e^{\frac{1}{2}\beta N} - e^{-\frac{1}{2}\beta N})(N-1) \\
\Leftrightarrow \frac{e^{-\frac{1}{2}\beta N}}{e^{\frac{1}{2}\beta N} - e^{-\frac{1}{2}\beta N}} &> N-1 \quad \Leftrightarrow \frac{1}{e^{\beta N} - 1} > N-1 \\
\Leftrightarrow e^{\beta N} - 1 < \frac{1}{N-1} &\quad \Leftrightarrow e^{\beta N} < \frac{N}{N-1},
\end{aligned}$$

which is condition (4.44). ■

4.7 Further results

In the present section we collect a number of technical results of a general nature that will be needed in later chapters. On a first reading, readers are advised to skip the present section and refer back to specific results when the need arises. The only result of the present section that is perhaps of some intrinsic value is Theorem 4.41 which together with Corollary 4.42 below implies that the transition probabilities of interacting particle systems on infinite lattices can be approximated by those on finite lattices, something that we have been using implicitly when doing simulations.

Let E be a compact metrizable space. By definition, a collection of functions $\mathcal{H} \subset \mathcal{C}(E)$ is *distribution determining* if

$$\mu f = \nu f \quad \forall f \in \mathcal{H} \quad \text{implies} \quad \mu = \nu.$$

We say that \mathcal{H} *separates points* if for all $x, y \in E$ such that $x \neq y$, there exists an $f \in \mathcal{H}$ such that $f(x) \neq f(y)$. We say that \mathcal{H} is *closed under products* if $f, g \in \mathcal{H}$ implies $fg \in \mathcal{H}$.

Lemma 4.37 (Application of Stone-Weierstrass) *Let E be a compact metrizable space. Assume that $\mathcal{H} \subset \mathcal{C}(E)$ separates points and is closed under products. Then \mathcal{H} is distribution determining.*

Proof If $\mu f = \nu f$ for all $f \in \mathcal{H}$, then we can add the constant function 1 to \mathcal{H} and retain this property. In a next step, we can add all linear combinations of functions in \mathcal{H} to the set \mathcal{H} ; by the linearity of the integral, it will then

still be true that $\mu f = \nu f$ for all $f \in \mathcal{H}$. But now \mathcal{H} is an algebra that separates points and vanishes nowhere, so by the Stone-Weierstrass theorem, \mathcal{H} is dense in $\mathcal{C}(E)$. If $f_n \in \mathcal{H}$, $f \in \mathcal{C}(E)$, and $\|f_n - f\|_\infty \rightarrow 0$, then $\mu f_n \rightarrow \mu f$ and likewise for ν , so we conclude that $\mu f = \nu f$ for all $f \in \mathcal{C}(E)$. If $A \subset E$ is a closed set, then the function $f(x) := d(x, A)$ is continuous, where d is a metric generating the topology on E and $d(x, A) := \inf_{y \in A} d(x, y)$ denotes the distance of x to A . Now the functions $f_n := 1 \wedge n f$ are also continuous and $f_n \uparrow 1_{A^c}$, so by the continuity of the integral with respect to increasing sequences we see that $\mu(O) = \nu(O)$ for every open set $O \subset E$. Since the open sets are closed under intersections, it follows that $\mu(A) = \nu(A)$ for every element A of the σ -algebra generated by the open sets, i.e., the Borel- σ -field $\mathcal{B}(E)$. ■

Lemma 4.38 (Weak convergence) *Let E be a compact metrizable space. Assume that $\mu_n \in \mathcal{M}_1(E)$ have the property that $\lim_{n \rightarrow \infty} \mu_n f$ exists for all $f \in \mathcal{H}$, where $\mathcal{H} \subset \mathcal{C}(E)$ is distribution determining. Then there exists a $\mu \in \mathcal{M}_1(E)$ such that $\mu_n \Rightarrow \mu$.*

Proof By Prohorov's theorem, the space $\mathcal{M}_1(E)$, equipped with the topology of weak convergence, is compact. Therefore, to prove the statement, it suffices to show that the sequence μ_n has not more than one cluster point, i.e., it suffices to show that if μ, μ' are subsequential limits, then $\mu' = \mu$. Clearly, μ, μ' must satisfy $\mu' f = \mu f$ for all $f \in \mathcal{H}$, so the claim follows from the assumption that \mathcal{H} is distribution determining. ■

Lemma 4.39 (Continuous probability kernels) *Let E be a compact metrizable space and let K be a continuous probability kernel on E . Then, for any $\mu_n, \mu \in \mathcal{M}_1(E)$ and $f_n, f \in \mathcal{C}(E)$,*

$$\begin{aligned} \mu_n \xrightarrow[n \rightarrow \infty]{} \mu \quad \text{implies} \quad \mu_n K \xrightarrow[n \rightarrow \infty]{} \mu K \\ \text{and} \quad \|f_n - f\|_\infty \xrightarrow[n \rightarrow \infty]{} 0 \quad \text{implies} \quad \|K f_n - K f\|_\infty \xrightarrow[n \rightarrow \infty]{} 0. \end{aligned}$$

Proof Since K is a continuous probability kernel, its associated operator maps the space $\mathcal{C}(E)$ into itself, so $\mu_n \Rightarrow \mu$ implies that $\mu_n(Kf) \Rightarrow \mu(Kf)$ for all $f \in \mathcal{C}(E)$, or equivalently $(\mu_n K)f \Rightarrow (\mu K)f$ for all $f \in \mathcal{C}(E)$, i.e., the measures $\mu_n K$ converge weakly to μ .

The second statement follows from the linearity and monotonicity of K and the fact that $K1 = 1$, which together imply that $\|K f_n - K f\|_\infty \leq \|f_n - f\|_\infty$. ■

Lemma 4.40 (Long-time limits) *Let E be a compact metrizable space and let $(P_t)_{t \geq 0}$ be the transition probabilities of a Feller process in E . Let $\mu \in \mathcal{M}_1(E)$ and assume that*

$$\mu P_t \xrightarrow[t \rightarrow \infty]{} \nu$$

for some $\nu \in \mathcal{M}_1(E)$. Then ν is an invariant law of the Feller process with transition probabilities $(P_t)_{t \geq 0}$.

Proof Using Lemma 4.39, this follows by writing

$$\nu P_t = \left(\lim_{s \rightarrow \infty} \mu P_s \right) P_t = \lim_{s \rightarrow \infty} \mu P_s P_t = \lim_{s \rightarrow \infty} \mu P_{s+t} = \nu.$$

■

The following theorem follows from [Kal97, Thm 17.25], where it is moreover shown that the condition (4.45) implies convergence in distribution of the associated Feller processes, viewed as random variables taking values in the space $\mathcal{D}_E[0, \infty)$ of cadlag paths with values in E . Note that in (4.45) below, \rightarrow (of course) means convergence in the topology we have defined on $\mathcal{C}(E)$, i.e., convergence w.r.t. the supremumnorm.

Theorem 4.41 (Limits of semigroups) *Let E be a compact metrizable space and let G_n, G be generators of Feller processes in E . Assume that there exists a linear operator on $\mathcal{C}(E)$ such that $\bar{A} = G$ and*

$$\forall f \in \mathcal{D}(A) \exists f_n \in \mathcal{D}(G_n) \text{ such that } f_n \rightarrow f \text{ and } G_n f_n \rightarrow A f. \quad (4.45)$$

Then the Feller semigroups $(P_t^n)_{t \geq 0}$ and $(P_t)_{t \geq 0}$ with generators G_n and G , respectively, satisfy

$$\sup_{t \in [0, T]} \|P_t^n f - P_t f\|_\infty \xrightarrow[n \rightarrow \infty]{} 0 \quad (f \in \mathcal{C}(E), T < \infty).$$

Moreover, if $\mu_n, \mu \in \mathcal{M}_1(E)$, then

$$\mu_n \xrightarrow[n \rightarrow \infty]{} \mu \text{ implies } \mu_n P_t^n \xrightarrow[n \rightarrow \infty]{} \mu P_t \quad (t \geq 0).$$

We note that in the case of interacting particle systems, Theorem 4.30 implies the following.

Corollary 4.42 (Convergence of particle systems) *Let S be a finite set and let Λ be countable. Let G_n, G be generators of interacting particle systems in S^Λ and assume that G_n, G can be written in the form (4.10) with rates satisfying (4.15). Assume moreover that*

$$\|G_n f - G f\|_\infty \xrightarrow[n \rightarrow \infty]{} 0 \quad (f \in \mathcal{C}_{\text{fin}}(S^\Lambda)).$$

Then the generators G_n, G satisfy (4.45).

Theorem 4.41 has the following useful consequence.

Proposition 4.43 (Limits of invariant laws) *Let E be a compact metrizable space and let G_n, G be generators of Feller processes in E satisfying (4.45). Let $\nu_n, \nu \in \mathcal{M}_1(E)$ and assume that for each n , the measure ν_n is an invariant law of the Feller process with generator G_n . Then $\nu_n \Rightarrow \nu$ implies that ν is an invariant law of the Feller process with generator G .*

Proof Using Theorem 4.41, this follows simply by observing that

$$\nu P_t = \lim_{n \rightarrow \infty} \nu_n P_t^n = \lim_{n \rightarrow \infty} \nu_n = \nu$$

for each $t \geq 0$. ■

Chapter 5

Monotonicity

5.1 The stochastic order

We recall that if S and T are partially ordered sets, then a function $f : S \rightarrow T$ is called *monotone* iff $x \leq y$ implies $f(x) \leq f(y)$. In particular, this definition also applies to real-valued functions (where we equip \mathbb{R} with the well-known order). If the local state space S of an interacting particle system is partially ordered, then we equip the product space with the *product order*

$$x \leq y \quad \text{iff} \quad x(i) \leq y(i) \quad \forall i \in \Lambda.$$

Many well-known interacting particle systems use the local state space $S = \{0, 1\}$, which is of course equipped with a natural order $0 \leq 1$. Often, it is useful to prove comparison results, that say that two interacting particle systems X and Y can be coupled in such a way that $X_t \leq Y_t$ for all $t \geq 0$. Here X and Y may be different systems, started in the same initial state, or also two copies of the same interacting particle system, started in initial states such that $X_0 \leq Y_0$.

The following theorem gives necessary and sufficient conditions for it to be possible to couple two random variables X and Y such that $X \leq Y$. A *coupling* of two random variables X and Y , in the most general sense of the word, is *a way to construct X and Y together on one underlying probability space* $(\Omega, \mathcal{F}, \mathbb{P})$. More precisely, if X and Y are random variables defined on different underlying probability spaces, then a coupling of X and Y is a pair of random variables (X', Y') defined on one underlying probability space $(\Omega, \mathcal{F}, \mathbb{P})$, such that X' is equally distributed with X and Y' is equally distributed with Y . Equivalently, since the laws of X and Y are all we really care about, we may say that a *coupling* of two probability laws μ, ν defined on measurable spaces (E, \mathcal{E}) and (F, \mathcal{F}) , respectively, is a probability measure

ρ on the product space $(E \times F, \mathcal{E} \otimes \mathcal{F})$ such that the first marginal of ρ is μ and its second marginal is ν .

Theorem 5.1 (Stochastic order) *Let S be a finite partially ordered set, let Λ be a countable set, and let μ, ν be probability laws on S^Λ . Then the following statements are equivalent:*

- (i) $\int \mu(dx) f(x) \leq \int \nu(dx) f(x) \quad \forall \text{ monotone } f \in \mathcal{C}(S^\Lambda),$
- (ii) $\int \mu(dx) f(x) \leq \int \nu(dx) f(x) \quad \forall \text{ monotone } f \in B(S^\Lambda),$
- (iii) *It is possible to couple random variables X, Y with laws $\mu = P[X \in \cdot]$ and $\nu = P[Y \in \cdot]$ in such a way that $X \leq Y$.*

Proof The implication (iii) \Rightarrow (ii) is easy: if X and Y are coupled such that $X \leq Y$ and f is monotone, then

$$\mathbb{E}[f(Y)] - \mathbb{E}[f(X)] = \mathbb{E}[f(Y) - f(X)] \geq 0,$$

since $f(Y) - f(X) \geq 0$ a.s. The implication (ii) \Rightarrow (i) is trivial.

For the nontrivial implication (i) \Rightarrow (iii) we refer to [Lig85, Theorem II.2.4]. For finite spaces, a nice intuitive proof based on the max flow min cut theorem can be found in [Pre74]. The theorem holds for more general spaces than spaces of the form S^Λ . For example, it holds also for \mathbb{R}^n ; see [KKO77]. ■

If two probability laws μ, ν satisfy the equivalent conditions of Theorem 5.1, then we say that μ and ν are *stochastically ordered* and we write¹ $\mu \leq \nu$. Clearly $\mu \leq \nu \leq \rho$ implies $\mu \leq \rho$. In light of this, the next lemma shows that the stochastic order is a bona fide partial order on $\mathcal{M}_1(S^\Lambda)$.

Lemma 5.2 (Monotone functions are distribution determining) *Let S be a finite partially ordered set and let Λ be countable. Then the set $\{f \in \mathcal{C}(S^\Lambda) : f \text{ is monotone}\}$ is distribution determining. In particular, $\mu \leq \nu$ and $\mu \geq \nu$ imply $\mu = \nu$.*

Proof Since the finite-dimensional distributions uniquely determine a probability measure on S^Λ , it suffices to prove the statement for finite Λ . In view of this, it suffices to show that if S is a finite partially ordered set, then the space of all monotone functions $f : S \rightarrow \mathbb{R}$ is distribution determining.

¹This notation may look a bit confusing at first sight, since, if μ, ν are probability measures on any measurable space (ω, \mathcal{F}) , then one might interpret $\mu \leq \nu$ in a pointwise sense, i.e., in the sense that $\mu(A) \leq \nu(A)$ for all $A \in \mathcal{F}$. In practice, this does not lead to confusion, since pointwise inequality for probability measures is a very uninteresting property. Indeed, it is easy to check that probability measures μ, ν satisfy $\mu \leq \nu$ in a pointwise sense if and only if $\mu = \nu$.

By definition, an *increasing subset* of S is a set $A \subset S$ such that $A \ni x \leq y$ implies $y \in A$. If A is increasing, then its indicator function 1_A is monotone, so it suffices to show that $\{1_A : A \text{ increasing}\}$ is distribution determining. By Lemma 4.37, it suffices to show that this class separates points and is closed under products.

If $x \neq y$, then either $x \notin \{z : z \geq y\}$ or $y \notin \{z : z \geq x\}$, so $\{1_A : A \text{ increasing}\}$ separates points. If A, B are increasing, then so is $A \cap B$, so by the fact that $1_A 1_B = 1_{A \cap B}$ we see that $\{1_A : A \text{ increasing}\}$ is closed under products. ■

We continue to consider spaces of the form S^Λ where S is a finite partially ordered set and Λ is countable. In particular, since Λ can be a set with only one element, this includes arbitrary finite partially ordered sets. By definition, a probability kernel K on S^Λ is *monotone* if it satisfies the following equivalent conditions. Note that in (i) below, \leq denotes the stochastic order. The equivalence of (i) and (ii) is a trivial consequence of Theorem 5.1.

- (i) $K(x, \cdot) \leq K(y, \cdot)$ for all $x \leq y$.
- (ii) Kf is monotone whenever $f \in \mathcal{C}(S^\Lambda)$ is monotone.

We note that if K is monotone, then

$$\mu \leq \nu \quad \text{implies} \quad \mu K \leq \nu K. \quad (5.1)$$

Indeed, for each monotone $f \in B(S^\Lambda)$, the function Kf is also monotone and hence $\mu \leq \nu$ implies that $\mu Kf \leq \nu Kf$.

Recall from (2.3) that a *random mapping representation* of a probability kernel K is a random map M such that

$$K(x, \cdot) = \mathbb{P}[M(x) \in \cdot] \quad \forall x. \quad (5.2)$$

We say that K can be represented in the class of monotone maps, or that K is *monotonically representable*, if there exists a random monotone map M such that (5.2) holds. We recall from Section 2.4 that when a Markov generator G is written in the form

$$Gf(x) = \sum_{m \in \mathcal{G}} r_m \{f(m(x)) - f(x)\}, \quad (5.3)$$

then we call (5.3) a *random mapping representation* of G . If the set \mathcal{G} can be chosen such that all maps $m \in \mathcal{G}$ are monotone, then we say that G is *monotonically representable*.

Lemma 5.3 (Monotone representability) *Each monotonically representable probability kernel is monotone. If the generator of an interacting particle system is monotonically representable, then, for each $t \geq 0$, the transition probability P_t is a monotonically representable probability kernel.*

Proof If a probability kernel K can be written in the form (5.2) with M a random monotone map, then for each $x \leq y$, the random variables $M(x)$ and $M(y)$ are coupled such that $M(x) \leq M(y)$ a.s., so their laws are stochastically ordered as $K(x, \cdot) \leq K(y, \cdot)$. Since this holds for all $x \leq y$, the kernel K is monotone.

Given a random mapping representation of the form (5.3) of the generator G of an interacting particle system, we can construct a stochastic flow $(\mathbf{X}_{s,t})_{s \leq t}$ as in (4.16) based on a Poisson set $\omega \subset \mathcal{G} \times \mathbb{R}$. If all maps $m \in \mathcal{G}$ are monotone, then the random maps $\mathbf{X}_{s,t} : S^\Lambda \rightarrow S^\Lambda$ are also monotone, since they are pointwise defined as the concatenation of finitely many maps from \mathcal{G} . It follows that

$$P_t(x, \cdot) = \mathbb{P}[\mathbf{X}_{0,t}(x) \in \cdot]$$

is a representation of P_t in terms of the random monotone map $\mathbf{X}_{0,t}$, so P_t is monotonically representable. ■

We say that an interacting particle system is *monotone* if its transition kernels are monotone probability kernels, and we say that it is *monotonically representable* if its generator is monotonically representable. Somewhat surprisingly, it turns out that for probability kernels, “monotonically representable” is a strictly stronger concept than being “monotone”. See [FM01] for an example of a probability kernel on $\{0, 1\}^2$ that is monotone but not monotonically representable. Nevertheless, it turns out that (almost) all monotone interacting particle systems that one encounters in practice are also monotonically representable.

The following maps are examples of monotone maps:

- The voter map \mathbf{vot}_{ij} defined in (1.4).
- The branching map \mathbf{bra}_{ij} defined in (1.6).
- The death map \mathbf{death}_i defined in (1.7).
- The coalescing random walk map \mathbf{rw}_{ij} defined in (1.20).
- The exclusion map \mathbf{excl}_{ij} defined in (1.23).
- The cooperative branching map \mathbf{coop}_{ij} defined in (1.25).

- The maps $m_{i,L}^{\pm}$ defined in (4.40).

As a result, the following interacting particle systems are monotonically representable (and hence, in particular, monotone):

- The voter model with generator as in (1.5).
- The contact process with generator as in (1.8).
- The ferromagnetic Ising model with Glauber dynamics, since its generator can be written as in (4.41).
- The biased voter model with generator as in (1.17).
- Systems of coalescing random walks with generator as in (1.21).
- The exclusion process with generator as in (1.24).
- Systems with cooperative branching and coalescence as in Figure 1.11.

On the other hand, the following maps are *not* monotone:

- The annihilating random walk map \mathbf{ann}_{ij} defined in (1.22).
- The killing map \mathbf{kill}_{ij} defined in (1.26).

Examples of interacting particle systems that are not monotone² are:

- The antiferromagnetic Ising model with Glauber dynamics.
- “Rebellious” voter models as in (1.18).
- Systems of annihilating random walks.
- The biased annihilating branching process of [Sud97, Sud99].

²Note that the fact that a given interacting particle system is represented in maps that are not monotone does not prove that the system is not monotone. Indeed, it is conceivable that the same system can also be monotonely represented.

5.2 The upper and lower invariant laws

In the present section, we assume that the local state space is $S = \{0, 1\}$, which covers all examples of monotone interacting particle systems mentioned in the previous section. We use the symbols $\underline{0}$ and $\underline{1}$ to denote the states in S^Λ that are identically 0 or 1, respectively. Below, $\delta_{\underline{0}}$ denotes the delta measure at the configuration that is identically 0, so $\delta_{\underline{0}}P_t$ denotes the law at time t of the process started in $X_0(i) = 0$ a.s. ($i \in \Lambda$).

Theorem 5.4 (Upper and lower invariant laws) *Let X be an interacting particle system with state space of the form $\{0, 1\}^\Lambda$ and transition probabilities $(P_t)_{t \geq 0}$. Assume that X is monotone. Then there exist invariant laws $\underline{\nu}$ and $\bar{\nu}$ such that*

$$\delta_{\underline{0}}P_t \xrightarrow[t \rightarrow \infty]{} \underline{\nu} \quad \text{and} \quad \delta_{\underline{1}}P_t \xrightarrow[t \rightarrow \infty]{} \bar{\nu}.$$

If ν is any other invariant law, then $\underline{\nu} \leq \nu \leq \bar{\nu}$.

The invariant laws $\underline{\nu}$ and $\bar{\nu}$ from Theorem 5.4 are called *lower* and *upper invariant law*, respectively. Before we give the proof of Theorem 5.4, we start with two preparatory lemmas.

Lemma 5.5 (Equal mean) *Let μ, ν be probability laws on $\{0, 1\}^S$ such that $\mu \leq \nu$ and*

$$\int \mu(dx) x(i) \geq \int \nu(dx) x(i) \quad (i \in \Lambda).$$

Then $\mu = \nu$.

Proof By Theorem 5.1, we can couple random variables with laws $\mathbb{P}[X \in \cdot] = \mu$ and $\mathbb{P}[Y \in \cdot] = \nu$ in such a way that $X \leq Y$. Now $\mathbb{E}[X(i)] \geq \mathbb{E}[Y(i)]$ implies $\mathbb{E}[Y(i) - X(i)] \leq 0$. Since $Y(i) - X(i) \geq 0$ a.s., it follows that $X(i) = Y(i)$. In particular, if this holds for all $i \in \Lambda$, then $\mu = \nu$. ■

Lemma 5.6 (Monotone convergence of probability laws) *Let $(\nu_n)_{n \geq 0}$ be a sequence of probability laws on $\{0, 1\}^\Lambda$ that are stochastically ordered as $\nu_k \leq \nu_{k+1}$ ($k \geq 0$). Then there exists a probability law ν on $\{0, 1\}^\Lambda$ such that $\nu_n \Rightarrow \nu$, i.e., the ν_n 's converge weakly to ν .*

Proof Since $\nu_n f$ increases to a finite limit for each monotone $f \in \mathcal{C}(\{0, 1\}^\Lambda)$, this is an immediate consequence of Lemmas 5.2 and 4.38. ■

Proof of Theorem 5.4 By symmetry, it suffices to prove the statement for $\underline{\nu}$. Since $\underline{0}$ is the lowest possible state, for each $t \geq 0$, we trivially have

$$\delta_{\underline{0}} \leq \delta_{\underline{0}}P_t$$

By (5.1), this implies that

$$\delta_0 P_s \leq \delta_0 P_t P_s = \delta_0 P_{t+s} \quad (s, t \geq 0),$$

which shows that $t \mapsto \delta_0 P_t$ is nondecreasing with respect to the stochastic order. By Lemma 5.6, each monotone sequence of probability laws has a weak limit, so there exists a probability law $\underline{\nu}$ on $\{0, 1\}^\Lambda$ such that

$$\delta_0 P_t \xrightarrow[t \rightarrow \infty]{} \underline{\nu}.$$

It follows from Lemma 4.40 that $\underline{\nu}$ is an invariant law.

To complete the proof of the theorem, we observe that if ν is any other invariant law, then, by (5.1), for any monotone $f \in \mathcal{C}(\{0, 1\}^\Lambda)$,

$$\delta_0 \leq \nu \quad \Rightarrow \quad \delta_0 P_t \leq \nu P_t = \nu \quad (t \geq 0).$$

Letting $t \rightarrow \infty$, it follows that $\underline{\nu} f \leq \nu f$ for all monotone $f \in \mathcal{C}(\{0, 1\}^\Lambda)$, which by Theorem 5.1 implies that $\underline{\nu} \leq \nu$. \blacksquare

Theorem 5.7 (Ergodicity of monotone systems) *Let X be a monotone interacting particle system with state space $\{0, 1\}^\Lambda$ and upper and lower invariant laws $\underline{\nu}$ and $\bar{\nu}$. If*

$$\int \underline{\nu}(dx) x(i) = \int \bar{\nu}(dx) x(i) \quad \forall i \in \Lambda, \quad (5.4)$$

then X has a unique invariant law $\nu := \underline{\nu} = \bar{\nu}$ and is ergodic in the sense that

$$\mathbb{P}^x[X_t \in \cdot] \xrightarrow[t \rightarrow \infty]{} \nu \quad (x \in \{0, 1\}^\Lambda).$$

On the other hand, if (5.4) does not hold, then X has at least two invariant laws.

Proof By Lemma 5.5, (5.4) is equivalent to the condition that $\underline{\nu} = \bar{\nu}$. It is clear that if $\underline{\nu} \neq \bar{\nu}$, then X has at least two invariant laws and ergodicity cannot hold. On the other hand, by Theorem 5.4, any invariant law ν satisfies $\underline{\nu} \leq \nu \leq \bar{\nu}$, so if $\underline{\nu} = \bar{\nu}$, then $\nu = \underline{\nu} = \bar{\nu}$.

To complete the proof, we must show that $\underline{\nu} = \bar{\nu} =: \nu$ implies $\delta_x P_t \Rightarrow \nu$ as $t \rightarrow \infty$ for all $x \in \{0, 1\}^\Lambda$. Since

$$\delta_0 P_t f \leq \delta_x P_t f \leq \delta_1 P_t f$$

for all monotone $f \in \mathcal{C}(\{0, 1\}^\Lambda)$, we see that

$$\underline{\nu} f \leq \liminf_{t \rightarrow \infty} P_t f \leq \limsup_{t \rightarrow \infty} P_t f \leq \bar{\nu} f$$

The claim now follows from Lemmas 4.38 and 5.2. ■

To state the final result of this section, we need a bit of theory. We observe that for any interacting particle system, the set \mathcal{I} of all invariant laws is a compact, convex subset of $\mathcal{M}_1(S^\Lambda)$. Indeed, if μ and ν are invariant laws and $p \in [0, 1]$, then clearly

$$(p\mu + (1-p)\nu)P_t = p\mu P_t + (1-p)\nu P_t = p\mu + (1-p)\nu \quad (t \geq 0),$$

proving that $p\mu + (1-p)\nu$ is an invariant law. The fact that \mathcal{I} is closed follows from Proposition 4.43. Since $\mathcal{M}_1(S^\Lambda)$ is compact, \mathcal{I} is also compact.

By definition, an element $\nu \in \mathcal{I}$ is called *extremal* if it cannot be written as a nontrivial convex combination of other elements of \mathcal{I} , i.e.,

$$\nu = p\nu_1 + (1-p)\nu_2 \quad (0 < p < 1, \nu_1, \nu_2 \in \mathcal{I}) \quad \text{implies} \quad \nu_1 = \nu_2 = \nu.$$

We let

$$\mathcal{I}_e := \{\nu \in \mathcal{I} : \nu \text{ is an extremal element of } \mathcal{I}\}.$$

Since \mathcal{I} is compact and convex, Choquet's theorem implies that each invariant law ν can be written as

$$\nu = \int \rho_\nu(d\mu)\mu,$$

where ρ_ν is a probability measure on \mathcal{I}_e . In practice, it happens quite often³ that \mathcal{I}_e is a finite set.⁴ In this case, Choquet's theorem simply says that each invariant law is a convex combination of the extremal invariant laws, i.e., each invariant law is of the form

$$\nu = \sum_{\mu \in \mathcal{I}_e} p(\mu)\mu,$$

where $(p(\mu))_{\mu \in \mathcal{I}_e}$ are nonnegative constants, summing up to one. In view of this, we are naturally interested in finding all extremal invariant laws of a given interacting particle system.

Lemma 5.8 (The lower and upper invariant law are extremal) *Let X be a monotone interacting particle system with state space $\{0, 1\}^\Lambda$ and upper and lower invariant laws $\underline{\nu}$ and $\bar{\nu}$. Then $\underline{\nu}$ and $\bar{\nu}$ are extremal invariant laws of X .*

³The voter model in dimensions $d \geq 3$ is a counterexample. The Ising model in dimensions $d \geq 3$ is also a counterexample, although for the Ising model, it is still true that $\underline{\nu}$ and $\bar{\nu}$ are the only extremal invariant measures that are moreover translation invariant.

⁴This may, however, be quite difficult to prove!

Proof By symmetry, it suffices to prove the statement for $\bar{\nu}$. Imagine that

$$\bar{\nu} = p\nu_1 + (1-p)\nu_2 \quad \text{for some } 0 < p < 1, \nu_1, \nu_2 \in \mathcal{I}.$$

By Theorem 5.4, for each monotone $f \in B(\{0,1\}^\Lambda)$, one has $\nu_1 f \leq \bar{\nu} f$ and $\nu_2 f \leq \bar{\nu} f$. Since

$$p(\bar{\nu} f - \nu_1 f) + (1-p)(\bar{\nu} f - \nu_2 f) = 0,$$

it follows that $\bar{\nu} f = \nu_1 f = \nu_2 f$. Since this holds for each monotone f , we conclude (by Lemma 5.2) that $\bar{\nu} = \nu_1 = \nu_2$. \blacksquare

Exercise 5.9 Let X be an interacting particle system with state space $\{0,1\}^\Lambda$ and generator G . Assume that G has a random mapping representation in terms of monotone maps and let $(\mathbf{X}_{s,t})_{s \leq t}$ be the corresponding stochastic flow as in (4.16). Show that the a.s. limits

$$\left. \begin{aligned} \underline{X}_t &:= \lim_{s \rightarrow -\infty} \mathbf{X}_{s,t}(\underline{0}), \\ \overline{X}_t &:= \lim_{s \rightarrow -\infty} \mathbf{X}_{s,t}(\underline{1}) \end{aligned} \right\} \quad (t \in \mathbb{R})$$

define stationary Markov processes $(\underline{X}_t)_{t \in \mathbb{R}}$ and $(\overline{X}_t)_{t \in \mathbb{R}}$ whose invariant laws

$$\underline{\nu} = \mathbb{P}[\underline{X}_t \in \cdot] \quad \text{and} \quad \bar{\nu} = \mathbb{P}[\overline{X}_t \in \cdot] \quad (t \in \mathbb{R})$$

are the lower and upper invariant law of X , respectively. Show that (5.4) implies that

$$\lim_{s \rightarrow -\infty} \mathbf{X}_{s,t}(x) = \underline{X}_t = \overline{X}_t \quad \text{a.s.} \quad (x \in \{0,1\}^\Lambda, t \in \mathbb{R}).$$

5.3 The contact process

We recall the definition of the contact process from (1.8). Since both the branching and death map are monotone, this is a monotonically representable interacting particle system, so by Theorem 5.4, it has a lower and upper invariant law $\underline{\nu}$ and $\bar{\nu}$. Since $\mathbf{bra}_{ij}(\underline{0}) = \underline{0}$ and $\mathbf{death}_i(\underline{0}) = \underline{0}$ for each $i, j \in \Lambda$, the all-zero configuration $\underline{0}$ is a trap for the contact process, so $\delta_{\underline{0}} P_t = \delta_{\underline{0}}$ for all $t \geq 0$ and hence

$$\underline{\nu} = \delta_{\underline{0}}.$$

Therefore, by Theorem 5.7, the contact process is ergodic if and only if the function

$$\theta(\lambda) := \int \bar{\nu}(dx) x(i) \quad (i \in \mathbb{Z}^d) \quad (5.5)$$

satisfies $\theta(\lambda) = 0$. Here λ denotes the infection rate and we stick to the convention to take the recovery rate δ (1.8) equal to 1. We note that by translation invariance, for the model on \mathbb{Z}^d (either nearest-neighbor or range R), the density $\int \bar{\nu}(dx) x(i)$ of the upper invariant law does not depend on $i \in \mathbb{Z}^d$. For reasons that will become clear in the next chapter, $\theta(\lambda)$ is actually the same as the survival probability started from a single occupied site, i.e., this is the function in Figure 1.4.

By definition, we say that a probability law μ on $\{0, 1\}^\Lambda$ is *nontrivial* if

$$\mu(\{\underline{0}\}) = 0,$$

i.e., if μ gives zero probability to the all-zero configuration.

Lemma 5.10 (Nontriviality of the upper invariant law) *For the contact process, if $\bar{\nu} \neq \delta_{\underline{0}}$, then $\bar{\nu}$ is nontrivial.*

Proof We can always write $\bar{\nu} = (1 - p)\delta_{\underline{0}} + p\mu$ where $p \in [0, 1]$ and μ is a nontrivial law. By assumption, $\bar{\nu} \neq \delta_{\underline{0}}$, so $p > 0$. Since $\bar{\nu}$ and $\delta_{\underline{0}}$ are invariant laws, μ must be an invariant law too. By Lemma 5.8, $\bar{\nu}$ cannot be written as a nontrivial convex combination of other invariant laws, so we conclude that $p = 1$. ■

Proposition 5.11 (Monotonicity in the infection rate) *Let $\bar{\nu}_\lambda$ denote the upper invariant law of the contact process with infection rate λ . Then $\lambda \leq \lambda'$ implies $\bar{\nu}_\lambda \leq \bar{\nu}_{\lambda'}$. In particular, the function $\lambda \mapsto \theta(\lambda)$ is nondecreasing.*

Proof Let X and X' be contact processes started in the initial state $X_0 = \underline{1} = X'_0$ and with infection rates λ and λ' . It suffices to prove that X and X' can be coupled such that $X_t \leq X'_t$ for all $t \geq 0$.

We use a Poisson construction, based on the random mapping representation (1.8). We write $\mathcal{G} = \mathcal{G}_{\text{bra}} \cup \mathcal{G}_{\text{death}}$ where

$$\mathcal{G}_{\text{bra}} := \{\mathbf{bra}_{ij} : (i, j) \in \mathcal{E}^d\} \quad \text{and} \quad \mathcal{G}_{\text{death}} := \{\mathbf{death}_i : i \in \mathbb{Z}^d\}.$$

Then X can be constructed as in Theorem 4.20 from a Poisson point set ω on

$$\mathcal{G} \times \mathbb{R} = (\mathcal{G}_{\text{bra}} \cup \mathcal{G}_{\text{death}}) \times \mathbb{R},$$

with intensity measure ρ_λ given by

$$\rho_\lambda(\{m\} \times A) := \begin{cases} \lambda \ell(A) & \text{if } m \in \mathcal{G}_{\text{bra}}, \\ \ell(A) & \text{if } m \in \mathcal{G}_{\text{death}}, \end{cases} \quad (A \in \mathcal{B}(\mathbb{R})),$$

where ℓ denotes the Lebesgue measure. Likewise, X' can be constructed from a Poisson point set ω' with intensity $\rho_{\lambda'}$. We claim that we can couple ω and ω' in such a way that the latter has more branching incidents, and the same death incidents as ω . This can be done as follows. Let ω'' be a Poisson point set on $\mathcal{G} \times \mathbb{R}$, independent of ω , with intensity measure $\rho'' := \rho_{\lambda'} - \rho_{\lambda}$, i.e.,

$$\rho''(\{m\} \times A) := \begin{cases} (\lambda' - \lambda)\ell(A) & \text{if } m \in \mathcal{G}_{\text{bra}}, \\ 0 & \text{if } m \in \mathcal{G}_{\text{death}}, \end{cases} \quad (A \in \mathcal{B}(\mathbb{R})).$$

Since the sum of two independent Poisson sets yields another Poisson set, setting

$$\omega' := \omega + \omega''$$

defines a Poisson point set with intensity $\rho_{\lambda'}$. We observe that

$$\begin{aligned} x \leq x' & \text{ implies } \mathbf{bra}_{ij}(x) \leq \mathbf{bra}_{ij}(x'), \\ x \leq x' & \text{ implies } \mathbf{death}_i(x) \leq \mathbf{death}_i(x'), \\ x \leq x' & \text{ implies } x \leq \mathbf{bra}_{ij}(x'). \end{aligned}$$

The first two statements just say that the maps \mathbf{bra}_{ij} and \mathbf{death}_i are monotone. The third statement says that if we apply a branching map only to the larger configuration x' , then the order between x and x' is preserved.

Since ω' has the same branching and death incidents as ω , plus some extra branching incidents, we conclude that the stochastic flows $(\mathbf{X}_{s,t})_{s \leq t}$ and $(\mathbf{X}'_{s,t})_{s \leq t}$ constructed from ω and ω' satisfy

$$x \leq x' \text{ implies } \mathbf{X}_{s,t}(x) \leq \mathbf{X}'_{s,t}(x') \quad (s \leq t).$$

In particular, setting $X_t := \mathbf{X}_{0,t}(1)$ and $X'_t := \mathbf{X}'_{0,t}(1)$ yields the desired coupling between X and X' . \blacksquare

Exercise 5.12 *Let X be a contact process on a graph Λ where each site i has exactly $|\mathcal{N}_i| = N$ neighbors. Calculate the constant K from (4.22) and apply Theorem 4.35 to conclude that*

$$\lambda N < 1 \text{ implies } \bar{\nu} = \delta_{\underline{0}}.$$

In Chapter 7, we will prove that $\theta(\lambda) > 0$ for λ sufficiently large.

5.4 Other examples

The Ising model with Glauber dynamics

We have seen in (4.41) that the generator of the Ising model with Glauber dynamics is monotonically representable, so by Theorem 5.4,⁵ it has a lower and upper invariant law $\underline{\nu}$ and $\bar{\nu}$. We let

$$m_*(\beta) := \int \bar{\nu}(dx) x(i),$$

which is independent of i if the processes has some translation invariant structure (like the nearest neighbor or range R processes on \mathbb{Z}^d). For reasons that cannot be explained here, this function is actually the same as the one defined in (1.15), i.e., this is the *spontaneous magnetization* of the Ising model, see Figure 1.6. By the symmetry between $+1$ and -1 spins, we clearly have

$$\int \underline{\nu}(dx) x(i) = -m_*(\beta).$$

By Theorem 4.36, we have

$$e^{\beta N} < \frac{N}{N-1} \quad \text{implies} \quad \underline{\nu} = \bar{\nu},$$

from which we conclude that $m_*(\beta) = 0$ for β sufficiently small,

The function $\beta \mapsto m_*(\beta)$ is nondecreasing, but this cannot be proved with the sort of techniques used in Proposition 5.11. The lower and upper invariant laws of the Ising model with Glauber dynamics are infinite volume Gibbs measures, and much of the analysis of the Ising model is based on this fact. In fact, the Ising model with Glauber dynamics is just one example of an interacting particle system that has these Gibbs measures as its invariant laws. In general, interacting particle systems with this property are called stochastic Ising models, and the Gibbs measures themselves are simply called the Ising model. We refer to [Lig85, Chapter IV] for an exposition of this material. In particular, in [Lig85, Thm IV.3.14], it is shown that for the nearest-neighbor model on \mathbb{Z}^2 , one has $m_*(\beta) > 0$ for β sufficiently large.

The voter model

Consider a voter model with local state space $S = \{0, 1\}$. Since the voter maps vot_{ij} from (1.4) are monotone, the voter model is monotonically rep-

⁵The difference between the local state space $\{-1, 1\}$ of the Ising model and $\{0, 1\}$ of Theorem 5.4 is of course entirely notational.

resentable. Since both the constant configurations $\underline{0}$ and $\underline{1}$ are traps,

$$\underline{\nu} = \delta_{\underline{0}} \quad \text{and} \quad \bar{\nu} = \delta_{\underline{1}},$$

so we conclude (recall Theorem 5.7) that the voter model is never ergodic. For the model on \mathbb{Z}^d , it is proved in [Lig85, Thm V.1.8] that if $d = 1, 2$, then $\delta_{\underline{0}}$ and $\delta_{\underline{1}}$ are the only extremal invariant laws. On the other hand, in dimensions $d \geq 3$, the set \mathcal{I}_e of extremal invariant laws is of the form $\{\nu_p : p \in [0, 1]\}$ where the invariant measure ν_n has intensity $\int \nu_p(dx) x(i) = p$. We will give a partial proof of these statements in Chapter 6.

5.5 Exercises

Exercise 5.13 Give an example of two probability measures μ, ν on a set of the form $\{0, 1\}^\Lambda$ that satisfy

$$\int \mu(dx) x(i) \leq \int \nu(dx) x(i) \quad (i \in \Lambda),$$

but that are not stochastically ordered as $\mu \leq \nu$.

Exercise 5.14 Let $(X_t^\lambda)_{t \geq 0}$ denote the contact process with infection rate λ (and death rate one), started in $X_0^\lambda = 1$. Apply Corollary 4.42 to prove that for each fixed $t \geq 0$, the function

$$\theta_t(\lambda) := \mathbb{P}[\mathbf{X}_{0,t}^\lambda(1)(i) = 1] \quad (5.6)$$

depends continuously on λ . Use this to conclude that the function $\theta(\lambda)$ from (5.5) is right-continuous. Hint: Use that the decreasing limit of continuous functions is upper semi-continuous.

For the next exercise, let us define a *double death* map

$$\text{death}_{ij}x(k) := \begin{cases} 0 & \text{if } k \in \{i, j\}, \\ x(k) & \text{otherwise.} \end{cases} \quad (5.7)$$

Recall the branching map bra_{ij} defined in (1.6), the death map death_i defined in (1.7), and the cooperative branching map coop_{ij} defined in (1.25). Consider the cooperative branching process X with values in $\{0, 1\}^\mathbb{Z}$ with generator

$$\begin{aligned} G_X f(x) = & \lambda \sum_{i \in \mathbb{Z}} \sum_{\sigma \in \{-1, +1\}} \{f(\text{coop}_{i, i+\sigma, i+2\sigma}x) - f(x)\} \\ & + \sum_{i \in \mathbb{Z}} \{f(\text{death}_i x) - f(x)\}, \end{aligned}$$

and the contact process with double deaths Y with generator

$$G_Y f(y) = \lambda \sum_{i \in \mathbb{Z}} \sum_{\sigma \in \{-1, +1\}} \{f(\text{bra}_{i, i+\sigma} y) - f(y)\} \\ + \sum_{i \in \mathbb{Z}} \{f(\text{death}_{i, i+1} y) - f(y)\},$$

Exercise 5.15 Let X be the process with cooperative branching defined above and set

$$X_t^{(2)}(i) := 1_{\{X_t(i)=1=X_t(i+1)\}} \quad (i \in \mathbb{Z}, t \geq 0).$$

Show that X can be coupled to a contact process with double deaths Y (with the same parameter λ) in such a way that

$$Y_0 \leq X_0^{(2)} \quad \text{implies} \quad Y_t \leq X_t^{(2)} \quad (t \geq 0).$$

Exercise 5.16 Show that a system $(X_t)_{t \geq 0}$ of annihilating random walks can be coupled to a system $(Y_t)_{t \geq 0}$ of coalescing random walks such that

$$X_0 \leq Y_0 \quad \text{implies} \quad X_t \leq Y_t \quad (t \geq 0).$$

Note that the annihilating random walks are not a monotone particle system.

Exercise 5.17 Let X be a system of branching and coalescing random walks with generator

$$G_X f(x) = \frac{1}{2}b \sum_{i \in \mathbb{Z}} \sum_{\sigma \in \{-1, +1\}} \{f(\text{bra}_{i, i+\sigma} x) - f(x)\} \\ + \frac{1}{2} \sum_{i \in \mathbb{Z}} \sum_{\sigma \in \{-1, +1\}} \{f(\text{rw}_{i, i+\sigma} x) - f(x)\},$$

and let Y be a system of coalescing random walks with positive drift, with generator

$$G_Y f(y) = \frac{1}{2}(1+b) \sum_{i \in \mathbb{Z}} \{f(\text{rw}_{i, i+1} y) - f(y)\} \\ + \frac{1}{2} \sum_{i \in \mathbb{Z}} \{f(\text{rw}_{i, i-1} y) - f(y)\}.$$

Show that X and Y can be coupled such that

$$Y_0 \leq X_0 \quad \text{implies} \quad Y_t \leq X_t \quad (t \geq 0).$$

Exercise 5.18 Let $d < d'$ and identify \mathbb{Z}^d with the subset of $\mathbb{Z}^{d'}$ consisting of all $(i_1, \dots, i_{d'})$ with $(i_{d+1}, \dots, i_{d'}) = (0, \dots, 0)$. Let X and X' denote the

nearest-neighbor contact processes on \mathbb{Z}^d and $\mathbb{Z}^{d'}$, respectively, with the same infection rate λ . Show that X and X' can be coupled such that

$$X_0(i) \leq X'_0(i) \quad (i \in \mathbb{Z}^d)$$

implies

$$X_t(i) \leq X'_t(i) \quad (t \geq 0, i \in \mathbb{Z}^d).$$

Prove the same when X is the nearest-neighbor process and X' is the range R process (both on \mathbb{Z}^d).

Chapter 6

Duality

6.1 Introduction

Imagine that we are studying an interacting particle system $(X_t)_{t \geq 0}$ that is constructed from a graphical representation as in Theorem 4.20 and we are interested in the state $X_u(i)$ of the system at a given site $i \in \Lambda$ and time u . Then it is natural to try to follow, backwards in time, all paths of potential influence that end at the space-time point (i, u) and try to figure out how the space-time point (i, u) got its type. We already used this idea in Chapter 4 where we showed that under the summability condition (4.22), the set $\zeta_{u-t}^u(\{i\})$ of all starting points of paths of potential influence that end at (i, u) grows at most exponentially in t , which formed the basis of the proof of Theorem 4.20. In Lemma 4.26, we moreover showed that the random map $\mathbf{X}_{u-t,u}[i]$, that tells us how $X_u(i)$ depends on X_{u-t} , depends only on coordinates in $\zeta_{u-t}^u(\{i\})$.

In the present chapter, we will show that the *backward in time process*

$$(\mathbf{X}_{u-t,u}[i])_{t \geq 0} \tag{6.1}$$

is in fact a Markov chain with a countable state space, i.e., a continuous-time Markov chain. Clearly, a lot of information about a given interacting particle system is hidden in its associated backward in time process. In general, the backward in time process is rather difficult to study since the maps $\mathbf{X}_{u-t,u}[i]$ quickly become complicated as t increases. We will see, however, that there are special classes of interacting particle systems (additive and cancellative systems) for which $\mathbf{X}_{u-t,u}[i]$ has a simple form. This naturally leads to the concept of *pathwise duality* and more generally *duality*.

To demonstrate the usefulness of these ideas, in Section 6.9, we will use the backward in time process to show that the voter model clusters in dimensions $d = 1, 2$, but not in dimensions $d \geq 3$. In Section 6.10, we use

the self-duality of the contact process to prove that for processes with some sort of translation invariant structure, the upper invariant law is the limit law started from any nontrivial translation invariant initial law, and we will show that this in turn implies that the function $\theta(\lambda)$ from (5.5) is continuous everywhere, except possibly at the critical point. Finally, in Section 6.11, we use duality to show that for a model with a mixture of voter model and contact process duality, the critical points associated with survival and nontriviality of the upper invariant law coincide.

6.2 Additive systems duality

Let Λ be a countable set. For any configuration $x \in \{0, 1\}^\Lambda$, we let

$$|x| := \sum_{i \in \Lambda} x(i)$$

denote the number of ones. Let Δ be another countable set. By definition, a map $m : \{0, 1\}^\Lambda \rightarrow \{0, 1\}^\Delta$ is *additive* iff

- (i) $m(\underline{0}) = \underline{0}$,
- (ii) $m(x \vee y) = m(x) \vee m(y) \quad (x, y \in \mathcal{S}(\Lambda)).$

It is easy to see that additive maps are monotone. Examples of additive maps are:

- The voter map \mathbf{vot}_{ij} defined in (1.4).
- The branching map \mathbf{bra}_{ij} defined in (1.6).
- The death map \mathbf{death}_i defined in (1.7).
- The coalescing random walk map \mathbf{rw}_{ij} defined in (1.20).
- The exclusion map \mathbf{excl}_{ij} defined in (1.23).

On the other hand, the following maps are monotone, but not additive:

- The cooperative branching map \mathbf{coop}_{ijk} defined in (1.25).
- The maps $m_{i,L}^\pm$ used to construct the Ising model with Glauber dynamics in (4.41).

An interacting particle system is called *additive* if its generator can be represented in additive local maps. Examples of additive particle systems are:

- The voter model with generator as in (1.5).
- The contact process with generator as in (1.8).
- The biased voter model with generator as in (1.17).
- Systems of coalescing random walks with generator as in (1.21).
- The exclusion process with generator as in (1.24).

There is a useful graphical way to describe an additive map $m : \{0, 1\}^\Lambda \rightarrow \{0, 1\}^\Lambda$, that works as follows:


- For each $i, j \in \Lambda$ with $i \neq j$ such that $m(1_{\{i\}})(j) = 1$, we draw an arrow from i to j .
- For each $i \in \Lambda$ such that $m(1_{\{i\}})(i) = 0$, we draw a blocking symbol \blacksquare at i .

The following lemma says that additive local maps are fully described by their arrows and blocking symbols.


Lemma 6.1 (Additive local maps) *An additive map $m : \{0, 1\}^\Lambda \rightarrow \{0, 1\}^\Lambda$ is local if and only if its description involves only finitely many arrows and blocking symbols. If m is an additive local map and $x \in \{0, 1\}^\Lambda$, then $m(x)(j) = 1$ if and only if at least one of the following conditions is satisfied:*

- there is an $i \in \Lambda$ such that $x(i) = 1$ and there is an arrow from i to j ,*
- $x(j) = 1$ and there is no blocking symbol at j .*


Although the proof of Lemma 6.1 is quite simple, we will postpone it until we have set up a more systematic framework for proving this and related claims. In terms of arrows and blocking symbols, the branching map \mathbf{bra}_{ij} , the death map \mathbf{death}_i , the voter map \mathbf{vot}_{ij} , the coalescing random walk map \mathbf{rw}_{ij} , and the exclusion map \mathbf{excl}_{ij} look as follows:



We use our conventions of representing additive maps in terms of arrows and blocking symbols to depict the graphical representation of an additive interacting particle system in a more suggestive way. In Figure 4.1, we drew the graphical representation of a contact process in the following fashion:



With our new conventions, the same graphical representation looks as follows:



It is easy to see that the contatenation of two additive maps is again additive. As a result, if $(\mathbf{X}_{s,t})_{s \leq t}$ is the stochastic flow associated with the graphical representation of an additive particle system, then the functions $\mathbf{X}_{s,t} : \{0,1\}^\Lambda \rightarrow \{0,1\}^\Lambda$ are additive maps. We claim that $\mathbf{X}_{s,t}(x)(j) = 1$ if and only if there is an $i \in \Lambda$ with $x(i) = 1$ and it is possible to walk through the graphical representation from the space-time point (i, s) to the space time point (j, t) along a path that may use arrows, but must avoid the blocking symbols. We now make this claim more precise.


For any $i, j \in \Lambda$ and $s < u$, by definition, an *open path* from (i, s) to (j, u) is a cadlag function $\gamma : [s, u] \rightarrow \Lambda$ such that $\gamma_s = i$, $\gamma_u = j$, and

- (i) if $\gamma_{t-} \neq \gamma_t$ for some $t \in (s, u]$, then there is an arrow from (γ_{t-}, t) to (γ_t, t) ,
 - (ii) there exist no $t \in (s, u]$ such that $\gamma_{t-} = \gamma_t$ while there is a blocking symbol at (γ_t, t) .
- (6.2)

In the context of additive systems, one can check that these open paths are exactly the paths of potential influence defined in (4.20). With these definitions, we can make our earlier claim precise. We claim that:

$$\mathbf{X}_{s,t}(x)(j) = 1 \quad \text{iff} \quad \begin{array}{l} \text{there exists an } i \in \Lambda \text{ such that } x(i) = 1 \\ \text{and an open path from } (i, s) \text{ to } (j, t). \end{array} \quad (6.3)$$

To prove (6.3), it suffices to observe that if X_t denotes the right-hand side of (6.3) as a function of t , then $(X_t)_{t \geq s}$ solves the evolution equation (4.14). For example, for the graphical representation of the contact process that we earlier used as an example, the time evolution of the process $X_t := \mathbf{X}_{0,t}(X_0)$ ($t \geq 0$) might look as follows:



We now turn our attention to the backward in time process from (6.1). Let us write $(i, s) \rightsquigarrow (j, u)$ if in the graphical representation, there is an open path from (i, s) to (j, u) . We fix $u \in \mathbb{R}$ and $j \in \Lambda$, and set


$$\zeta_{u-t}^u(\{j\}) := \{i \in \Lambda : (i, u-t) \rightsquigarrow (j, u)\} \quad (t \geq 0).$$

Note that since for additive interacting particle systems, open paths and paths of potential influence are the same, this is actually the same process

as the one defined in (4.21). Then (6.3) tells us that the backward in time process from (6.1) has the following simple form:


$$\mathbf{X}_{u-t,t}[j](x) = 1_{\{x(i) = 1 \text{ for some } i \in \zeta_{u-t}^u(\{j\})\}}$$

($u \in \mathbb{R}$, $t \geq 0$, $j \in \Lambda$, $x \in \{0,1\}^\Lambda$). It is easy to see that the process $(\zeta_{u-t}^u(\{j\}))_{t \geq 0}$ is a Markov process:



Indeed, we observe that the process $(\zeta_{u-t}^u(\{j\}))_{t \geq 0}$, or rather its indicator function, evolves as a contact process started with a single infection at j . There is only one complication: contrary to our usual conventions, the process $(\zeta_{u-t}^u(\{j\}))_{t \geq 0}$ has left-continuous sample paths, but we ignore this slight complication for the moment.

In fact, there is no need to restrict ourselves to initial states with only one infected site. If we reverse the direction of all arrows and reverse the direction of time, then we can use our graphical representation to construct an additive process that runs backwards in time, as follows:



In this example, the backward in time process is also a contact process, but in general, the backward in time process can have a different dynamics from the forward in time process. Applying the principle “reverse the arrows and keep the blocking symbols” to the voter model map $\text{vot}_{i,j}$, we obtain the coalescing random walk map $\text{rw}_{j,i}$, and vice versa:



We introduce the notation

$$\mathcal{S}(\Lambda) := \{0, 1\}^\Lambda \quad \text{and} \quad \mathcal{S}_{\text{fin}}(\Lambda) := \{x \in \mathcal{S}(\Lambda) : |x| < \infty\}.$$

To describe the relation between the “forward” process $(X_t)_{t \geq 0}$ and the “backward” process $(Y_t)_{t \geq 0}$, we define a function $\psi_{\text{add}} : \mathcal{S}(\Lambda) \times \mathcal{S}(\Lambda) \rightarrow \{0, 1\}$ by

$$\psi_{\text{add}}(x, y) := 1_{\{x \wedge y \neq \underline{0}\}} \quad (x, y \in \mathcal{S}(\Lambda)), \quad (6.4)$$

where $x \wedge y := x(i) \wedge y(i)$ ($i \in \Lambda$) denotes the pointwise minimum of x and y

Lemma 6.2 (Dual additive local maps) *For each additive local map $m : \mathcal{S}(\Lambda) \rightarrow \mathcal{S}(\Lambda)$, there exists a unique “dual” map $\hat{m} : \mathcal{S}(\Lambda) \rightarrow \mathcal{S}(\Lambda)$ such that*

$$\psi_{\text{add}}(m(x), y) = \psi_{\text{add}}(x, \hat{m}(y)) \quad (x, y \in \mathcal{S}(\Lambda)). \quad (6.5)$$

This dual map \hat{m} is also additive.

One can check that \hat{m} from Lemma 6.2 is precisely the map that is constructed from m by the principle “reverse the arrows and keep the blocking symbols”. Although the proof of Lemma 6.2 is not difficult, we prefer to give the proof later in a more abstract setting.

We are now ready to formulate a pathwise duality result for general additive interacting particle systems. Let \mathcal{G} be a collection of additive local maps and let $(r_m)_{m \in \mathcal{G}}$ be nonnegative constants. We will be interested in the additive interacting particle systems with generators

$$\begin{aligned} Gf(x) &:= \sum_{m \in \mathcal{G}} r_m \{f(m(x)) - f(x)\}, \\ \hat{G}f(y) &:= \sum_{m \in \mathcal{G}} r_m \{f(\hat{m}(y)) - f(y)\}, \end{aligned} \quad (6.6)$$

where \hat{m} is the dual map associated with m in the sense of Lemma 6.2. We let ω be a Poisson point set on $\mathcal{G} \times \mathbb{R}$ with intensity measure $\rho(\{m\} \times A) := r_m \ell(A)$. Assume that the generator G satisfies the summability condition (4.15). Then by Theorem 4.19, we can use ω to define a stochastic flow $(\mathbf{X}_{s,u})_{s \leq u}$ on $\mathcal{S}(\Lambda)$. Let $\hat{\mathcal{G}} := \{\hat{m} : m \in \mathcal{G}\}$. Then setting

$$\hat{\omega} := \{(\hat{m}, -t) : (m, t) \in \omega\} \quad (6.7)$$

defines a Poisson point set $\hat{\omega}$ on $\hat{\mathcal{G}} \times \mathbb{R}$ with intensity measure $\rho(\{\hat{m}\} \times A) = r_m \ell(A)$. Assuming that also the generator \hat{G} satisfies the summability condition (4.15), we can use $\hat{\omega}$ to define a stochastic flow $(\mathbf{Y}_{s,u})_{s \leq u}$ on $\mathcal{S}(\Lambda)$. In our pictures, this corresponds to the stochastic flow we get when we reverse the direction of all arrows and the direction of time. We call $(\mathbf{Y}_{s,u})_{s \leq u}$ the *dual* stochastic flow and we call the interacting particle system $(Y_t)_{t \geq 0}$ with generator \hat{G} the *dual* interacting particle system.

It turns out that if we are only interested in constructing the dual process for initial states in $\mathcal{S}_{\text{fin}}(\Lambda)$, then in fact we do not need the summability condition (4.15) for the dual generator \hat{G} . Indeed, if we only assume the summability condition (4.15) for the forward generator G , then we have seen in Chapter 4 that the process $(\zeta_{u-t}^u(\{j\}))_{t \geq 0}$ is non-explosive, and as we have just seen this is basically the same as the dual process $(Y_t)_{t \geq 0}$ started with a single site in the state 1.

Exercise 6.3 Give an example of an additive particle system for which the forward generator G satisfies the summability condition (4.15) but the dual generator \hat{G} does not satisfy (4.15).

Before we can formulate the main result of this section, we have to take care of one more technical detail. We have already noticed that contrary to our usual conventions, the process $(\zeta_{u-t}^u(\{j\}))_{t \geq 0}$ has sample paths that are left-continuous with right limits, or *caglad*¹ for short. The choice to have cadlag sample paths in the forward time direction is, of course, mainly a matter of convention. Instead of working with cadlag functions $t \mapsto X_t$ that solve the evolution equation (4.14), we could instead have chosen to work with caglad (!) functions $[s, \infty) \ni t \mapsto X_t^- \in S^\Lambda$ that solve the equation

$$X_s^- = x \quad \text{and} \quad X_{t+}^- = \mathbf{m}_t^\omega(X_t) \quad (t \geq s), \quad (6.8)$$

where $X_{t+}^- := \lim_{u \downarrow t} X_u^-$. Then exactly the same argument as in the proof of Theorem 4.19 tells us that (6.8) has a unique solution for each $x \in S^\Lambda$ and $s \in \mathbb{R}$, which we can use to define a stochastic flow $(\mathbf{X}_{s,u}^-)_{s \leq u}$ by

$$\mathbf{X}_{s,u}^-(x) := X_u^- \quad \text{where } (X_t^-)_{t \geq s} \text{ solves (6.8).} \quad (6.9)$$

¹From the French *continue à gauche limite à droite*.

Informally, we can think of $\mathbf{X}_{s,u}^-$ (instead of $\mathbf{X}_{s,u}^+$) as the concatenation of all maps m such that $(m, t) \in \omega$ for some $t \in [s, u]$ (instead of $t \in (s, u]$). In order to have a more symmetric notation, in this chapter, we will often write $\mathbf{X}_{s,u}^+$ instead of $\mathbf{X}_{s,u}$. Note that since jumps happen only at the times of a Poisson process, we have $\mathbf{X}_{s,u}^- = \mathbf{X}_{s,u}^+$ a.s. for all deterministic $s \leq u$.

With this notation, the main result of this section reads as follows. Note that formula (6.10) below says that the map $\mathbf{Y}_{-u,-s}^-$ is dual to $\mathbf{X}_{s,u}^+$ in the sense of (6.5), and similarly $\mathbf{Y}_{-u,-s}^+$ is dual to $\mathbf{X}_{s,u}^-$.

Proposition 6.4 (Additive systems pathwise duality) *Assume that the generator G from (6.6) satisfies the summability condition (4.15). Then*

$$\psi_{\text{add}}(\mathbf{X}_{s,u}^\pm(x), y) = \psi_{\text{add}}(x, \mathbf{Y}_{-u,-s}^\mp(y)) \quad (6.10)$$

for all $s \leq u$, $x \in \mathcal{S}(\Lambda)$, and $y \in \mathcal{S}_{\text{fin}}(\Lambda)$. If the dual generator \hat{G} from (6.6) also satisfies the summability condition (4.15), then (6.10) holds for all $x, y \in \mathcal{S}(\Lambda)$.

The basic idea of the proof of Proposition 6.4 is very simple. One has $\psi_{\text{add}}(\mathbf{X}_{s,u}^+(x), y) = 1$ if and only if

$$\begin{aligned} & \exists j \in \Lambda \text{ s.t. } \mathbf{X}_{s,u}^+(x)(j) = 1 = y(j) \\ & \Leftrightarrow \exists i, j \in \Lambda \text{ s.t. } x(i) = 1, y(j) = 1, (i, s) \rightsquigarrow (j, u) \\ & \Leftrightarrow \exists i \in \Lambda \text{ s.t. } x(i) = 1 = \mathbf{Y}_{-u,-s}^-(y)(j), \end{aligned}$$

which in turn is equivalent to $\psi_{\text{add}}(x, \mathbf{Y}_{-u,-s}^-(y)) = 1$. Proposition 6.4 is an almost sure statement, which is why we talk about “pathwise” duality. As an immediate consequence, we also obtain a distributional statement.

Lemma 6.5 (Additive systems duality) *Assume that the generators G and \hat{G} in (6.6) satisfy the summability condition (4.15). Let $(X_t)_{t \geq 0}$ and $(Y_t)_{t \geq 0}$ be interacting particle systems with generators G and \hat{G} , respectively. Fix $t \geq 0$. Then*

$$\mathbb{P}[X_t \wedge Y_0 \neq \underline{0}] = \mathbb{P}[X_0 \wedge Y_t \neq \underline{0}] \quad (t \geq 0) \quad (6.11)$$

whenever X_t is independent of Y_0 and Y_t is independent of X_0 .

Proof Let $(\mathbf{X}_{s,u}^+)_{s \leq u}$ and $(\mathbf{Y}_{s,u}^-)_{s \leq u}$ be the stochastic flows constructed from the Poisson sets ω and $\hat{\omega}$. Let X_0 and Y_0 be independent of each other and of the Poisson sets ω and $\hat{\omega}$. Fix $t \geq 0$. Then setting

$$X_s := \mathbf{X}_{0,s}^+(X_0) \quad \text{and} \quad Y_s := \mathbf{Y}_{-t,s-t}^-(Y_0) \quad (s \geq 0)$$

defines interacting particle systems $(X_s)_{s \geq 0}$ and $(Y_s)_{s \geq 0}$ with generators G and \hat{G} . We claim that X_s is independent of Y_{t-s} for each $0 \leq s \leq t$. Indeed, X_s is a function of X_0 and $\{(m, u) \in \omega : 0 < u \leq s\}$, while Y_{t-s} is a function of Y_0 and $\{(m, u) \in \omega : s < u \leq t\}$. Since restrictions of Poisson sets to disjoint parts of space are independent, the claim follows. In particular, setting $s = 0$ or $s = t$, this shows that X_0 is independent of Y_t and Y_0 is independent of X_t . Now Proposition 6.4 tells us that almost surely

$$\psi_{\text{add}}(X_t, Y_0) = \psi_{\text{add}}(\mathbf{X}_{0,t}^+(X_0), Y_0) = \psi_{\text{add}}(X_0, \mathbf{Y}_{-t,0}^-(Y_0)) = \psi_{\text{add}}(X_0, Y_t).$$

Taking expectations, we see that

$$\mathbb{E}[\psi_{\text{add}}(X_t, Y_0)] = \mathbb{E}[\psi_{\text{add}}(X_0, Y_t)].$$

Recalling the definition of ψ_{add} , this gives (6.11). ■

We note that if we know the expression in (6.11) for all finite deterministic initial states $Y_0 = y$, then this determines the law of X_t uniquely. Indeed:

Lemma 6.6 (Distribution determining functions) *The functions $\{f_y : y \in \{0, 1\}^\Lambda, |y| < \infty\}$ with $f_y(x) := 1_{\{x \wedge y \neq \underline{0}\}}$ are distribution determining.*

Proof We may equivalently prove that the functions $g_y(x) := 1 - f_y(x) = 1_{\{x \wedge y = \underline{0}\}}$ are distribution determining. Since $x \wedge 1_{\{i\}}(i) = x(i)$, the class $\{g_y : |y| < \infty\}$ separates points, and since $g_y g_{y'} = g_{y \vee y'}$, this class is closed under products. The claim now follows from Lemma 4.37. ■

6.3 Cancellative systems duality

Let \oplus denote addition modulo two, i.e.,

$$0 \oplus 0 := 0, \quad 0 \oplus 1 := 1, \quad 1 \oplus 0 := 1, \quad \text{and} \quad 1 \oplus 1 := 0.$$

Let Λ and Δ be countable sets. For $x, y \in \mathcal{S}(\Lambda)$, we define $(x \oplus y) := x(i) \oplus y(i)$ ($i \in \Lambda$) in a pointwise way. By definition, a map $m : \mathcal{S}(\Lambda) \rightarrow \mathcal{S}(\Delta)$ is *cancellative* iff

- (i) $m(\underline{0}) = \underline{0}$,
- (ii) $m(x \oplus y) = m(x) \oplus m(y) \quad (x, y \in \mathcal{S}(\Lambda)).$

In the same way, we can also define cancellative maps that are defined only on $\mathcal{S}_{\text{fin}}(\Lambda)$ instead of $\mathcal{S}(\Lambda)$. An interacting particle system is called *cancellative* if its generator can be represented in cancellative local maps. Examples of cancellative maps are:

- The voter map vot_{ij} defined in (1.4).
- The *annihilating branching map* bran_{ij} defined in (6.12) below.
- The death map death_i defined in (1.7).
- The annihilating random walk map rw_{ij} defined in (1.22).
- The exclusion map excl_{ij} defined in (1.23).

Here, we define an *annihilating branching map* $\text{bran}_{ij} : \{0, 1\}^\Lambda \rightarrow \{0, 1\}^\Lambda$ as

$$\text{bran}_{ij}(x)(k) := \begin{cases} x(i) \oplus x(j) & \text{if } k = j, \\ x(k) & \text{otherwise.} \end{cases} \quad (6.12)$$

We represent additive maps by arrows and blocking symbols exactly in the same way as we did for additive maps:

- For each $i, j \in \Lambda$ with $i \neq j$ such that $m(1_{\{i\}})(j) = 1$, we draw an arrow from i to j .
- For each $i \in \Lambda$ such that $m(1_{\{i\}})(i) = 0$, we draw a blocking symbol \blacksquare at i .


The following lemma, which is similar to Lemma 6.1, shows additive local maps are fully described by their arrows and blocking symbols.

Lemma 6.7 (Cancellative local maps) *A cancellative map $m : \{0, 1\}^\Lambda \rightarrow \{0, 1\}^\Lambda$ is local if and only if its description involves only finitely many arrows and blocking symbols. Let m be a cancellative local map, let $j \in \Lambda$, and let $J \subset \Lambda$ be defined as:*


$$J := \{i \in \Lambda : i \neq j \text{ and there is an arrow from } i \text{ to } j \\ \text{or } i = j \text{ and there is no blocking symbol at } j\}.$$

Then for each $x \in \mathcal{S}(\Lambda)$, one has $m(x)(j) = 1$ if and only if $|J|$ is odd.

Every graphical representation involving arrows and blocking symbols that can be used to define an additive particle system can also be used to define a cancellative particle system. The cancellative maps mentioned above have the following representations in terms of arrows and blocking symbols:



If we interpret the graphical representation of a contact process in a cancellative way, then it becomes a graphical representation for an interacting particle system involving the annihilating branching map \mathbf{bran}_{ij} and the death map \mathbf{death}_i .



We define a *cancellative duality function* $\psi_{\text{canc}} : \mathcal{S}(\Lambda) \times \mathcal{S}_{\text{fin}}(\Lambda) \rightarrow \{0, 1\}$ by

$$\psi_{\text{canc}}(x, y) := \bigoplus_{i \in \Lambda} x(i)y(i) \quad (x \in \mathcal{S}(\Lambda), y \in \mathcal{S}_{\text{fin}}(\Lambda)). \quad (6.13)$$

Note that since $y \in \mathcal{S}_{\text{fin}}(\Lambda)$, all but finitely many of the summands are zero, so the infinite sum modulo two is well-defined. Unlike in the additive case, there is no way to make sense of $\psi_{\text{canc}}(x, y)$ for general $x, y \in \mathcal{S}(\Lambda)$.² The following lemma is similar to Lemma 6.2.

²For interacting particle systems on $\Lambda = \mathbb{Z}$, it is sometimes useful to consider the case that $\sup\{i : x(i) = 1\} < \infty$ and $\inf\{i : y(i) = 1\} > -\infty$. Clearly, $\psi_{\text{canc}}(x, y)$ is well-defined for such x, y , even though both may be infinite.

Lemma 6.8 (Dual cancellative local maps) *For each cancellative local map $m : \mathcal{S}(\Lambda) \rightarrow \mathcal{S}(\Lambda)$, there exists a unique dual map $\hat{m} : \mathcal{S}(\Lambda) \rightarrow \mathcal{S}(\Lambda)$ such that*

$$\psi_{\text{canc}}(m(x), y) = \psi_{\text{canc}}(x, \hat{m}(y)) \quad (x, y \in \mathcal{S}(\Lambda)). \quad (6.14)$$

This dual map \hat{m} is also cancellative.

One can check that \hat{m} from Lemma 6.8 is precisely the map that is constructed from m by the principle “reverse the arrows and keep the blocking symbols”. Although the proof of Lemma 6.2 is not difficult, we postpone it for the moment being. Instead, we immediately formulate a result that says that each cancellative interacting particle system has a unique dual interacting particle system.

As we did in the cancellative case, we let \mathcal{G} be a collection of cancellative local maps and let $(r_m)_{m \in \mathcal{G}}$ be nonnegative constants. We will be interested in the cancellative interacting particle systems with generators

$$\begin{aligned} Gf(x) &:= \sum_{m \in \mathcal{G}} r_m \{f(m(x)) - f(x)\}, \\ \hat{G}f(y) &:= \sum_{m \in \mathcal{G}} r_m \{f(\hat{m}(y)) - f(y)\}, \end{aligned} \quad (6.15)$$

where \hat{m} is the dual map associated with m in the sense of Lemma 6.8. We let ω be a Poisson point set on $\mathcal{G} \times \mathbb{R}$ with intensity measure $\rho(\{m\} \times A) := r_m \ell(A)$ and define $\hat{\omega}$ as in (6.7). We assume that the generator G satisfies the summability condition (4.15). We let $(\mathbf{X}_{s,u}^\pm)_{s \leq u}$ denote the cadlag and caglad stochastic flows on $\mathcal{S}(\Lambda)$ constructed from ω , and we let $(\mathbf{Y}_{s,u}^\pm)_{s \leq u}$ denote the cadlag and caglad stochastic flows on $\mathcal{S}_{\text{fin}}(\Lambda)$ constructed from $\hat{\omega}$. As in the additive case, one can check that the summability condition (4.15) for the *forward* generator G is exactly what one needs to ensure that the *dual* stochastic flow is well-defined on the space $\mathcal{S}_{\text{fin}}(\Lambda)$ of finite configurations and maps $\mathcal{S}_{\text{fin}}(\Lambda)$ into itself. We will later prove this in more detail.

The following result is similar to Proposition 6.4. Note that this time, there is no statement when the initial states x and y are both infinite, since in such a situation the duality function is not even defined.

Proposition 6.9 (Cancellative systems pathwise duality) *Assume that the generator G from (6.15) satisfies the summability condition (4.15). Then*

$$\psi_{\text{canc}}(\mathbf{X}_{s,u}^\pm(x), y) = \psi_{\text{canc}}(x, \mathbf{Y}_{-u,-s}^\mp(y)) \quad (6.16)$$

for all $s \leq u$, $x \in \mathcal{S}(\Lambda)$, and $y \in \mathcal{S}_{\text{fin}}(\Lambda)$.

The basic idea of the proof of Proposition 6.9 is similar to that of Proposition 6.4. One has $\psi_{\text{canc}}(\mathbf{X}_{s,u}^+(x), y) = 1$ if and only if

$$\begin{aligned} & |\{j \in \Lambda : \mathbf{X}_{s,u}^+(x)(j) = 1 = y(j)\}| \text{ is odd} \\ \Leftrightarrow & \text{ the number of open paths starting at a point } (i, s) \text{ such that} \\ & x(i) = 1 \text{ and ending at a point } (j,) \text{ such that } y(j) = 1 \text{ is odd} \\ \Leftrightarrow & |\{i \in \Lambda : x(i) = 1 = \mathbf{Y}_{-u,-s}^-(y)(j)\}| \text{ is odd,} \end{aligned}$$

which in turn is equivalent to $\psi_{\text{canc}}(x, \mathbf{Y}_{-u,-s}^-(y)) = 1$. Proposition 6.9 is an almost sure statement. By the same arguments used to prove Lemma 6.5, it implies the following distributional statement.

Lemma 6.10 (Cancellative systems duality) *Assume that the generators G and \hat{G} in (6.15) satisfy the summability condition (4.15). Let $(X_t)_{t \geq 0}$ and $(Y_t)_{t \geq 0}$ be interacting particle systems with generators G and \hat{G} , respectively. Fix $t \geq 0$. Then*

$$\mathbb{P}[|X_t \wedge Y_0| \text{ is odd}] = \mathbb{P}[|X_0 \wedge Y_t| \text{ is odd}] \quad (t \geq 0) \quad (6.17)$$

whenever X_t is independent of Y_0 and Y_t is independent of X_0 .

If we known the expression in (6.17) for all finite initial states $Y_0 = y$, then this determines the law of X_t uniquely. Indeed:

Lemma 6.11 (Distribution determining functions) *The set of functions $\{\psi_{\text{canc}}(\cdot, y) : y \in \mathcal{S}_{\text{fin}}(\Lambda)\}$ is distribution determining.*

Proof We may equivalently show that the functions

$$g_y(x) := 1 - 2\psi_{\text{canc}}(\cdot, y) = (-1)^{\psi_{\text{canc}}(x, y)}$$

are distribution determining. Since $\psi_{\text{canc}}(x, 1_{\{i\}}) = x(i)$, the class $\{g_y : |y| < \infty\}$ separates points, and since

$$g_y g_{y'} = g_{y \oplus y'},$$

this class is closed under products. The claim now follows from Lemma 4.37. ■

Some models that a priori do not look like cancellative systems turn out to be representable in cancellative maps. An example is the Neuhauser-Pacala model, defined by its transition rates in (1.18). We define a *rebellious map* by

$$\text{rebel}_{ijk}(x)(l) := \begin{cases} x(i) \oplus x(j) \oplus x(k) & \text{if } l = k, \\ x(l) & \text{otherwise.} \end{cases} \quad (6.18)$$

In words, this says that $x(k)$ changes its state if $x(i) \neq x(j)$.

Exercise 6.12 Show that the map rebel_{ijk} is cancellative. Show that the generator of the Neuhauser-Pacala model defined in (1.18) can be represented as

$$\begin{aligned} G_{\text{NP}}f(x) &= \frac{\alpha}{|\mathcal{N}_i|} \sum_i \sum_{j \in \mathcal{N}_i} \{f(\text{vot}_{ji}(x)) - f(x)\} \\ &= \frac{1-\alpha}{|\mathcal{N}_i|^2} \sum_i \sum_{\substack{j, k \in \mathcal{N}_i \\ j \neq k}} \{f(\text{rebel}_{kji}(x)) - f(x)\}. \end{aligned}$$

Exercise 6.13 In the threshold voter model, the site i changes its type $x(i)$ from 0 to 1 with rate one as long as at least one site in its neighborhood \mathcal{N}_i has type 1, and likewise, i flips from 1 to 0 with rate one as long as at least one site in \mathcal{N}_i has type 0. Show that the generator of the threshold voter model can be written as

$$G_{\text{thresh}}f(x) = 2^{-|\mathcal{N}_i|+1} \sum_i \sum_{\substack{\Delta \subset \mathcal{N}_i \cup \{i\} \\ |\Delta| \text{ is even}}} \{f(m_{\Delta,i}(x)) - f(x)\},$$

where $m_{\Delta,i}$ is the cancellative map defined by

$$m_{\Delta,i}(x)(k) := \begin{cases} x(i) \oplus \bigoplus_{j \in \Delta} x(j) & \text{if } k = i, \\ x(k) & \text{otherwise.} \end{cases}$$

Exercise 6.14 Show that the threshold voter model is monotone.

6.4 The backward in time process

In this and the coming sections, we will study duality of interacting particle systems in a more general setting. The aim is not only to fill in the details of some of the proofs of the results stated in the preceding sections, but also to highlight general principles that are applicable beyond the setting of additive and cancellative processes.

In the present section, we work in the general setting of Chapter 4. S is a finite set and Λ is countable. \mathcal{G} is a countable collection of local maps $m : S^\Lambda \rightarrow S^\Lambda$ and $(r_m)_{m \in \mathcal{G}}$ are nonnegative rates satisfying the summability condition (4.15). We construct a Poisson point set ω on $\mathcal{G} \times \mathbb{R}$ with intensity $r_m dt$ and let $(\mathbf{X}_{s,u}^\pm)_{s \leq u}$ be the cadlag and caglad versions of the stochastic flow constructed from ω as in (4.16) and (6.9).

Let T be a finite set and let $\mathcal{C}(S^\Lambda, T)$ denote the space of all continuous functions $f : S^\Lambda \rightarrow T$. Since by Lemma 4.13, functions in $\mathcal{C}(S^\Lambda, T)$ depend

on finitely many coordinates, it is easy to see that $\mathcal{C}(S^\Lambda, T)$ is a countable set. The following proposition generalizes the “backward in time” process first defined in (6.1) and shows that it is, indeed, a Markov process.

Proposition 6.15 (Backward in time process) *With the definitions above, assuming (4.15), setting*

$$\mathbf{F}_{s,u}^\pm(f) := f \circ \mathbf{X}_{-u,-s}^\mp \quad (s \leq u, f \in \mathcal{C}(S^\Lambda, T)). \quad (6.19)$$

defines stochastic flows $(\mathbf{F}_{s,u}^\pm)_{s \leq u}$ with independent increments, consisting of random maps $\mathbf{F}_{s,u}^\pm : \mathcal{C}(S^\Lambda, T) \rightarrow \mathcal{C}(S^\Lambda, T)$. Moreover, if F_0 is a $\mathcal{C}(S^\Lambda, T)$ -valued random variable, independent of ω , and $s \in \mathbb{R}$, then setting

$$F_t^\pm := \mathbf{F}_{s,s+t}^\pm(F_0) \quad (t \geq 0)$$

defines Markov processes $(F_t^\pm)_{t \geq 0}$ with countable state space $\mathcal{C}(S^\Lambda, T)$, where $(F_t^+)_{t \geq 0}$ has cadlag sample paths and $(F_t^-)_{t \geq 0}$ has caglad sample paths. The process $(F_t^+)_{t \geq 0}$ is the continuous-time Markov chain that evolves in such a way that

$$F_t^+ = \begin{cases} F_{t-}^+ \circ m & \text{if } (m, -t) \in \omega \text{ for some } m \in \mathcal{G}, \\ F_{t-}^+ & \text{otherwise.} \end{cases} \quad (6.20)$$

In particular, this continuous-time Markov chain is nonexplosive.

Proof By Lemma 4.26, the maps $\mathbf{X}_{-u,-s}^\mp$ are continuous, so $f \circ \mathbf{X}_{-u,-s}^\mp \in \mathcal{C}(S^\Lambda, T)$ for all $f \in \mathcal{C}(S^\Lambda, T)$. The fact that $(\mathbf{F}_{s,u}^\pm)_{s \leq u}$ are stochastic flows with independent increments now follows immediately from the analogue properties of $(\mathbf{X}_{s,u}^\pm)_{s \leq u}$. The proof that $(F_s^\pm)_{0 \leq s \leq t}$ are Markov processes follows in exactly the same way as in the proof of Proposition 2.7.

The fact that $(F_t^+)_{t \geq 0}$ evolves as in (6.20) follows immediately from its definition. We claim that for each $f \in \mathcal{C}(S^\Lambda, T)$,

$$\sum_{\substack{m \in \mathcal{G} \\ f \circ m \neq f}} r_m < \infty.$$

Indeed, $f \circ m \neq f$ implies that $\mathcal{D}(m) \cap \mathcal{R}(f) \neq \emptyset$, so our claim follows from the fact that by (4.15),

$$\sum_{\substack{m \in \mathcal{G} \\ \mathcal{D}(m) \ni R}} < \infty$$

for each finite $R \subset \Lambda$. This shows that condition (2.18) of Proposition 2.8 is satisfied, so by that proposition we know that the equation (6.20) has a

unique solution up to some possibly finite explosion time. Our construction in terms of the stochastic flow $(\mathbf{F}_{s,u}^+)_{s \leq u}$ shows that in fact, (6.20) has a solution that is defined for all time, so the continuous-time Markov chain $(F_t^+)_{t \geq 0}$ is nonexplosive. \blacksquare

Applying Proposition 6.15 to the function $f : S^\Lambda \rightarrow S$ defined as $f(x) := x(i)$, we see in particular that the backward in time process $(\mathbf{X}_{u-t,u}[i])_{t \geq 0}$ from (6.1) is a Markov chain with a countable state space. In general, this is a complicated process to study, but as might be imagined, it becomes easier in the mean-field limit. The paper [MSS20] studies the mean-field limit of $(\mathbf{X}_{u-t,u}[i])_{t \geq 0}$ and shows that it can be used to derive a stochastic representation to solutions of mean-field equations.

In the special cases of additive and cancellative interacting particle systems, we will see in a moment that the stochastic flows $(\mathbf{F}_{s,u}^\pm)_{s \leq u}$ from Proposition 6.15 map additive functions into additive functions and cancellative functions into cancellative functions, respectively, and this naturally leads to the duality functions ψ_{add} and ψ_{canc} defined in (6.4) and (6.13), respectively. In the next section, we first take a more general look at Markov process duality.

6.5 Markov process duality

Let \mathcal{X} , \mathcal{Y} , and T be sets and let $m : \mathcal{X} \rightarrow \mathcal{X}$, $\hat{m} : \mathcal{Y} \rightarrow \mathcal{Y}$, and $\psi : \mathcal{X} \times \mathcal{Y} \rightarrow T$ be functions. Then we say that \hat{m} is *dual* to m with *duality function* ψ if

$$\psi(m(x), y) = \psi(x, \hat{m}(y)) \quad (x \in \mathcal{X}, y \in \mathcal{Y}).$$

If $(\mathbf{X}_{s,t}^\pm)_{s \leq t}$ and $(\mathbf{Y}_{s,t}^\pm)_{s \leq t}$ are the cadlag and caglad versions of two stochastic flows, then we say that these stochastic flows are *dual* with *duality function* ψ if

$$\psi(\mathbf{X}_{s,t}^\pm(x), y) = \psi(x, \mathbf{Y}_{-t,-s}^\mp(y)) \quad (s \leq t, x \in \mathcal{X}, y \in \mathcal{Y}). \quad (6.21)$$

If two Markov processes $(X_t)_{t \geq 0}$ and $(Y_t)_{t \geq 0}$ can be constructed from dual stochastic flows, then we say that these Markov processes are *pathwise dual*.³

If $(X_t)_{t \geq 0}$ is pathwise dual to $(Y_t)_{t \geq 0}$ with a duality function ψ that takes values in a finite dimensional linear space \mathbb{V} , then we claim that

$$\mathbb{E}[\psi(X_t, Y_0)] = \mathbb{E}[\psi(X_0, Y_t)] \quad (t \geq 0) \quad (6.22)$$

³This terminology was first introduced in [JK14b].

whenever X_t is independent of Y_0 and Y_t is independent of X_0 . Indeed, this follows by writing

$$\mathbb{E}[\psi(X_t, Y_0)] = \mathbb{E}[\psi(\mathbf{X}_{0,t}^+(X_0), Y_0)] = \mathbb{E}[\psi(X_0, \mathbf{Y}_{-t,0}^-(Y_0))] = \mathbb{E}[\psi(X_0, Y_t)].$$

More generally, if (6.22) holds, then we say that the Markov process $(X_t)_{t \geq 0}$ is *dual* to $(Y_t)_{t \geq 0}$ with *duality function* ψ . As we will see, it can happen that (6.22) holds even though it may not be possible⁴ to construct $(X_t)_{t \geq 0}$ and $(Y_t)_{t \geq 0}$ from dual stochastic flows.

Let $(\mathbf{X}_{s \leq u}^\pm)$ be the cadlag and caglad stochastic flows constructed from the graphical representation ω of an interacting particle system $(X_t)_{t \geq 0}$ as in (4.16) and (6.9). Let T be a finite set and let $(\mathbf{F}_{s \leq u}^\pm)$, defined in (6.19), be the stochastic flows of the backward in time process $(F_t)_{t \geq 0}$ that takes values in the space $\mathcal{C}(S^\Lambda, T)$. We claim that $(X_t)_{t \geq 0}$ is pathwise dual to $(F_t)_{t \geq 0}$ with duality function $\psi : S^\Lambda \times \mathcal{C}(S^\Lambda, T) \rightarrow T$ defined as

$$\psi(x, f) := f(x) \quad (x \in S^\Lambda, f \in \mathcal{C}(S^\Lambda, T)). \quad (6.23)$$

Indeed, this follows immediately from the definition of $(\mathbf{F}_{s \leq u}^\pm)$ in (6.19), since

$$\psi(\mathbf{X}_{s,u}^\pm(x), f) = f(\mathbf{X}_{s,u}^\pm(x)) = \mathbf{F}_{-u,-s}^\mp(f)(x) = \psi(x, \mathbf{F}_{-u,-s}^\mp(f))$$

for all $x \in S^\Lambda$ and $f \in \mathcal{C}(S^\Lambda, T)$. In practice, this duality is hard to work with, since the space $\mathcal{C}(S^\Lambda, T)$ is very big. However, it sometimes happens that the stochastic flow $(\mathbf{F}_{s \leq u}^\pm)$ of the backward in time process leaves certain subspaces of $\mathcal{C}(S^\Lambda, T)$ invariant. In the next section, we will show that additive and cancellative stochastic flows leave the spaces of additive and cancellative invariant, respectively, and this naturally leads to the duality functions ψ_{add} and ψ_{canc} defined in (6.4) and (6.13).

6.6 Additive duality revisited

In this section, we return to the additive systems of Section 6.2. We fill in the details of some proofs that have only been sketched or that have been skipped altogether and at the same time reveal the general principles that underlie this duality.

As before, we let $\mathcal{C}(\mathcal{S}(\Lambda), \{0, 1\})$ denote the space of all continuous functions from $\mathcal{S}(\Lambda) = \{0, 1\}^\Lambda$ (equipped with the product topology) to $\{0, 1\}$. We let

$$\mathcal{C}_{\text{add}}(\mathcal{S}(\Lambda), \{0, 1\}) := \{m \in \mathcal{C}(\mathcal{S}(\Lambda), \{0, 1\}) : m \text{ is additive}\}$$

⁴We will not formally prove this, but it seems that the dualities covered by Theorem 6.24 can usually not be obtained in a pathwise way, except when $r \in \{1, 2\}$.

denote the subspace of additive continuous maps. Recall that if E is any topological space, then a function $f : E \rightarrow \mathbb{R}$ is called *lower semi-continuous* if one (and hence both) of the following equivalent conditions are satisfied:

- (i) $\liminf_{n \rightarrow \infty} f(x_n) \geq f(x)$ whenever $x_n \rightarrow x$,
- (ii) the level set $\{x \in E : f(x) \leq a\}$ is closed for each $a \in \mathbb{R}$.

We let $\mathcal{L}(\mathcal{S}(\Lambda), \{0, 1\})$ denote the space of all lower semi-continuous functions from $\mathcal{S}(\Lambda)$ to $\{0, 1\}$ and let

$$\mathcal{L}_{\text{add}}(\mathcal{S}(\Lambda), \{0, 1\}) := \{m \in \mathcal{L}(\mathcal{S}(\Lambda), \{0, 1\}) : m \text{ is additive}\}$$

denote the subspace of additive lower semi-continuous maps. The following lemma gives an explicit characterisation of additive lower semi-continuous maps.

Lemma 6.16 (Additive functions) *For each $y \in \mathcal{S}(\Lambda)$, let $f_y : \mathcal{S}(\Lambda) \rightarrow \{0, 1\}$ be defined as*

$$f_y(x) := \psi_{\text{add}}(x, y) = \bigvee_{i \in \Lambda} x(i)y(i) \quad (x \in \mathcal{S}(\Lambda)).$$

Then

$$\begin{aligned} \mathcal{C}_{\text{add}}(\mathcal{S}(\Lambda), \{0, 1\}) &= \{f_y : y \in \mathcal{S}_{\text{fin}}(\Lambda)\}, \\ \mathcal{L}_{\text{add}}(\mathcal{S}(\Lambda), \{0, 1\}) &= \{f_y : y \in \mathcal{S}(\Lambda)\}. \end{aligned}$$

Proof It is straightforward to check that f_y is additive for each $y \in \mathcal{S}(\Lambda)$. To see that it is moreover lower semi-continuous, assume that $x_n \rightarrow x$ pointwise. If $f_y(x) = 0$ then trivially $\liminf_{n \rightarrow \infty} f_y(x_n) \geq f_y(x)$ so without loss of generality we assume $f_y(x) = 1$. Then there exists some $i \in \Lambda$ such that $x(i) = 1 = y(i)$ and hence $x_n(i) = 1$ for all n large enough, which implies $\liminf_{n \rightarrow \infty} f_y(x_n) \geq 1$. This proves that $\{f_y : y \in \mathcal{S}(\Lambda)\} \subset \mathcal{L}_{\text{add}}(\mathcal{S}(\Lambda), \{0, 1\})$.

To prove the converse inclusion, assume that $f \in \mathcal{L}_{\text{add}}(\mathcal{S}(\Lambda), \{0, 1\})$. Define $I := \{i \in \Lambda : f(1_{\{i\}}) = 0\}$ and $L_0 := \{x : f(x) = 0\}$. Then $1_{\{i\}} \in L_0$ for all $i \in I$. Since f is additive it follows that $\bigvee_{i \in J} 1_{\{i\}} \in L_0$ for all finite $J \subset I$. Using moreover that f is lower semi-continuous and hence L_0 is closed, we see that $\bigvee_{i \in J} 1_{\{i\}} \in L_0$ for each $J \subset I$. This proves that $f(x) = 0$ for all $x \in \mathcal{S}(\Lambda)$ such that $x(j) = 0$ for all $j \in \Lambda \setminus I$. Since additive functions are monotone, we see that conversely, if $x(j) = 1$ for some $j \in \Lambda \setminus I$, then $f(x) \geq f(1_{\{j\}}) = 1$. Setting $y(i) := 0$ if $i \in I$ and $:= 1$ otherwise, it follows that $f = f_y$.

This completes the proof that $\{f_y : y \in \mathcal{S}(\Lambda)\} = \mathcal{L}_{\text{add}}(\mathcal{S}(\Lambda), \{0, 1\})$. The fact that f_y is continuous if and only if $y \in \mathcal{S}_{\text{fin}}(\Lambda)$ follows from Lemma 4.13. \blacksquare

Since the concatenation of two additive maps is again additive, for an additive interacting particle system, the stochastic flow $(\mathbf{F}_{s,u}^\pm)_{s \leq u}$ of the backward in time process, defined in (6.19), maps the space $\mathcal{C}_{\text{add}}(\mathcal{S}(\Lambda), \{0, 1\})$ into itself. By Lemma 6.16, we can identify $\mathcal{S}_{\text{fin}}(\Lambda)$ with $\mathcal{C}_{\text{add}}(\mathcal{S}(\Lambda), \{0, 1\})$ through the bijection $y \mapsto f_y$. In this identification, the abstract duality function from (6.23) takes the concrete form

$$\psi(x, f_y) = f_y(x) = \psi_{\text{add}}(x, y) \quad (x \in \mathcal{S}(\Lambda), y \in \mathcal{S}_{\text{fin}}(\Lambda)).$$

These arguments show that for each additive interacting particle system $(X_t)_{t \geq 0}$, we can find a Markov process $(Y_t)_{t \geq 0}$ that takes values in $\mathcal{S}_{\text{fin}}(\Lambda)$ so that $(X_t)_{t \geq 0}$ is pathwise dual to $(Y_t)_{t \geq 0}$ with respect to the duality function ψ_{add} . To complete our new, more precise derivation of the results of Section 6.2, we must identify this dual process more explicitly and show also an additive interacting particle system. The following lemma is very similar to the representation of additive maps in terms of arrows and blocking symbols described in Section 6.2.

Lemma 6.17 (Additive maps) *Let Λ be a countable set and let $M = (M(i, j))_{i, j \in \Lambda}$ be a matrix with values in $\{0, 1\}$. Then setting*

$$m(x)(j) := \bigvee_{i \in \Lambda} x(i) M(i, j) \quad (j \in \Lambda, x \in \mathcal{S}(\Lambda)) \quad (6.24)$$

defines an additive lower semi-continuous map $m : \mathcal{S}(\Lambda) \rightarrow \mathcal{S}(\Lambda)$. Conversely, each additive lower semi-continuous map $m : \mathcal{S}(\Lambda) \rightarrow \mathcal{S}(\Lambda)$ is of this form, and m is a local map if and only if the set

$$\{(i, j) \in \Lambda^2 : M(i, j) \neq 1(i, j)\} \quad (6.25)$$

is finite, where $1(i, j) := 1_{\{i=j\}}$ ($i, j \in \Lambda$) denotes the identity matrix.

Proof If $m : \mathcal{S}(\Lambda) \rightarrow \mathcal{S}(\Lambda)$ is additive and lower semi-continuous, then since $\mathcal{S}(\Lambda) = \{0, 1\}^\Lambda$ is equipped with the product topology, for each $j \in \Lambda$, the map $m[j] : \mathcal{S}(\Lambda) \rightarrow \{0, 1\}$ is additive and lower semi-continuous, so by Lemma 6.16 there exists an $y_j \in \mathcal{S}(\Lambda)$ such that $m[j] = f_{y_j}$. Setting $M(i, j) := y_j(i)$, this means that m is of the form (6.24). Conversely, if m is of the form (6.24), then Lemma 6.16 tells us that $m[j] : \mathcal{S}(\Lambda) \rightarrow \{0, 1\}$ is additive and lower semi-continuous for each $j \in \Lambda$, and hence m is additive and lower semi-continuous.

We recall that a map m is local if and only if it is continuous and $\mathcal{D}(m)$ is finite. By Lemma 4.13, a map of the form (6.24) is continuous if and only if $\sum_{i \in \Lambda} M(i, j) < \infty$ for all $j \in \Lambda$. Moreover, for a map of the form (6.24), we observe that $\mathcal{D}(m) = \{j \in \Lambda : M(\cdot, j) \neq 1(\cdot, j)\}$. It follows that m is local if and only if the set in (6.25) is finite. ■

An additive lower semi-continuous map $m : \mathcal{S}(\Lambda) \rightarrow \mathcal{S}(\Lambda)$ clearly determines the matrix M from (6.24) uniquely. We denote this matrix by M_m . Our next aim is to show that each additive map has a dual map, which is also additive.

Lemma 6.18 (Duals of additive maps) *Each additive lower semi-continuous map $m : \mathcal{S}(\Lambda) \rightarrow \mathcal{S}(\Lambda)$ has a unique dual map $\hat{m} : \mathcal{S}(\Lambda) \rightarrow \mathcal{S}(\Lambda)$ with respect to the duality function ψ_{add} , and the dual map \hat{m} is also an additive lower semi-continuous map. The matrix of \hat{m} is the transpose of the matrix of m , i.e., $M_{\hat{m}}(j, i) = M_m(i, j)$ ($i, j \in \Lambda$). If m is a local map, then so is \hat{m} .*

Proof Define \hat{m} by $M_{\hat{m}}(j, i) = M_m(i, j)$ ($i, j \in \Lambda$). Then

$$\psi_{\text{add}}(m(x), y) = \bigvee_{j \in \Lambda} \bigvee_{i \in \Lambda} x(i) M(i, j) y(j) = \psi_{\text{add}}(x, \hat{m}(y))$$

for all $x, y \in \mathcal{S}(\Lambda)$, which shows that m is dual to \hat{m} w.r.t. ψ . To prove that the dual of m is unique, assume that m is also dual to m' with respect to ψ_{add} . Then

$$\begin{aligned} m'(y)(i) &= \psi_{\text{add}}(1_{\{i\}}, m'(y)) = \psi_{\text{add}}(m(1_{\{i\}}), y) \\ &= \psi_{\text{add}}(1_{\{i\}}, \hat{m}(y)) = \hat{m}(y)(i) \end{aligned}$$

for all $i \in \Lambda$ and hence $m' = \hat{m}$. Now if m is a local map, then the set in (6.25) is finite for M_m and hence also for its transpose $M_{\hat{m}}$, which proves that \hat{m} is a local map. ■

We can now fill in the missing proof details from Section 6.2.

Proof of Lemmas 6.1 and 6.2 Let m be an additive map and let M_m be its matrix. Then we obtain the representation of m in terms of arrows and blocking symbols by drawing an arrow from i to j for each $i \neq j$ such that $M_m(i, j) = 1$, and drawing a blocking symbol at i for each i such that $M_m(i, i) = 0$. Then Lemma 6.17 says that each lower semi-continuous additive map can be represented in terms of arrows and blocking symbols, and such a map is local if and only if its representation contains only finitely many arrows and blocking symbols. In particular this implies Lemma 6.1. Since

taking the transpose of M_m corresponds to the recipe “reverse the arrows, keep the blocking symbols”, Lemma 6.2 follows from Lemma 6.18. ■

Proof of Proposition 6.4 Fix $y \in \mathcal{S}_{\text{fin}}(\Lambda)$, let f_y be defined as in Lemma 6.16, and let $(F_t^+)_{t \geq 0}$ be defined by

$$F_t^+ := \mathbf{F}_{0,t}^+(f_y) = f_y \circ \mathbf{X}_{-t,0}^-,$$

where $(\mathbf{F}_{s,u}^+)_{s \leq u}$ is the cadlag stochastic flow of the backward in time process, defined in (6.19). As we already remarked above Lemma 6.17, since the concatenation of two additive maps is again additive, the stochastic flow $(\mathbf{F}_{s,u}^+)_{s \leq u}$ of the backward in time process, defined in (6.19), maps the space $\mathcal{C}_{\text{add}}(\mathcal{S}(\Lambda), \{0, 1\})$ into itself. By Lemma 6.16, it follows that there exists a cadlag function $[0, \infty) \ni t \mapsto Y_t \in \mathcal{S}_{\text{fin}}(\Lambda)$ such that

$$F_t^+ = f_{Y_t} \quad (t \geq 0). \quad (6.26)$$

We have seen in Proposition 6.15 that $(F_t^+)_{t \geq 0}$ is a nonexplosive continuous-time Markov chain that evolves as in (6.20). The fact that m is dual to \hat{m} w.r.t. ψ_{add} means that

$$f_y \circ m(x) = \psi_{\text{add}}(m(x), y) = \psi_{\text{add}}(x, \hat{m}(y)) = f_{\hat{m}(y)}(x)$$

($x \in \mathcal{S}(\Lambda)$, $y \in \mathcal{S}_{\text{fin}}(\Lambda)$). It follows that $(Y_t)_{t \geq 0}$ is a nonexplosive continuous-time Markov chain that evolves as

$$Y_t = \begin{cases} \hat{m}(Y_{t-}) & \text{if } (\hat{m}, t) \in \hat{\omega} \text{ for some } m \in \mathcal{G}, \\ Y_{t-} & \text{otherwise.} \end{cases}$$

In other words, this is the process with generator \hat{G} as in (6.6). To prove (6.10), it now suffices to observe that

$$\begin{aligned} \psi_{\text{add}}(x, \mathbf{Y}_{0,t}^+(y)) &= \psi_{\text{add}}(x, Y_t) = f_{Y_t}(x) \\ &= F_t^+(x) = f_y \circ \mathbf{X}_{-t,0}^-(x) = \psi_{\text{add}}(\mathbf{X}_{-t,0}^-(x), y). \end{aligned}$$

The more general case where $\mathbf{X}_{-t,0}^-$ is replaced by $\mathbf{X}_{s,u}^\pm$ follows by the same arguments.

It remains to show that (6.10) holds for all $x, y \in \mathcal{S}(\Lambda)$ if the dual generator \hat{G} from (6.6) also satisfies the summability condition (4.15). To prove this, we observe that the duality function ψ_{add} is continuous with respect to increasing sequences, i.e., if $y_1 \leq y_2 \leq \dots$ and $y_n \rightarrow y$ pointwise, then $\psi_{\text{add}}(x, y_n) \rightarrow \psi_{\text{add}}(x, y)$. By Lemma 4.26, the random maps $\mathbf{X}_{s,u}^\pm$ and $\mathbf{Y}_{-u,-s}^\mp$ are continuous with respect to pointwise convergence. Since $\mathbf{X}_{s,u}^\pm$ and $\mathbf{Y}_{-u,-s}^\mp$ are additive, they are monotone. Using this, the final claim of Proposition 6.4 follows by approximating general $y \in \mathcal{S}(\Lambda)$ with an increasing sequence $y_n \in \mathcal{S}_{\text{fin}}(\Lambda)$. ■

6.7 Cancellative duality revisited

In this section, we fill in some of the remaining proofs for the cancellative systems of Section 6.2. The arguments are very similar to those of the previous section.

We let $\mathcal{C}_{\text{canc}}(\mathcal{S}(\Lambda), \{0, 1\})$ denote the space of cancellative continuous functions $f : \mathcal{S}(\Lambda) \rightarrow \{0, 1\}$ and we let $\mathcal{C}_{\text{canc}}(\mathcal{S}_{\text{fin}}(\Lambda), \{0, 1\})$ denote the space of all cancellative functions $f : \mathcal{S}_{\text{fin}}(\Lambda) \rightarrow \{0, 1\}$.⁵ The following lemma is similar to Lemma 6.16.

Lemma 6.19 (Cancellative functions) *One has*

$$\begin{aligned}\mathcal{C}_{\text{canc}}(\mathcal{S}(\Lambda), \{0, 1\}) &= \{\psi_{\text{canc}}(\cdot, y) : y \in \mathcal{S}_{\text{fin}}(\Lambda)\}, \\ \mathcal{C}_{\text{canc}}(\mathcal{S}_{\text{fin}}(\Lambda), \{0, 1\}) &= \{\psi_{\text{canc}}(x, \cdot) : x \in \mathcal{S}(\Lambda)\}.\end{aligned}$$

Proof It is straightforward to check that $\mathcal{S}(\Lambda) \ni x \mapsto \psi_{\text{canc}}(x, y)$ and $\mathcal{S}_{\text{fin}}(\Lambda) \ni y \mapsto \psi_{\text{canc}}(x, y)$ are cancellative. Since $\psi_{\text{canc}}(\cdot, y)$ depends on finitely many coordinates, by Lemma 4.13, it is continuous. If $f : \mathcal{S}_{\text{fin}}(\Lambda) \rightarrow \{0, 1\}$ is cancellative, then

$$f(y) = f\left(\bigoplus_{i: y(i)=1} 1_{\{i\}}\right) = \bigoplus_{i: y(i)=1} f(1_{\{i\}}) \quad (y \in \mathcal{S}_{\text{fin}}(\Lambda)),$$

so setting $x(i) := 1$ if $f(1_{\{i\}}) = 1$ and $:= 0$ otherwise, we see that $f(y) = \psi_{\text{canc}}(x, y)$ ($y \in \mathcal{S}_{\text{fin}}(\Lambda)$). If $f : \mathcal{S}(\Lambda) \rightarrow \{0, 1\}$ is cancellative and continuous, then by Lemma 4.13, there exists a finite set $\Delta \subset \Lambda$ so that f depends only on the coordinates in Δ . Each $x \in \mathcal{S}(\Lambda)$ can uniquely be written as $x = x_{\Delta} \oplus x_{\Lambda \setminus \Delta}$ where $x_{\Delta}(i) = 0$ for all $i \in \Lambda \setminus \Delta$ and $x_{\Lambda \setminus \Delta}(i) = 0$ for all $i \in \Delta$. Let $\Delta_x := \{i \in \Delta : x(i) = 1\}$. Then

$$f(x) = f\left(x_{\Lambda \setminus \Delta} \oplus \bigoplus_{i \in \Delta_x} 1_{\{i\}}\right) = f(x_{\Lambda \setminus \Delta}) \oplus \bigoplus_{i \in \Delta_x} f(1_{\{i\}}).$$

Here $f(x_{\Lambda \setminus \Delta}) = f(\underline{0}_{\Lambda \setminus \Delta}) = f(\underline{0}) = 0$ since f is cancellative and does not depend on the coordinates in $\Lambda \setminus \Delta$. Defining $y \in \mathcal{S}_{\text{fin}}(\Lambda)$ by $y := 1$ for all $i \in \Delta$ such that $f(1_{\{i\}}) = 1$ and $:= 0$ otherwise, we see that $f(x) = \psi_{\text{canc}}(x, y)$ ($x \in \mathcal{S}(\Lambda)$). \blacksquare

The following lemma replaces Lemma 6.17 in the cancellative setting.

⁵In view of our habit to equip $\mathcal{S}_{\text{fin}}(\Lambda)$ with the discrete topology, any function on $\mathcal{S}_{\text{fin}}(\Lambda)$ is automatically continuous.

Lemma 6.20 (Cancellative local maps) *Let Λ be a countable set and let $M = (M(i, j))_{i, j \in \Lambda}$ be a matrix with values in $\{0, 1\}$ such that the set*

$$\{(i, j) \in \Lambda^2 : M(i, j) \neq 1(i, j)\} \quad (6.27)$$

is finite. Then setting

$$m(x)(j) := \bigoplus_{i \in \Lambda} x(i)M(i, j) \quad (j \in \Lambda, x \in \mathcal{S}(\Lambda)) \quad (6.28)$$

defines a cancellative local map $m : \mathcal{S}(\Lambda) \rightarrow \mathcal{S}(\Lambda)$. Conversely, each cancellative local map $m : \mathcal{S}(\Lambda) \rightarrow \mathcal{S}(\Lambda)$ is of this form.

Proof This follows from the arguments used in the proof of Lemma 6.17, replacing \vee by \oplus . ■

The following lemma is similar to Lemma 6.18.

Lemma 6.21 (Duals of cancellative local maps) *Each cancellative local map $m : \mathcal{S}(\Lambda) \rightarrow \mathcal{S}(\Lambda)$ has a unique dual map $\hat{m} : \mathcal{S}(\Lambda) \rightarrow \mathcal{S}(\Lambda)$ with respect to the duality function ψ_{canc} . The dual map \hat{m} is also a cancellative local map. The matrix of \hat{m} is the transpose of the matrix of m , i.e., $M_{\hat{m}}(j, i) = M_m(i, j)$ ($i, j \in \Lambda$).*

Proof This follows from the arguments used in the proof of Lemma 6.18, replacing \vee by \oplus . ■

Proof of Lemmas 6.7 and 6.8 Just as in the additive case, the representation of a cancellative map m in terms of its matrix M_m is just a more formal way of describing the representation of a cancellative map in terms of arrows and blocking symbols. Lemmas 6.7 and 6.8 now follow immediately from Lemmas 6.20 and 6.21 exactly in the same way as we derived Lemmas 6.1 and 6.2 from Lemmas 6.17 and 6.18. ■

Proof Proposition 6.9 This is very similar to the proof of Proposition 6.4. The main idea is that the backward in time stochastic flow $(\mathbf{F}_{s,u}^\omega)_{s \leq u}$ maps the space $\mathcal{C}_{\text{canc}}(\mathcal{S}(\Lambda), \{0, 1\})$ into itself, so by Lemma 6.19, for each $y \in \mathcal{S}_{\text{fin}}$ and $s \in \mathbb{R}$, the cadlag and caglad backward in time processes started at time s in the initial state $\psi_{\text{canc}}(\cdot, y)$

$$F_t^\pm := \mathbf{F}_{s,t}^\pm(\psi_{\text{canc}}(\cdot, y)) \quad (t \geq s)$$

are of the form

$$F_t^\pm = \psi_{\text{canc}}(\cdot, Y_t^\pm) \quad (t \geq s)$$

for some $(Y_t^\pm)_{t \geq s}$ with values in \mathcal{S}_{fin} . One can then check that $\mathbf{Y}_{s,t}^\pm(y) := Y_t^\pm$ ($s \leq t$) are the cadlag and caglad stochastic flow of the continuous-time Markov chain with generator \hat{G} and that the stochastic flows $(\mathbf{X}_{s,u}^\pm)_{s \leq u}$ are dual to $(\mathbf{Y}_{s,u}^\pm)_{s \leq u}$ as in (6.16). ■

6.8 Other dualities

The additive systems duality function (6.4) and cancellative systems duality function (6.13) are not the only choices of ψ that lead to useful dualities. There are two approaches to finding useful duality functions: the pathwise approach, that aims to find dualities between stochastic flows in the sense of (6.21), and the distributional approach, that only aims to prove distributional relations of the form (6.22).

The pathwise approach always depends on finding a clever stochastic flow and then finding a suitable space of functions on S^Λ that is mapped into itself by the stochastic flow of the backward in time process. As we have seen, for additive and cancellative systems, the spaces $\mathcal{C}_{\text{add}}(\{0,1\}^\Lambda, \{0,1\})$ and $\mathcal{C}_{\text{canc}}(\{0,1\}^\Lambda, \{0,1\})$ are invariant, and this naturally leads to additive and cancellative duality. More generally, for monotone interacting particle systems, the stochastic flow maps monotone functions into monotone functions, which leads to a form of duality studied in [Gra86, SS15b]. Generalisations of additive and cancellative duality to local state spaces with three or more elements have been studied in [LS21].

To explain a bit about the distributional approach, which only aims to prove relations of the form (6.22) without proving a duality of stochastic flows, for technical simplicity, for the remainder of this section we will mostly restrict ourselves to finite state spaces. Using approximation results such as Theorem 4.41 and Corollary 4.42, many of the claims below can be extended to interacting particle systems on infinite lattices.

Lemma 6.22 (Duality of finite Markov processes) *Let $(X_t)_{t \geq 0}$ and $(Y_t)_{t \geq 0}$ be Markov processes with finite state spaces S and R , generators G and H , and Markov semigroups $(P_t)_{t \geq 0}$ and $(Q_t)_{t \geq 0}$. Then one has*

$$\mathbb{E}[\psi(X_t, Y_0)] = \mathbb{E}[\psi(X_0, Y_t)] \quad (x \in S, y \in R, t \geq 0) \quad (6.29)$$

if and only if

$$G\psi(\cdot, y)(x) = H\psi(x, \cdot)(y). \quad (6.30)$$

Proof The duality relation (6.29) says that

$$\sum_{x' \in S} P_t(x, x') \psi(x', y) = \sum_{y' \in R} \psi(x, y') Q_t(y, y') \quad (x \in S, y \in R, t \geq 0),$$

which can in matrix form be written as

$$P_t \psi = \psi Q_t^\dagger \quad (t \geq 0), \quad (6.31)$$

where $Q_t^\dagger(y', y) := Q_t(y, y')$ denotes the transpose of Q_t . Differentiating with respect to t and setting $t = 0$, it follows that

$$G\psi = \psi H^\dagger$$

which is just a more formal way of writing (6.30). Conversely, if (6.30) holds, then $G^2\psi = G\psi H^\dagger = \psi(H^\dagger)^2$ and by induction $G^n\psi = \psi(H^\dagger)^n$ for all $n \geq 0$. Using the fact that

$$P_t = \sum_{n=0}^{\infty} \frac{1}{n!} t^n G^n \quad \text{and} \quad Q_t = \sum_{n=0}^{\infty} \frac{1}{n!} t^n H^n,$$

it follows that $P_t\psi = \psi Q_t^\dagger$ ($t \geq 0$) and hence (6.29) holds. \blacksquare


Let $(X_t)_{t \geq 0}$ and $(Y_t)_{t \geq 0}$ be Markov processes with finite state spaces S and R , generators G and H , and Markov semigroups $(P_t)_{t \geq 0}$ and $(Q_t)_{t \geq 0}$. Let K be a probability kernel from S to R . A relation of the form (compare (6.31))

$$P_t K = K Q_t \quad (t \geq 0) \tag{6.32}$$

is called an *intertwining* of Markov processes. Note that (6.32) says that the following two procedures are equivalent for each S -valued random variable X_0 :

- Evolve the state X_0 for time t under the evolution of the Markov process $(X_t)_{t \geq 0}$, then map the outcome X_t into a random variable Y_t using the kernel K .
- Map X_0 into a random variable Y_0 using the kernel K , then evolve Y_0 for time t under the evolution of the Markov process $(Y_t)_{t \geq 0}$.

We can summarize the situation in the following commutative diagram:



Lemma 6.23 (Intertwining of finite Markov processes) *The intertwining relation (6.32) is equivalent to*

$$GK = KH. \tag{6.33}$$

Proof Analogue to the proof of Lemma 6.22. ■

As one might guess, there is a close relationship between duality and intertwining. If $(P_t)_{t \geq 0}$, $(Q_t)_{t \geq 0}$, and $(R_t)_{t \geq 0}$ are Markov semigroups, K is a probability kernel, and ψ a duality function such that

$$P_t K = K Q_t \quad \text{and} \quad Q_t \psi = \psi R_t^\dagger \quad (t \geq 0),$$

then trivially

$$P_t(K\psi) = K Q_t \psi = (K\psi) R_t^\dagger \quad (t \geq 0), \quad (6.34)$$

which says that the Markov processes with semigroups $(P_t)_{t \geq 0}$ and $(R_t)_{t \geq 0}$ are dual with duality function $K\psi$.

To see these general principles at work, let us look at interacting particle systems with state space of the form $\{0, 1\}^\Lambda$ where Λ is finite. For each $r > 0$, we let ψ_r denote the duality function

$$\psi_r(x, y) := \prod_{i \in \Lambda} (1 - r)^{x(i)y(i)} \quad (x, y \in \{0, 1\}^\Lambda). \quad (6.35)$$

Using the fact that $0^n = 1_{\{n=0\}}$, we observe that

$$\left. \begin{aligned} \psi_1(x, y) &= 1 - \psi_{\text{add}}(x, y) \\ \psi_2(x, y) &= (-1)^{\psi_{\text{canc}}(x, y)} \end{aligned} \right\} \quad (x, y \in \{0, 1\}^\Lambda).$$

Therefore, duality with respect to duality functions of the form (6.35) includes additive and cancellative duality as special cases.

There is a close connection between duality functions of the form (6.35) and *thinning*, as we now explain. Let $(\chi_p(i))_{i \in \Lambda}$ be i.i.d. with $\mathbb{P}[\chi_p(i) = 1] = p$ and $\mathbb{P}[\chi_p(i) = 0] = 1 - p$. Then

$$K_p(x, y) := \mathbb{P}[y(i) = \chi_p(i)x(i) \ \forall i \in \Lambda] \quad (x, y \in \{0, 1\}^\Lambda) \quad (6.36)$$

defines a *thinning kernel*. Note that if we interpret sites i with $x(i) = 1$ as being occupied by a particle, then the effect of K_p is to independently throw away some of these particles, where each particle has a probability p to remain. We claim that

$$K_p K_q = K_{pq} \quad \text{and} \quad K_p \psi_r = \psi_{pr} \quad (0 \leq p, q \leq 1, r > 0). \quad (6.37)$$

The first relation is clear from the interpretation in terms of thinning, while the second relation follows by writing

$$\begin{aligned} K_p \psi_r(x, z) &= \sum_y K_p(x, y) \prod_{i \in \Lambda} (1 - r)^{y(i)z(i)} = \mathbb{E} \left[\prod_{i \in \Lambda} (1 - r)^{\chi_p(i)x(i)z(i)} \right] \\ &= \prod_{i \in \Lambda} \mathbb{E}[(1 - r)^{\chi_p(i)x(i)z(i)}] = \prod_{i \in \Lambda} (1 - pr)^{x(i)z(i)} = \psi_{pr}(x, z). \end{aligned}$$

Let (Λ, E) be a finite graph. The paper [Sud00] considers interacting particle systems on graphs where the configuration along each edge makes the following transitions with the following rates:⁶

“annihilation”	$11 \mapsto 00$	at rate a ,
“branching”	$01 \mapsto 11$ and $10 \mapsto 11$	each at rate b ,
“coalescence”	$11 \mapsto 01$ and $11 \mapsto 10$	each at rate c ,
“death”	$01 \mapsto 00$ and $10 \mapsto 00$	each at rate d ,
“exclusion”	$01 \mapsto 10$ and $10 \mapsto 01$	each at rate e .

More formally, for each $i, j \in \Lambda$, we can define a map $m_{ij}^{01 \mapsto 11}$ on $\{0, 1\}^\Lambda$ as follows:

$$m_{ij}^{01 \mapsto 11}(x)(k) = \begin{cases} 1 & \text{if } k = i \text{ and } (x(i), x(j)) = (0, 1), \\ 1 & \text{if } k = j \text{ and } (x(i), x(j)) = (0, 1), \\ x(k) & \text{in all other cases.} \end{cases}$$

Defining $m_{ij}^{11 \mapsto 00}$ etc. in a similar way, the generator of the process we are interested in can be written as

$$\begin{aligned} Gf(x) = & \sum_{\{i,j\}} a \{f(m_{ij}^{11 \mapsto 00}(x)) - f(x)\} \\ & + \sum_{(i,j)} \left[b \{f(m_{ij}^{01 \mapsto 11}(x)) - f(x)\} + c \{f(m_{ij}^{11 \mapsto 01}(x)) - f(x)\} \right. \\ & \left. + d \{f(m_{ij}^{01 \mapsto 00}(x)) - f(x)\} + e \{f(m_{ij}^{01 \mapsto 10}(x)) - f(x)\} \right], \end{aligned} \tag{6.38}$$

where the first sum runs over all (unordered) edges $\{i, j\} \in E$ and the second sum runs over all ordered pairs (i, j) such that $\{i, j\} \in E$.

Theorem 6.24 (Lloyd-Sudbury duality) *Let G and G' be defined as in (6.38) in terms of rates a, b, c, d, e and a', b', c', d', e' , respectively, and let $r > 0$. Then one has*

$$G\psi_r = \psi_r G' \tag{6.39}$$

if and only if $a' = a + 2(1 - r)\gamma$, $b' = b + \gamma$, $c' = c - (2 - r)\gamma$, $d' = d + \gamma$, and $e' = e - \gamma$, where $\gamma := (a + c - d + (1 - r)b)/r$.

⁶The meaning of the words “annihilation”, “branching”, ... here is a bit different from the way we have used these words so far. In particular, the “death” rate d refers only to “deaths while the neighboring site is empty”, while “deaths while the neighboring site is occupied” are called “coalescence”.

Proof This follows from Lemma 6.22 by checking (6.30). The calculations are a bit tedious, so we omit them here. They can be found in [Sud00, formula (9)], which is a simplification of [SL95, formula (21)]. ■

As we have already seen, $\psi_1 = 1 - \psi_{\text{add}}$ and $\psi_2 = (-1)^{\psi_{\text{canc}}}$ correspond to additive and cancellative duality. It seems that for $r \neq 1, 2$, dualities of the form (6.39) are almost never⁷ pathwise dualities. To give an example with $r \neq 1, 2$, consider an interacting particle system on a (possibly infinite) graph (Λ, E) whose dynamics are a mixture of contact process and voter model dynamics, with generator of the form:

$$\begin{aligned} G_{\text{covov}} f(x) := & \lambda \sum_{(i,j) \in \mathcal{E}} \{f(\text{bra}_{ij}(x)) - f(x)\} \\ & + \sum_{i \in \Lambda} \{f(\text{death}_i(x)) - f(x)\} \\ & + \alpha \sum_{(i,j) \in \mathcal{E}} \{f(\text{vot}_{ij}(x)) - f(x)\} \quad (x \in \{0, 1\}^\Lambda), \end{aligned} \quad (6.40)$$

where \mathcal{E} denotes the set of oriented edges associated with E . Letting $\mathcal{N}_i := \{j \in \Lambda : \{i, j\} \in E\}$ denote the set of neighbors of i , we assume that Λ is countable and

$$\sup_{i \in \Lambda} |\mathcal{N}_i| < \infty,$$

which implies that the generator in (6.40) satisfies the summability condition (4.15) and hence, by Theorem 4.20, corresponds to a well-defined interacting particle system. Such systems are studied in [DLZ14], who are especially interested in the fast-voting limit $\alpha \rightarrow \infty$. The contact-voter model is additive (but not cancellative, because the branching map is not), and by Lemma 6.18 and Proposition 6.4 dual with respect to the duality function $\psi_1 = 1 - \psi_{\text{add}}$ to the interacting particle system with generator

$$\begin{aligned} G_{\text{covw}} f(y) := & \lambda \sum_{(i,j) \in \mathcal{E}} \{f(\text{bra}_{ij}(y)) - f(y)\} \\ & + \sum_{i \in \Lambda} \{f(\text{death}_i(y)) - f(y)\} \\ & + \alpha \sum_{(i,j) \in \mathcal{E}} \{f(\text{rw}_{ij}(y)) - f(y)\} \quad (y \in \{0, 1\}^\Lambda), \end{aligned} \quad (6.41)$$

which corresponds to a system of branching and coalescing random walks. Perhaps surprisingly, the contact-voter model is also self-dual.

⁷Except some very trivial and pathological cases.

Proposition 6.25 (Self-duality of the contact-voter model) *Assume that $\lambda > 0$. Then the contact-voter model with generator as in (6.40) is self-dual with respect to the duality function ψ_r with $r := \lambda/(\alpha + \lambda)$.*

Proof We first consider the case that the graph (Λ, E) is finite. The generator G_{covo} is a special case of the generators considered in Theorem 6.24 and corresponds to the choice of parameters

$$a = 0, \quad b = \lambda + \alpha, \quad c = 1, \quad d = 1 + \alpha, \quad e = 0.$$

We observe that setting $r := \lambda/(\alpha + \lambda)$ makes the parameter γ from Theorem 6.24 zero, which has the effect that $a' = a$, $b' = b$, $c' = c$, $d' = d$, and $e' = e$, i.e., we have found a self-duality.

To extend the result to infinite graphs, we use an approximation argument. We need to show that

$$\mathbb{E}[\psi_r(\mathbf{X}_{0,t}(x), x')] = \mathbb{E}[\psi_r(x, \mathbf{X}_{0,t}(x'))] \quad (t \geq 0, x, x' \in \{0, 1\}^\Lambda), \quad (6.42)$$

where $(\mathbf{X}_{s,u})_{s \leq u} = (\mathbf{X}_{s,u}^+)_{s \leq u}$ denotes the stochastic flow defined by the graphical representation of the contact-voter model, and

$$\psi_r(x, y) := \prod_{i \in \Lambda} (1 - r)^{x(i)y(i)} \quad (x, y \in \{0, 1\}^\Lambda). \quad (6.43)$$

Let (Λ_n, E_n) be finite subgraphs of (Λ, E) that increase to the whole graph. For each n , let $x_n(i) := x(i)$ if $i \in \Lambda_n$ and $:= 0$ otherwise, and define x'_n similarly in terms of x' . Note that if x and y are zero outside Λ , then in (6.43) it does not matter if we take the product over Λ or Λ_n . If we restrict the sums in the definition of the generator (6.40) to vertices and edges in (Λ_n, E_n) , then for the restricted process $(X_t^n)_{t \geq 0}$, the coordinates $X_t^n(i)$ with $i \in \Lambda \setminus \Lambda_n$ do not evolve (keep their initial value) while the coordinates in Λ_n form a contact-voter model on a finite graph for which Theorem 6.24 is applicable. Thus, letting $(\mathbf{X}_{s,u}^n)_{s \leq u}$ denote the stochastic flow of the restricted process, we see that for each n , we have

$$\mathbb{E}[\psi_r(\mathbf{X}_{0,t}^n(x_n), x'_n)] = \mathbb{E}[\psi_r(x_n, \mathbf{X}_{0,t}^n(x'_n))] \quad (t \geq 0, x, x' \in \{0, 1\}^\Lambda). \quad (6.44)$$

Since the contact-voter model is additive, we can express its stochastic flow in terms of open paths as in (6.3). There is a natural coupling of the graphical representations of the finite processes and the infinite process. In this coupling, $\mathbf{X}_{s,t}^n(x_n)(j) = 1$ if and only if there exists an $i \in \Lambda_n$ such that $x(i) = 1$ and an open path from (i, s) to (j, t) that never leaves the finite

set Λ_n . Using this, we see that in this coupling, $\mathbf{X}_{s,t}^n(x_n)$ a.s. increases to $\mathbf{X}_{s,t}(x)$, i.e.,

$$\mathbf{X}_{s,t}^n(x_n) \leq \mathbf{X}_{s,t}^{n+1}(x_{n+1}) \quad \forall n \quad \text{and} \quad \mathbf{X}_{s,t}^n(x_n) \xrightarrow[n \rightarrow \infty]{} \mathbf{X}_{s,t}(x) \quad \text{pointwise.} \quad (6.45)$$

Since $r \leq 1$, it is easy to check that if x_n increases to x and y_n increases to y , then $\psi_r(x_n, y_n)$ decreases to $\psi_r(x, y)$. Using this, taking the limit in (6.44), we arrive at (6.42).

Note that although the duality function ψ_r is continuous with respect to increasing sequences in the way we have just described, it is in general not true that $x_n \rightarrow x$ and $y_n \rightarrow y$ pointwise imply that $\psi_r(x_n, y_n) \rightarrow \psi_r(x, y)$. This is why we based our approximation argument on a clever monotone coupling and did not use the more general Corollary 4.42 to approximate infinite systems with finite systems. ■

We have already seen in (6.37) that there is a close connection between the duality functions ψ_r and thinning. The following proposition demonstrates this on our example of the contact-voter model.

Proposition 6.26 (Thinning of the contact-voter model) *Let $(P_t)_{t \geq 0}$ and $(Q_t)_{t \geq 0}$ denote the semigroups of the contact-voter model with generator as in (6.40) and the system of branching and coalescing random walks with generator as in (6.41), respectively. Let K_r denote the thinning kernel defined in (6.36) with $r := \lambda/(\alpha + \lambda)$. Then*

$$P_t K_r = K_r Q_t \quad (t \geq 0).$$

Proof We first prove the statement for finite graphs. Additive duality tells us that (i) $P_t \psi_1 = \psi_1 Q_t^\dagger$, and Proposition 6.25 tells us that (ii) $P_t \psi_r = \psi_r P_t^\dagger$. We want to show that this implies that (iii) $P_t K_r = K_r Q_t$. Let us first argue a bit differently and show that (i) and (iii) imply (ii). Indeed, using also (6.37), we see that (i) and (iii) imply that

$$P_t \psi_r = P_t K_r \psi_1 = K_r Q_t \psi_1 = K_r \psi_1 P_t^\dagger = \psi_r P_t^\dagger \quad (t \geq 0),$$

which is (ii). Reversing the argument, we see that (i) and (ii) imply

$$P_t K_r \psi_1 = K_r Q_t \psi_1 \quad (t \geq 0).$$

In view of this, the claim of the proposition will follow if we show that ψ_1 , as a matrix, is invertible. We can view the linear space of all functions $f : \{0, 1\}^\Lambda \rightarrow \mathbb{R}$ as the tensor product $\bigotimes_{i \in \Lambda} \mathbb{R}^{\{0, 1\}}$. In this picture, the matrix ψ_r is the tensor product over Λ of single-site matrices of the form

$$\begin{pmatrix} 1 & 1 \\ 1 & 1 - r \end{pmatrix}.$$




Figure 6.1: Graphical representation of a one-dimensional voter model and its dual system of coalescing random walks. At the final time, the points 2,3,4, and 5 have the same type, because they descend from the same ancestor.

Since these single-site matrices are invertible, so is ψ_r for all $r > 0$.

To also get the result for infinite graphs, we use approximation with finite graphs. In this case, the argument is simpler than in the proof of Proposition 6.25 since thinning is a continuous operation in the sense that if X^n are random variables with values in $\{0,1\}^\Lambda$ that converge weakly in law to X , and Y^n and Y are obtained from X^n and X by thinning with the kernel K_r , then the Y_n converge weakly in law to Y . As a result, we can use Corollary 4.42 to approximate infinite systems with finite systems and take the limit to get the result for infinite systems. ■

6.9 Invariant laws of the voter model

By Lemma 6.18 and Proposition 6.4, the voter model X is pathwise dual, with respect to the additive duality function ψ_{add} from (6.4), to a collection Y of coalescing random walks. Due to the fact that $|Y_t|$ is a nonincreasing function of t (i.e., the number of walkers can only decrease), it is much easier to work with this dual system than with the voter model itself, so duality is really the key to understanding the voter model.

Proposition 6.27 (Clustering in low dimensions) *Let X be a nearest-neighbor or range R voter model on \mathbb{Z}^d . Assume that $d = 1, 2$. Then, regardless of the initial law,*

$$\mathbb{P}[X_t(i) = X_t(j)] \xrightarrow[t \rightarrow \infty]{} 1 \quad \forall i, j \in \mathbb{Z}^d.$$

Moreover, the delta measures $\delta_{\underline{0}}$ and $\delta_{\underline{1}}$ on the constant configurations are the only extremal invariant laws.

Proof In the graphical representation of the voter model, for each $(i, t) \in \mathbb{Z}^d \times \mathbb{R}$ and $s \geq 0$, there is a unique site

$$j =: \xi_s^{(i,t)} \in \mathbb{Z}^d \text{ such that } (j, t-s) \rightsquigarrow (i, t).$$

Here $(\xi_s^{(i,t)})_{s \geq 0}$ is the path of a random walk starting at $\xi_0^{(i,t)} = i$ and “running downwards in the graphical representation”. Two such random walks started from different space-time points (i, t) and (i', t') are independent up to the first time they meet, and coalesce as soon as they meet. Moreover, if $X_t = \mathbf{X}_{0,t}(X_0)$, then, as demonstrated in Figure 6.1,

$$X_t(i) = X_{t-s}(\xi_s^{(i,t)}) \quad (0 \leq s \leq t),$$

i.e., $\xi_s^{(i,t)}$ traces back where the site i at time t got its type from.⁸

Since the difference $\xi_s^{(i,t)} - \xi_s^{(j,t)}$ of two such random walks is a random walk with absorption in the origin, and since random walk on \mathbb{Z}^d in dimensions $d = 1, 2$ is recurrent, we observe that

$$\mathbb{P}[X_t(i) = X_t(j)] \geq \mathbb{P}[\xi_t^{(i,t)} = \xi_t^{(j,t)}] = \mathbb{P}[\xi_t^{(i,0)} = \xi_t^{(j,0)}] \xrightarrow[t \rightarrow \infty]{} 1 \quad \forall i, j \in \mathbb{Z}^d.$$

This clearly implies that all invariant laws must be concentrated on constant configurations, i.e., a general invariant law is of the form $p\delta_{\underline{0}} + (1-p)\delta_{\underline{1}}$ with $p \in [0, 1]$. ■

For product initial laws we can be more precise. Although we state the following theorem for two-type processes only, it is clear from the proof that the statement generalizes basically unchanged to multitype voter models.

Theorem 6.28 (Process started in product law) *Let X be a nearest neighbor or range R voter model on \mathbb{Z}^d . Assume that the $(X_0(i))_{i \in \mathbb{Z}^d}$ are i.i.d. with intensity $\mathbb{P}[X_0(i) = 1] = p \in [0, 1]$. Then*

$$\mathbb{P}[X_t \in \cdot] \xrightarrow[t \rightarrow \infty]{} \nu_p, \quad (6.46)$$

where ν_p is an invariant law of the process. If $d = 1, 2$, then

$$\nu_p = (1-p)\delta_{\underline{0}} + p\delta_{\underline{1}}. \quad (6.47)$$

On the other hand, if $d \geq 3$ and $0 < p < 1$, then the measures ν_p are concentrated on configurations that are not constant.

⁸This construction works in fact generally for multitype voter models, where the local state space S can be any finite set, and which are in general of course not additive systems. For simplicity, we will focus on the two-type voter model here.

Proof As in the proof of Proposition 6.27, let $(\xi_s^{(i,t)})_{s \geq 0}$ be the backward random walk in the graphical representation starting at (i, t) . Define a random equivalence relation \sim on \mathbb{Z}^d by

$$i \sim j \quad \text{iff} \quad \xi_s^{(i,0)} = \xi_s^{(j,0)} \text{ for some } s \geq 0.$$

We claim that if we color the equivalence classes of \sim in an i.i.d. fashion such that each class gets the color 1 with probability p and the color 0 with probability $1 - p$, then this defines an invariant law ν_p such that (6.46) holds. Since random walk in dimensions $d = 1, 2$ is recurrent, there is a.s. only one equivalence class, and $\nu_p = (1 - p)\delta_0 + p\delta_1$. On the other hand, since random walk in dimensions $d \geq 3$ is transient, there are a.s. infinitely many⁹ equivalence classes and hence for $p \neq 0, 1$ the measure ν_p is concentrated on configurations that are not constant.

To prove (6.46), we use coupling. Let $(\chi(i))_{i \in \mathbb{Z}^d}$ be i.i.d. $\{0, 1\}$ -valued with $\mathbb{P}[\chi(i) = 1] = p$. For each $t \geq 0$, we define a random equivalence relation \sim_t on \mathbb{Z}^d by

$$i \sim_t j \quad \text{iff} \quad \xi_s^{(i,0)} = \xi_s^{(j,0)} \text{ for some } 0 \leq s \leq t.$$

We enumerate the elements of \mathbb{Z}^d in some arbitrary way and define

$$\tilde{X}_t(i) := \chi(j) \quad \text{where } j \text{ is the smallest element of } \{k \in \mathbb{Z}^d : i \sim_t k\}. \quad (6.48)$$

Then \tilde{X}_t is equally distributed with X_t and converges a.s. as $t \rightarrow \infty$ to a random variable with law ν_p . ■

Remark In dimensions $d \geq 3$, it is in fact known that the measures ν_p are extremal, and each extremal invariant law of the voter model is of this form. See [Lig85, Thm V.1.8].

6.10 Homogeneous invariant laws

In the present section, we show how the self-duality of the contact process can be used to prove that for contact processes with some sort of translation

⁹Although this is intuitively plausible, it requires a bit of work to prove this. A quick proof, that however requires a bit of ergodic theory, is as follows: since Poisson point processes are spatially ergodic, and the number N of equivalence classes is a translation-invariant random variable, this random number N must in fact be a.s. constant. Since the probability that two paths coalesce tends to zero as the distance between their starting points tends to infinity, for each finite n we can find n starting points sufficiently far from each other so that with positive probability, none of the paths started at these points coalesce. This implies that $\mathbb{P}[N \geq n] > 0$ for each finite n and hence by the fact that N is a.s. constant $\mathbb{P}[N = \infty] = 1$.

invariant structure, the upper invariant law is the limit law started from any nontrivial translation invariant initial law, and we will show that this in turn implies that the function $\theta(\lambda)$ from (5.5) is continuous everywhere, except possibly at the critical point. The methods of the present section are not restricted to additive particle systems. Applications of the technique to cancellative systems can be found in [SS08, CP14]. Applications to systems whose duals are systems of interacting diffusion processes can be found in [AS04, AS09, AS12].

We start with a simpler observation, that has been anticipated before, and which says that the functions $\theta(\lambda)$ from (1.9) and (5.5) are the same.

Lemma 6.29 (The function theta) *Let X denote the contact process with infection rate λ on a graph Λ and let $\bar{\nu}$ denote its upper invariant law. Then*

$$\int \bar{\nu}(dx) x(i) = \mathbb{P}^{1_{\{i\}}}[X_t \neq \underline{0} \ \forall t \geq 0] \quad (i \in \Lambda).$$

More generally, for any $y \in \{0, 1\}^\Lambda$ such that $|y| < \infty$,

$$\int \bar{\nu}(dx) 1_{\{x \wedge y \neq \underline{0}\}} = \mathbb{P}^y[X_t \neq \underline{0} \ \forall t \geq 0].$$

Proof By Lemma 6.18 and Proposition 6.4, the contact process X is self-dual with respect to the additive systems duality function, i.e.,

$$\mathbb{P}^x[X_t \wedge y = \underline{0}] = \mathbb{P}^y[x \wedge X_t = \underline{0}] \quad (t \geq 0).$$

In particular, setting $x = 1$, we see that

$$\begin{aligned} \int \bar{\nu}(dx) 1_{\{x \wedge y \neq \underline{0}\}} &= \lim_{t \rightarrow \infty} \mathbb{P}^1[X_t \wedge y \neq \underline{0}] \\ &= \lim_{t \rightarrow \infty} \mathbb{P}^y[1 \wedge X_t \neq \underline{0}] = \mathbb{P}^y[X_t \neq \underline{0} \ \forall t \geq 0]. \end{aligned}$$

■

In what follows, we will be interested in contact processes that have some sort of translation invariant structure. For simplicity, we will concentrate on processes on \mathbb{Z}^d with a nearest-neighbor or range R graph structure, even though the arguments can be generalized to other graphs such as infinite regular trees.

We define translation operators $T_i : \{0, 1\}^{\mathbb{Z}^d} \rightarrow \{0, 1\}^{\mathbb{Z}^d}$ by

$$T_i(x)(j) := x(j - i) \quad (i \in \mathbb{Z}^d).$$

We say that a probability law μ on $\{0, 1\}^{\mathbb{Z}^d}$ is *homogeneous* or *translation invariant* if $\mu \circ T_i^{-1} = \mu$ for all $i \in \mathbb{Z}^d$.

The main aim of the present section is to prove the following result, which is originally due to Harris [Har76]. We can think of this result as a sort of spatial analogue of the observation in Section 3.5 that for the mean-field contact process, solutions of the differential equation (3.23) started in any nonzero initial state converge to the upper fixed point. Recall from Chapter 5 that a probability law μ on $\{0, 1\}^{\mathbb{Z}^d}$ is *nontrivial* if $\mu(\{0\}) = 0$, i.e., if μ gives zero probability to the all-zero configuration.

Theorem 6.30 (Convergence to upper invariant law) *Let $(X_t)_{t \geq 0}$ be a contact process started in a homogeneous nontrivial initial law $\mathbb{P}[X_0 \in \cdot]$. Then*

$$\mathbb{P}[X_t \in \cdot] \xrightarrow[t \rightarrow \infty]{} \bar{\nu},$$

where $\bar{\nu}$ is the upper invariant law.

We start with two preparatory lemmas. We will use the graphical representation of the contact process as an additive particle system (see Section 6.2) and use the shorthand

$$X_t^x := \mathbf{X}_{0,t}(x) \quad (t \geq 0, x \in \{0, 1\}^{\mathbb{Z}^d}),$$

where $(\mathbf{X}_{s,t})_{s \leq t}$ is the stochastic flow constructed from the graphical representation as in (6.3). We continue to use the notation $|x| := \sum_i x(i)$. We say that x is *finite* if $|x| < \infty$.

Lemma 6.31 (Extinction versus unbounded growth) *For each finite $x \in \{0, 1\}^{\mathbb{Z}^d}$, one has*

$$X_t^x = \underline{0} \text{ for some } t \geq 0 \quad \text{or} \quad |X_t^x| \xrightarrow[t \rightarrow \infty]{} \infty \quad \text{a.s.} \quad (6.49)$$

Proof Define

$$\rho(x) := \mathbb{P}[X_t^x \neq \underline{0} \forall t \geq 0] \quad (x \in \{0, 1\}^{\mathbb{Z}^d}, |x| < \infty).$$

It is not hard to see that for each $N \geq 0$ there exists an $\varepsilon > 0$ such that

$$|x| \leq N \quad \text{implies} \quad \rho(x) \leq 1 - \varepsilon. \quad (6.50)$$

We first argue why it is plausible that this implies (6.49) and then give a rigorous proof. Imagine that $|X_t^x| \not\rightarrow \infty$. Then, in view of (6.50), the process infinitely often gets a chance of at least ε to die out, hence eventually it should die out.

To make this rigorous, let

$$\mathcal{A}_x := \{X_t^x \neq \underline{0} \forall t \geq 0\} \quad (x \in \{0, 1\}^{\mathbb{Z}^d}, |x| < \infty).$$

denote the event that the process $(X_t^x)_{t \geq 0}$ survives and let \mathcal{F}_t be the σ -field generated by the Poisson point processes used in our graphical representation till time t . Then

$$\rho(X_t^x) = \mathbb{P}[\mathcal{A}_x \mid \mathcal{F}_t] \xrightarrow[t \rightarrow \infty]{} 1_{\mathcal{A}_x} \quad \text{a.s.}, \quad (6.51)$$

where we have used an elementary result from probability theory that says that if \mathcal{F}_n is an increasing sequence of σ -fields and $\mathcal{F}_\infty = \sigma(\bigcup_n \mathcal{F}_n)$, then $\lim_n \mathbb{P}[\mathcal{A} \mid \mathcal{F}_n] = \mathbb{P}[\mathcal{A} \mid \mathcal{F}_\infty]$ a.s. for each measurable event \mathcal{A} . (See [Loe63, § 29, Complement 10 (b)].) In view of (6.50), formula (6.51) implies (6.49). ■

Lemma 6.32 (Nonzero intersection) *Let $(X_t)_{t \geq 0}$ be a contact process with a homogeneous nontrivial initial law $\mathbb{P}[X_0 \in \cdot]$. Then for each $s, \varepsilon > 0$ there exists an $N \geq 1$ such that for any $x \in \{0, 1\}^{\mathbb{Z}^d}$*

$$|x| \geq N \quad \text{implies} \quad \mathbb{P}[x \wedge X_s = \underline{0}] \leq \varepsilon.$$

Proof By duality,

$$\mathbb{P}[x \wedge X_s = \underline{0}] = \mathbb{P}[X_s^x \wedge X_0 = \underline{0}]$$

where X_0 is independent of the graphical representation used to define X_s^x . Set $\Lambda_M := \{-M, \dots, M\}^d$. It is not hard to see that for each $x \in \{0, 1\}^{\mathbb{Z}^d}$ with $|x| \geq N$ we can find an $x' \leq x$ with $|x'| \geq N/|\Lambda_M|$ such that the sets

$$\{i + \Lambda_M : x'(i) = 1\}$$

are disjoint, where we define $i + \Lambda_M := \{i + j : j \in \Lambda_M\}$. Write $\rightsquigarrow_{i+\Lambda_M}$ to indicate the presence of an open path that stays in $i + \Lambda_M$ and set

$$X_s^{\{i\}^{(M)}} := \{j \in \mathbb{Z}^d : (i, 0) \rightsquigarrow_{i+\Lambda_M} (j, s)\}.$$

Then, using Hölder's inequality¹⁰ in the inequality marked with an exclamation-

¹⁰Recall that Hölder's inequality says that $1/p + 1/q = 1$ implies $\|fg\|_1 \leq \|f\|_p \|g\|_q$, where $\|f\|_p := (\int |f|^p d\mu)^{1/p}$. By induction, this gives $\|\prod_{i=1}^n f_i\|_1 \leq \prod_{i=1}^n \|f_i\|_n$.

tion mark, we have

$$\begin{aligned}
\mathbb{P}[X_s^x \wedge X_0 = \underline{0}] &= \int \mathbb{P}[X_0 \in dy] \mathbb{P}[X_s^x \wedge y = \underline{0}] \\
&\leq \int \mathbb{P}[X_0 \in dy] \mathbb{P}\left[\bigvee_{i: x'(i)=1} X_s^{\{i\}(M)} \wedge y = \underline{0}\right] \\
&= \int \mathbb{P}[X_0 \in dy] \prod_{i: x'(i)=1} \mathbb{P}[X_s^{\{i\}(M)} \wedge y = \underline{0}] \\
&\stackrel{!}{\leq} \prod_{i: x'(i)=1} \left(\int \mathbb{P}[X_0 \in dy] \mathbb{P}[X_s^{\{i\}(M)} \wedge y = \underline{0}]^{|x'|} \right)^{1/|x'|} \\
&= \prod_{i: x'(i)=1} \left(\int \mathbb{P}[X_0 \in dy] \mathbb{P}[X_s^{\{0\}(M)} \wedge y = \underline{0}]^{|x'|} \right)^{1/|x'|} \\
&= \int \mathbb{P}[X_0 \in dy] \mathbb{P}[X_s^{\{0\}(M)} \wedge y = \underline{0}]^{|x'|},
\end{aligned}$$

where we have used the homogeneity of $\mathbb{P}[X_0 \in \cdot]$ in the last but one equality. Our arguments so far show that $|x| \geq N$ implies that

$$\mathbb{P}[x \wedge X_s = \underline{0}] \leq \int \mathbb{P}[X_0 \in dy] \mathbb{P}[X_s^{\{0\}(M)} \wedge y = \underline{0}]^{N/|\Lambda_M|} =: f(N, M).$$

Here, using the fact that

$$\mathbb{P}[X_s^{\{0\}(M)} \wedge y = \underline{0}] < 1 \quad \text{if } y(i) = 1 \text{ for some } i \in \Lambda_M,$$

we see that

$$\lim_{N \uparrow \infty} f(N, M) = \int \mathbb{P}[X_0 \in dy] 1_{\{y(i)=0 \ \forall i \in \Lambda_M\}} = \mathbb{P}[X_0(i) = 0 \ \forall i \in \Lambda_M].$$

Since $\mathbb{P}[X_0 \in \cdot]$ is nontrivial, we have that

$$\lim_{M \uparrow \infty} \mathbb{P}[X_0(i) = 0 \ \forall i \in \Lambda_M] = \mathbb{P}[X_0 = \underline{0}] = 0.$$

Together with our previous equation, this shows that

$$\lim_{M \rightarrow \infty} \lim_{N \rightarrow \infty} f(N, M) = 0.$$

By a diagonal argument, for each $\varepsilon > 0$ we can choose N and M_N such that $f(N, M_N) \leq \varepsilon$, proving our claim. \blacksquare

Exercise 6.33 Show by counterexample that the statement of Lemma 6.32 is false for $s = 0$.

Proof of Theorem 6.30 As in the proof of Lemma 6.31, we set

$$\rho(x) := \mathbb{P}[X_t^x \neq \underline{0} \ \forall t \geq 0] \quad (x \in \{0, 1\}^{\mathbb{Z}^d}, \ |x| < \infty).$$

By Lemmas 4.38, 6.6, and 6.29, it suffices to show that

$$\lim_{t \rightarrow \infty} \mathbb{P}[x \wedge X_t \neq \underline{0}] = \rho(x)$$

for all finite $x \in \{0, 1\}^{\mathbb{Z}^d}$. By duality, this is equivalent to showing that

$$\lim_{t \rightarrow \infty} \mathbb{P}[X_{t-s}^x \wedge X_s \neq \underline{0}] = \rho(x) \quad (x \in \{0, 1\}^{\mathbb{Z}^d}, \ |x| < \infty),$$

where $(X_t^x)_{t \geq 0}$ and $(X_t)_{t \geq 0}$ are independent and $s > 0$ is some fixed constant. For each $\varepsilon > 0$, we can choose N as in Lemma 6.32, and write

$$\begin{aligned} \mathbb{P}[X_t^x \wedge X_s \neq \underline{0}] &= \mathbb{P}[X_t^x \wedge X_s \neq \underline{0} \mid |X_t^x| = 0] \mathbb{P}[|X_t^x| = 0] \\ &\quad + \mathbb{P}[X_t^x \wedge X_s \neq \underline{0} \mid 0 < |X_t^x| < N] \mathbb{P}[0 < |X_t^x| < N] \\ &\quad + \mathbb{P}[X_t^x \wedge X_s \neq \underline{0} \mid |X_t^x| \geq N] \mathbb{P}[|X_t^x| \geq N]. \end{aligned}$$

Here, by Lemma 6.31 and our choice of N ,

- (i) $\mathbb{P}[X_t^x \wedge X_s \neq \underline{0} \mid |X_t^x| = 0] = 0,$
- (ii) $\lim_{t \rightarrow \infty} \mathbb{P}[0 < |X_t^x| < N] = 0,$
- (iii) $\liminf_{t \rightarrow \infty} \mathbb{P}[X_t^x \wedge X_s \neq \underline{0} \mid |X_t^x| \geq N] \geq 1 - \varepsilon,$
- (iv) $\lim_{t \rightarrow \infty} \mathbb{P}[|X_t^x| \geq N] = \rho(x),$

from which we conclude that

$$(1 - \varepsilon)\rho(x) \leq \liminf_{t \rightarrow \infty} \mathbb{P}[X_t^x \wedge X_s \neq \underline{0}] \leq \limsup_{t \rightarrow \infty} \mathbb{P}[X_t^x \wedge X_s \neq \underline{0}] \leq \rho(x).$$

Since $\varepsilon > 0$ is arbitrary, our proof is complete. ■

Theorem 6.30 has a simple corollary.

Corollary 6.34 (Homogeneous invariant laws) *All homogeneous invariant laws of a contact process are convex combinations of $\delta_{\underline{0}}$ and $\bar{\nu}$.*

Proof Let ν be any homogeneous invariant law. We will show that ν is a convex combination of $\delta_{\underline{0}}$ and $\bar{\nu}$. If $\nu = \delta_{\underline{0}}$ we are done. Otherwise, as in the proof of Lemma 5.10, we can write $\nu = (1 - p)\delta_{\underline{0}} + p\mu$ where $p \in (0, 1]$ and μ is a nontrivial homogeneous invariant law. But now Theorem 6.30 implies that

$$\mu = \mu P_t \xrightarrow[t \rightarrow \infty]{} \bar{\nu},$$

so we conclude that $\mu = \bar{\nu}$. ■

Recall from Exercise 5.14 that the function $\lambda \mapsto \theta(\lambda)$ from (5.5) is right-continuous everywhere. We let

$$\lambda_c := \inf\{\lambda \in \mathbb{R} : \theta(\lambda) > 0\} \quad (6.52)$$

denote the *critical point* of the contact process. As an application of Theorem 6.30, we prove the following result.

Proposition 6.35 (Continuity above the critical point) *The function $\lambda \mapsto \theta(\lambda)$ is left-continuous on (λ_c, ∞) .*

Proof Let $\bar{\nu}_\lambda$ denote the upper invariant law of the contact process with infection rate λ . Fix $\lambda > \lambda_c$ and choose $\lambda_n \uparrow \lambda$. Since the space $\mathcal{M}_1(\{0, 1\}^{\mathbb{Z}^d})$ of probability measures on $\{0, 1\}^{\mathbb{Z}^d}$, equipped with the topology of weak convergence, is compact, it suffices to show that each subsequential limit ν_* of the measures $\bar{\nu}_{\lambda_n}$ equals $\bar{\nu}_\lambda$. By Proposition 4.43, each such subsequential ν_* limit is an invariant law. It clearly is also homogeneous. Since $\lambda > \lambda_c$, by Lemma 5.10, the measures $\bar{\nu}_{\lambda_n}$ are nontrivial for n large enough, and hence, using also Proposition 5.11, the same is true for ν_* . By Corollary 6.34, we conclude that $\nu_* = \bar{\nu}$. This argument shows that the map

$$(\lambda_c, \infty) \ni \lambda \mapsto \bar{\nu}_\lambda$$

is left-continuous w.r.t. the topology of weak convergence. Since $x \mapsto x(i)$ is a continuous function and $\theta(\lambda)$ is its expectation under $\bar{\nu}_\lambda$, the claim follows. ■

6.11 Equality of critical points

The contact voter model X on \mathbb{Z}^d , that has a mixture of contact process and voter model dynamics, has been introduced in (6.40). It has two parameters: the infection rate λ and the voter rate α . We say that X *survives* if

$$\mathbb{P}^{1_{\{0\}}}[X_t \neq \underline{0} \ \forall t \geq 0] > 0.$$

For each $\alpha \geq 0$, we define critical infection rates $\lambda_c(\alpha)$ and $\lambda'_c(\alpha)$ by

$$\begin{aligned} \lambda_c(\alpha) &:= \inf \{ \lambda \in \mathbb{R} : \text{the upper invariant law is nontrivial} \}, \\ \lambda'_c(\alpha) &:= \inf \{ \lambda \in \mathbb{R} : \text{the process survives} \}. \end{aligned}$$

The paper [DLZ14] studies the asymptotics of $\lambda_c(\alpha)$ as $\alpha \rightarrow \infty$. Here, we will use duality to prove a more simple statement, namely, that $\lambda_c(\alpha) = \lambda'_c(\alpha)$ for all $\alpha \geq 0$.

For $\alpha = 0$ (i.e., the pure contact process), we already know this, as it is a direct consequence of Lemma 6.29, which follows from the self-duality of the contact process. We will use a similar argument here using Proposition 6.25, which says that the contact voter model is self-dual with respect to the duality function ψ_r from (6.35) with $r := \lambda/(\alpha + \lambda)$. Note that if $\alpha = 0$ (the pure contact process), then $r = 1$ which corresponds to additive systems duality.

Proposition 6.36 (Characterization of the upper invariant law) *Let $r := \lambda/(\alpha + \lambda)$. The upper invariant law $\bar{\nu}$ of the contact voter model satisfies*

$$\int \bar{\nu}(dx) \psi_r(x, y) = \mathbb{P}^y[X_t = \underline{0} \text{ for some } t \geq 0] \quad (6.53)$$

for all finite $y \in \{0, 1\}^{\mathbb{Z}^d}$. In particular, $\lambda_c(\alpha) = \lambda'_c(\alpha)$ for all $\alpha \geq 0$.

Proof Letting X^1 and X^y denote the processes started in $X_0^1 = 1$ and $X_0^y = y$, we observe that by Proposition 6.25,

$$\begin{aligned} \int \bar{\nu}(dx) \psi_r(x, y) &= \lim_{t \rightarrow \infty} \mathbb{E}[\psi_r(X_t^1, y)] \\ &= \lim_{t \rightarrow \infty} \mathbb{E}[\psi_r(1, X_t^y)] = \lim_{t \rightarrow \infty} \mathbb{E}[(1 - r)^{|X_t^y|}]. \end{aligned}$$

The proof of Lemma 6.31 carries over without a change to the contact voter model, so

$$X_t^y = \underline{0} \text{ for some } t \geq 0 \quad \text{or} \quad |X_t^y| \xrightarrow[t \rightarrow \infty]{} \infty \quad \text{a.s.}$$

Using this, we see that

$$\lim_{t \rightarrow \infty} \mathbb{E}[(1 - r)^{|X_t^y|}] = \mathbb{P}^y[X_t = \underline{0} \text{ for some } t \geq 0],$$

completing the proof of (6.53).

Inserting $y = 1_{\{0\}}$ into (6.53), we see that

$$\int \bar{\nu}(dx) (1 - r)^{x(0)} = \mathbb{P}^{1_{\{0\}}}[X_t = \underline{0} \text{ for some } t \geq 0],$$

or equivalently, using the fact that $1 - (1 - r)^{x(0)} = rx(0)$ with $r = \lambda/(\alpha + \lambda)$,

$$\frac{\lambda}{\alpha + \lambda} \int \bar{\nu}(dx) x(0) = \mathbb{P}^{1_{\{0\}}}[X_t \neq \underline{0} \text{ } \forall t \geq 0].$$

This shows that $\bar{\nu} = \delta_{\underline{0}}$ if and only if the process survives. ■

Chapter 7

Oriented percolation

7.1 Introduction

Although we have seen phase transitions in our simulations of interacting particle systems in Chapter 1, and we have seen how phase transitions are defined and can be calculated in the mean-field limit in Chapter 3, we have not yet proved the existence of a phase transition for any of the spatial models that we have seen so far.

In the present chapter, we fill this gap by proving that the contact process on \mathbb{Z}^d undergoes a phase transition by showing that the critical point λ_c defined in (6.52) is nontrivial in the sense that $0 < \lambda_c < \infty$. Note that by Lemma 6.29,

$$\begin{aligned}\lambda_c &= \inf\{\lambda \in \mathbb{R} : \text{the contact process survives}\} \\ &= \inf\{\lambda \in \mathbb{R} : \text{the upper invariant law is nontrivial}\}.\end{aligned}$$

In Exercise 5.12, which is based on Theorem 4.35, we have already proved for the process that

$$\frac{1}{|\mathcal{N}_0|} \leq \lambda_c,$$

where $|\mathcal{N}_0| = 2d$ or $=(2R+1)^d - 1$ is the size of the neighborhood of the origin for the nearest-neighbor process and for the range R process, respectively. In view of this, it suffices to prove that $\lambda_c < \infty$. A simple comparison argument (Exercise 5.18) shows that if the nearest-neighbor one-dimensional contact process survives for some value of λ , then the same is true for the nearest-neighbor and range R processes in dimensions $d \geq 2$. Thus, it suffices to show that $\lambda_c < \infty$ for the nearest-neighbor process in dimension one.

The method we will use is comparison with oriented percolation. This neither leads to a particularly short proof nor does it yield a very good up-


per bound on λ_c , but it has the advantage that it is a very robust method that can be applied to many other interacting particle systems. For example, in [SS08] and [SS15a], the method is applied to rebellious voter models and systems with cooperative branching and coalescing random walk dynamics, respectively. An important paper for propagating the technique was [Dur91], where this was for the first time applied to non-monotone systems and it was shown that “basically, all one needs” to prove survival is that a particle system spreads into empty areas at a positive speed.

7.2 Oriented percolation

In order to prepare for the proof that the critical infection rate of the contact process is finite, in the present section, we will study *oriented* (or *directed*) *bond percolation* on \mathbb{Z}^d . For $i, j \in \mathbb{Z}^d$, we write $i \leq j$ if $i = (i_1, \dots, i_d)$ and $j = (j_1, \dots, j_d)$ satisfy $i_k \leq j_k$ for all $k = 1, \dots, d$. Let

$$\mathcal{A} := \{(i, j) : i, j \in \mathbb{Z}^d, i \leq j, |i - j| = 1\}. \quad (7.1)$$


We view \mathbb{Z}^d as an infinite directed graph, where elements $(i, j) \in \mathcal{A}$ represent arrows (or directed *bonds*) between neighbouring sites. Note that all arrows point ‘upwards’ in the sense of the natural order on \mathbb{Z}^d .



Now fix some *percolation parameter* $p \in [0, 1]$ and let $(\omega_{(i,j)})_{(i,j) \in \mathcal{A}}$ be a collection of i.i.d. Bernoulli random variables with $\mathbb{P}[\omega_{(i,j)} = 1] = p$. We say that there is an *open path* from a site $i \in \mathbb{Z}^d$ to $j \in \mathbb{Z}^d$ if there exist $n \geq 0$ and a function $\gamma : \{0, \dots, n\} \rightarrow \mathbb{Z}^d$ such that $\gamma(0) = i$, $\gamma(n) = j$, and

$$(\gamma(k-1), \gamma(k)) \in \mathcal{A} \quad \text{and} \quad \omega_{(\gamma(k-1), \gamma(k))} = 1 \quad (k = 1, \dots, n).$$

We denote the presence of an open path by \rightsquigarrow . Note that open paths must walk upwards in the sense of the order on \mathbb{Z}^d . We write $0 \rightsquigarrow \infty$ to indicate the existence of an infinite open path starting at the origin $0 \in \mathbb{Z}^d$.



Theorem 7.2 (Critical percolation parameter) *For oriented percolation in dimensions $d \geq 2$, there exists a critical parameter $p_c = p_c(d)$ such that $\mathbb{P}[0 \rightsquigarrow \infty] = 0$ for $p < p_c$ and $\mathbb{P}[0 \rightsquigarrow \infty] > 0$ for $p > p_c$. One has*

$$\frac{1}{d} \leq p_c(d) \leq \frac{8}{9}.$$

$$p_c := \inf \{p \in [0, 1] : \mathbb{P}[0 \rightsquigarrow \infty] > 0\}.$$

To prove that $0 < p_c$, let N_n denote the number of open paths of length n starting in 0. Since there are d^n different upward paths of length n starting at the origin, and each path has probability p^n to be open, we see that

$$\mathbb{P}[N_n \neq 0] < \mathbb{E}[N_n] = d^n p^n.$$

To prove that $p_c(d) \leq 8/9$ for $d \geq 2$ it suffices to consider the case $d = 2$, for we may view \mathbb{Z}^2 as a subset of \mathbb{Z}^d ($d \geq 3$) and then, if there is an open path that stays in \mathbb{Z}^2 , then certainly there is an open path in \mathbb{Z}^d . (Note, by the way, that in $d = 1$ one has $\mathbb{P}[0 \rightsquigarrow \infty] = 0$ for all $p < 1$ hence $p_c(1) = 1$.)

We will use a Peierls argument, named after R. Peierls who used a similar argument in 1936 for the Ising model [Pei36]. In Figure 7.1, we have drawn a piece of \mathbb{Z}^2 with a random collection of open arrows. Sites $i \in \mathbb{Z}^2$ such that $0 \rightsquigarrow i$ are drawn green. These sites are called *wet*. Consider the *dual lattice*

$$\hat{\mathbb{Z}}^2 := \{(n + \tfrac{1}{2}, m + \tfrac{1}{2}) : (n, m) \in \mathbb{Z}^2\}.$$




Figure 7.1: Peierls argument for oriented percolation. The green cluster of points reachable from the origin is surrounded by a red contour. The *north* and *west* steps of this contour cannot cross open arrows.

If there are only finitely many wet sites, then the set of all non-wet sites contains one infinite connected component. (Here ‘connected’ is to be interpreted in terms of the unoriented graph \mathbb{N}^2 with nearest-neighbor edges.) Let γ be the boundary of this infinite component. Then γ is a nearest-neighbor path in $\hat{\mathbb{Z}}^2$, starting in some point $(k + \frac{1}{2}, -\frac{1}{2})$ and ending in some point $(-\frac{1}{2}, m + \frac{1}{2})$ with $k, m \geq 0$, such that all sites immediately to the left of γ are wet, and no open arrows starting at these sites cross γ . In Figure 7.1, we have indicated γ with red arrows.

From these considerations, we see that the following statement is true: one has $0 \not\rightsquigarrow \infty$ if and only if there exists a path in $\hat{\mathbb{Z}}^2$, starting in some point $(k + \frac{1}{2}, -\frac{1}{2})$ ($k \geq 0$), ending in some point $(-\frac{1}{2}, m + \frac{1}{2})$ ($m \geq 0$), and passing to the northeast of the origin, such that all arrows of γ in the north and west directions (solid red arrows in the figure) are not crossed by an open arrow. Let M_n be the number of paths of length n with these properties. Since there are $n - 1$ dual sites from where such a path of length n can start, and since in each step, there are three directions where it can go, there are less than $n3^n$ paths of length n with these properties. Since each path must make at least half of its steps in the north and west directions, the expected number of these paths satisfies

$$\mathbb{E}\left[\sum_{n=2}^{\infty} M_n\right] \leq \sum_{n=2}^{\infty} n3^n(1-p)^{n/2} < \infty \quad (p > \frac{8}{9})$$

and therefore

$$\mathbb{P}[0 \not\rightsquigarrow \infty] \leq \mathbb{P}\left[\sum_{n=2}^{\infty} M_n \geq 1\right] \leq \mathbb{E}\left[\sum_{n=2}^{\infty} M_n\right] < \infty.$$

This does not quite prove what we want yet, since we need the right-hand side of this equation to be less than one. To fix this, we use a trick. (This part of the argument comes from [Dur88].) Set $D_m := \{0, \dots, m\}^2$ and let us say that a set i is “wet” if $j \rightsquigarrow i$ for some $j \in D_m$. If $D_m \not\rightsquigarrow \infty$, then the set of wet sites must be finite, and, just as before, there must be a dual path surrounding this set of wet sites. Then, by the same arguments as before

$$\mathbb{P}[D_m \not\rightsquigarrow \infty] \leq \mathbb{P}\left[\sum_{n=2m}^{\infty} M_n \geq 1\right] \leq \mathbb{E}\left[\sum_{n=2m}^{\infty} M_n\right] \leq \sum_{n=2m}^{\infty} n3^n(1-p)^{n/2},$$

where now the sum starts at $2m$ since the dual path must surround D_m and hence have length $2m$ at least. If $p > \frac{8}{9}$, then the sum is finite so it can be made arbitrarily small by choosing m sufficiently large. It follows that $\mathbb{P}[D_m \rightsquigarrow \infty] > 0$ for some m , hence $\mathbb{P}[i \rightsquigarrow \infty] > 0$ for at least one $i \in D_m$, and therefore, by translation invariance, also $\mathbb{P}[0 \rightsquigarrow \infty] > 0$. ■

7.3 Survival

The main result of the present chapter is the following theorem, which rigorously establishes the existence of a phase transition for the contact process on \mathbb{Z}^d .

Theorem 7.3 (Nontrivial critical point) *For the nearest-neighbor or range R contact process on \mathbb{Z}^d ($d \geq 1$), the critical infection rate satisfies $0 < \lambda_c < \infty$.*

Proof As already mentioned in Section 7.1, the fact that $0 < \lambda_c$ has already been proved in Exercise 5.12. By Exercise 5.18, to prove that $\lambda_c < \infty$, it suffices to consider the one-dimensional nearest-neighbor case.

We will set up a comparison between the graphical representation of the one-dimensional nearest-neighbor contact process and oriented bond percolation on \mathbb{Z}^2 ; see Figure 7.2.

We fix $T > 0$ and define a map $\psi : \mathbb{Z}^2 \rightarrow \mathbb{Z} \times \mathbb{R}$ by

$$\psi(i) = (\kappa_i, \sigma_i) := (i_1 - i_2, T(i_1 + i_2)) \quad (i = (i_1, i_2) \in \mathbb{Z}^2).$$

The points (κ_i, σ_i) with $i \in \mathbb{N}^2$ are indicated by open circles in Figure 7.2. As before, we make \mathbb{Z}^2 into an oriented graph by defining a collection of arrows \mathcal{A} as in (7.1). We wish to define a collection $(\omega_{(i,j)})_{(i,j) \in \mathcal{A}}$ of Bernoulli random variables such that

$$\omega_{(i,j)} = 1 \quad \text{implies} \quad (\kappa_i, \sigma_i) \rightsquigarrow (\kappa_j, \sigma_j) \quad ((i,j) \in \mathcal{A}).$$




Figure 7.2: Comparison with oriented percolation. Good events in the graphical representation of the contact process (blue) correspond to open percolation arrows (black). An infinite open path along percolation arrows implies an infinite open path in the graphical representation of the contact process.

For each $i \in \mathbb{Z}^2$ we let

$$\tau_i^\pm := \inf\{t \geq \sigma_i : \text{at time } t \text{ there is an infection arrow from } \kappa_i \text{ to } \kappa_i \pm 1\}$$

denote the first time after σ_i that an arrow points out of κ_i to the left or right, respectively, and we define “good events”

$$\mathcal{G}_i^\pm := \left\{ \tau_i^\pm < \sigma_i + T \text{ and there are no blocking symbols on } \{\kappa_i\} \times (\sigma_i, \tau_i^\pm] \text{ and } \{\kappa_i \pm 1\} \times (\tau_i^\pm, \sigma_i + T] \right\}.$$

Clearly,

$$\begin{aligned} \mathcal{G}_i^- &\text{ implies } \psi(i_1, i_2) \rightsquigarrow \psi(i_1, i_2 + 1), \\ \text{and } \mathcal{G}_i^+ &\text{ implies } \psi(i_1, i_2) \rightsquigarrow \psi(i_1 + 1, i_2). \end{aligned}$$

In view of this, we set

$$\omega((i_1, i_2), (i_1, i_2 + 1)) := 1_{\mathcal{G}_i^-} \quad \text{and} \quad \omega((i_1, i_2), (i_1 + 1, i_2)) := 1_{\mathcal{G}_i^+}.$$

Then the existence of an infinite open path in the oriented percolation model defined by the $(\omega_{(i,j)})_{(i,j) \in \mathcal{A}}$ implies the existence of an infinite open path in the graphical representation of the contact process, and hence survival of the latter.




Figure 7.3: Good events use information from partially overlapping regions of space-time.

We observe that

$$p := \mathbb{P}[\omega_{(i,j)} = 1] = \mathbb{P}(\mathcal{G}_i^\pm) = (1 - e^{-\lambda T})e^{-T} \quad ((i,j) \in \mathcal{A}), \quad (7.2)$$

which tends to one as $\lambda \rightarrow \infty$ while $T \rightarrow 0$ in such a way that $\lambda T \rightarrow \infty$. It follows that for λ sufficiently large, by a suitable choice of T , we can make p as close to one as we wish. We would like to conclude from this that $\mathbb{P}[(0,0) \rightsquigarrow \infty] > 0$ for the oriented percolation defined by the $\omega_{(i,j)}$'s, and therefore also $\mathbb{P}[(0,0) \rightsquigarrow \infty] > 0$ for the contact process.

Unfortunately, life is not quite so simple, since as shown in Figure 7.3, the good events \mathcal{G}_i^\pm have been defined using information from partially overlapping space-time regions of the graphical representation of the contact process, and in view of this are not independent. They are, however, 3-dependent in the sense of Theorem 7.4 below, so by applying that result we can estimate the Bernoulli random variables $(\omega_{(i,j)})_{(i,j) \in \mathcal{A}}$ from below by i.i.d. Bernoulli random variables $(\tilde{\omega}_{(i,j)})_{(i,j) \in \mathcal{A}}$ whose success probability \tilde{p} can be made arbitrarily close to one, so we are done. ■

7.4 K-dependence

To finish the proof of Theorem 7.3 we need to provide the proof of Theorem 7.4 below, which states that k -dependent random variables with success probability p can be estimated from below by i.i.d. random variables with a success probability \tilde{p} that tends to one as $p \rightarrow 1$.

By definition, for $k \geq 0$, one says that a collection $(X_i)_{i \in \mathbb{Z}^d}$ of random variables, indexed by the integer square lattice, is k -dependent if for any

$A, B \subset \mathbb{Z}^d$ with

$$\inf\{|i - j| : i \in A, j \in B\} > k,$$

the collections of random variables $(X_i)_{i \in A}$ and $(X_j)_{j \in B}$ are independent of each other. Note that in particular, 0-dependence means independence.

It is a bit unfortunate that the traditional definition of k -dependence is strictly tied to the integer lattice \mathbb{Z}^d , while the structure of \mathbb{Z}^d has little to do with the essential idea. Therefore, in these lecture notes, we will deviate from tradition and replace(!) the definition above by the following definition.

Let Λ be countable and let $(X_i)_{i \in \Lambda}$ be a countable collection of random variables. Then we will say that the $(X_i)_{i \in \Lambda}$ are K -dependent if for each $i \in \Lambda$ there exists a $\Delta_i \subset \Lambda$ with $i \in \Delta_i$ and $|\Delta_i| \leq K$, such that

$$\chi_i \text{ is independent of } (\chi_j)_{j \in \Lambda \setminus \Delta_i}.$$

Note that according to our new definition, 1-dependence means independence. The next theorem is taken from [Lig99, Thm B26], who in turn cites [LSS97].

Theorem 7.4 (K -dependence) *Let Λ be a countable set and let $p \in (0, 1)$, $K < \infty$. Assume that $(\chi_i)_{i \in \Lambda}$ are K -dependent Bernoulli random variables with $P[\chi_i = 1] \geq p$ ($i \in \Lambda$), and that*

$$\tilde{p} := (1 - (1 - p)^{1/K})^2 \geq \frac{1}{4}.$$

Then it is possible to couple $(\chi_i)_{i \in \Lambda}$ to a collection of independent Bernoulli random variables $(\tilde{\chi}_i)_{i \in \Lambda}$ with

$$P[\tilde{\chi}_i = 1] = \tilde{p} \quad (i \in \Lambda), \tag{7.3}$$

in such a way that $\tilde{\chi}_i \leq \chi_i$ for all $i \in \Lambda$.

Proof Since we can always choose some arbitrary denumeration of Λ , we may assume that $\Lambda = \mathbb{N}$. Our strategy will be as follows. We will choose $\{0, 1\}$ -valued random variables $(\psi_i)_{i \in \mathbb{N}}$ with $P[\psi_i = 1] = r$, independent of each other and of the $(\chi_i)_{i \in \mathbb{N}}$, and put

$$\chi'_i := \psi_i \chi_i \quad (i \in \mathbb{N}).$$

Note that the $(\chi'_i)_{i \in \mathbb{N}}$ are a ‘thinned out’ version of the $(\chi_i)_{i \in \mathbb{N}}$. In particular, $\chi'_i \leq \chi_i$ ($i \in \mathbb{N}$). We will show that for an appropriate choice of r ,

$$P[\chi'_n = 1 \mid \chi'_0, \dots, \chi'_{n-1}] \geq \tilde{p} \tag{7.4}$$

for all $n \geq 0$, and we will show that this implies that the $(\chi'_i)_{i \in \mathbb{N}}$ can be coupled to independent $(\tilde{\chi}_i)_{i \in \Lambda}$ as in (7.3) in such a way that $\tilde{\chi}_i \leq \chi'_i \leq \chi_i$ ($i \in \mathbb{N}$).

We start with the latter claim. Imagine that (7.4) holds. Set $p'_0 := P[\chi'_0 = 1]$ and

$$p'_n(\varepsilon_0, \dots, \varepsilon_{n-1}) := P[\chi'_n = 1 \mid \chi'_0 = \varepsilon_0, \dots, \chi'_{n-1} = \varepsilon_{n-1}]$$

whenever $P[\chi'_0 = \varepsilon_0, \dots, \chi'_{n-1} = \varepsilon_{n-1}] > 0$. Let $(U_n)_{n \in \mathbb{N}}$ be independent, uniformly distributed $[0, 1]$ -valued random variables. Set

$$\tilde{\chi}_n := 1_{\{U_n < \tilde{p}\}} \quad (n \in \mathbb{N})$$

and define inductively

$$\chi'_n := 1_{\{U_n < p'_n(\chi'_0, \dots, \chi'_{n-1})\}} \quad (n \in \mathbb{N}).$$

Then

$$P[\chi'_n = \varepsilon_n, \dots, \chi'_0 = \varepsilon_0] = p'_n(\varepsilon_0, \dots, \varepsilon_{n-1}) \cdots p'_1(\varepsilon_0) \cdot p'_0.$$

This shows that these new χ'_n 's have the same distribution as the old ones, and they are coupled to $\tilde{\chi}_i$'s as in (7.3) in such a way that $\tilde{\chi}_i \leq \chi'_i$.

What makes life complicated is that (7.4) does not always hold for the original $(\chi_i)_{i \in \mathbb{N}}$, which is why we have to work with the thinned variables $(\chi'_i)_{i \in \mathbb{N}}$.¹ We observe that

$$\begin{aligned} P[\chi'_n = 1 \mid \chi'_0 = \varepsilon_0, \dots, \chi'_{n-1} = \varepsilon_{n-1}] \\ = rP[\chi_n = 1 \mid \chi_0 = \varepsilon_0, \dots, \chi_{n-1} = \varepsilon_{n-1}]. \end{aligned} \quad (7.5)$$

We will prove by induction that for an appropriate choice of r ,

$$P[\chi_n = 0 \mid \chi'_0 = \varepsilon_0, \dots, \chi'_{n-1} = \varepsilon_{n-1}] \leq 1 - r. \quad (7.6)$$

Note that this is true for $n = 0$ provided that $r \leq p$. Let us put

$$\begin{aligned} E_0 &:= \{i \in \Delta_n : 0 \leq i \leq n-1, \varepsilon_i = 0\}, \\ E_1 &:= \{i \in \Delta_n : 0 \leq i \leq n-1, \varepsilon_i = 1\}, \\ F &:= \{i \notin \Delta_n : 0 \leq i \leq n-1\}. \end{aligned}$$

¹Indeed, let $(\phi_n)_{n \geq 0}$ be independent $\{0, 1\}$ -valued random variables with $P[\phi_n = 1] = \sqrt{p}$ for some $p < 1$, and put $\chi_n := \phi_n \phi_{n+1}$. Then the $(\chi_n)_{n \geq 0}$ are 3-dependent with $P[\chi_n = 1] = p$, but $P[\chi_n = 1 \mid \chi_{n-1} = 0, \chi_{n-2} = 1] = 0$.

Then

$$\begin{aligned}
& P[\chi_n = 0 \mid \chi'_0 = \varepsilon_0, \dots, \chi'_{n-1} = \varepsilon_{n-1}] \\
&= P[\chi_n = 0 \mid \chi'_i = 0 \ \forall i \in E_0, \chi_i = 1 = \psi_i \ \forall i \in E_1, \chi'_i = \varepsilon_i \ \forall i \in F] \\
&= P[\chi_n = 0 \mid \chi'_i = 0 \ \forall i \in E_0, \chi_i = 1 \ \forall i \in E_1, \chi'_i = \varepsilon_i \ \forall i \in F] \\
&= \frac{P[\chi_n = 0, \chi'_i = 0 \ \forall i \in E_0, \chi_i = 1 \ \forall i \in E_1, \chi'_i = \varepsilon_i \ \forall i \in F]}{P[\chi'_i = 0 \ \forall i \in E_0, \chi_i = 1 \ \forall i \in E_1, \chi'_i = \varepsilon_i \ \forall i \in F]} \\
&\leq \frac{P[\chi_n = 0, \chi'_i = \varepsilon_i \ \forall i \in F]}{P[\psi_i = 0 \ \forall i \in E_0, \chi_i = 1 \ \forall i \in E_1, \chi'_i = \varepsilon_i \ \forall i \in F]} \\
&= \frac{P[\chi_n = 0 \mid \chi'_i = \varepsilon_i \ \forall i \in F]}{P[\psi_i = 0 \ \forall i \in E_0, \chi_i = 1 \ \forall i \in E_1 \mid \chi'_i = \varepsilon_i \ \forall i \in F]} \\
&\leq \frac{1-p}{(1-r)^{|E_0|} P[\chi_i = 1 \ \forall i \in E_1 \mid \chi'_i = \varepsilon_i \ \forall i \in F]} \leq \frac{1-p}{(1-r)^{|E_0|} r^{|E_1|}}, \tag{7.7}
\end{aligned}$$

where in the last step we have used K -dependence and the (nontrivial) fact that

$$P[\chi_i = 1 \ \forall i \in E_1 \mid \chi'_i = \varepsilon_i \ \forall i \in F] \geq r^{|E_1|}. \tag{7.8}$$

We claim that (7.8) is a consequence of the induction hypothesis (7.6). Indeed, we may assume that the induction hypothesis (7.6) holds regardless of the ordering of the first n elements, so without loss of generality we may assume that $E_1 = \{n-1, \dots, m\}$ and $F = \{m-1, \dots, 0\}$, for some m . Then the left-hand side of (7.8) may be written as

$$\begin{aligned}
& \prod_{k=m}^{n-1} P[\chi_k = 1 \mid \chi_i = 1 \ \forall m \leq i < k, \chi'_i = \varepsilon_i \ \forall 0 \leq i < m] \\
&= \prod_{k=m}^{n-1} P[\chi_k = 1 \mid \chi'_i = 1 \ \forall m \leq i < k, \chi'_i = \varepsilon_i \ \forall 0 \leq i < m] \geq r^{n-m}.
\end{aligned}$$

If we assume moreover that $r \geq \frac{1}{2}$, then $r^{|E_1|} \geq (1-r)^{|E_1|}$ and therefore the right-hand side of (7.7) can be further estimated as

$$\frac{1-p}{(1-r)^{|E_0|} r^{|E_1|}} \leq \frac{1-p}{(1-r)^{|\Delta_n \cap \{0, \dots, n-1\}|}} \leq \frac{1-p}{(1-r)^{K-1}}.$$

We see that in order for our proof to work, we need $\frac{1}{2} \leq r \leq p$ and

$$\frac{1-p}{(1-r)^{K-1}} \leq 1-r. \tag{7.9}$$

In particular, choosing $r = 1 - (1 - p)^{1/K}$ yields equality in (7.9). Having proved (7.6), we see by (7.5) that (7.4) holds provided that we put $\tilde{p} := r^2$. ■

Exercise 7.5 *Combine Theorem 7.2 and formulas (7.2) and (7.3) to derive an explicit upper bound on the critical infection rate λ_c of the one-dimensional contact process.*

Exercise 7.6 *The one-dimensional contact process with double deaths has been introduced just before Exercise 5.15. Use comparison with oriented percolation to prove that the one-dimensional contact process with double deaths survives with positive probability if its branching rate λ is large enough. When you apply Theorem 7.4, what value of K do you (at least) need to use?*

Exercise 7.7 *Use the previous exercise and Exercise 5.15 to conclude that for the cooperative branching process considered there, if λ is large enough, then: 1° If the process is started with at least two particles, then there is a positive probability that the number of particles will always be at least two. 2° The upper invariant law is nontrivial.*

Exercise 7.8 *Assume that there exists some $t > 0$ such that the contact process satisfies*

$$r := \mathbb{E}^{1_{\{0\}}} [|X_t|] < 1.$$

Show that this then implies that

$$\mathbb{E}^{1_{\{0\}}} [|X_{nt}|] \leq r^n \quad (n \geq 0)$$

and the process started in any finite initial state dies out a.s. Can you use this to improve the lower bound $1/|\mathcal{N}_i| \leq \lambda_c$ from Exercise 5.12, e.g., for the one-dimensional nearest-neighbor process?

Bibliography

- [ADS15] M. Aizenman, H. Duminil-Copin, and V. Sidoravicius. Random currents and continuity of Ising model's spontaneous magnetization. *Commun. Math. Phys.* 334 (2015), 719–742.
- [AS04] S.R. Athreya and J.M. Swart. Branching-coalescing particle systems. *Prob. Theory Relat. Fields* 131(3) (2005), 376–414.
- [AS09] S.R. Athreya and J.M. Swart. Branching-coalescing particle systems. Erratum. *Prob. Theory Relat. Fields.* 145(3-4) (2009), 639–640.
- [AS12] S.R. Athreya and J.M. Swart. Systems of branching, annihilating, and coalescing particles. *Electron. J. Probab.* 17 (2012), paper No. 80, 1–32.
- [AU29] P. Alexandroff and P. Urysohn. Mémoire sur les espaces topologiques compacts, dédié à Monsieur D. Egoroff. *Verhandelingen Amsterdam* 14(1) (1929) 1–93.
- [BD01] W.A. Brock and S.N. Durlauf. Discrete choice with social interactions. *The Review of Economic Studies* 68(2) (2001), 235–260.
- [BDT19] O. Blondel, A. Deshayes, and C. Toninelli. Front evolution of the Fredrickson-Andersen one spin facilitated model. *Electron. J. Probab.* 24 (2019), paper No. 1, 1–32.
- [BG90] C. Bezuidenhout and G. Grimmett. The critical contact process dies out. *Ann. Probab.* 18(4) (1990), 1462–1482.
- [CDFT20] R. Cerf, P. Dai Pra, M. Formentin, and D. Tovazzi. Rhythmic behavior of an Ising Model with dissipation at low temperature. Preprint (2020), arXiv:2002.08244.

- [CFT16] F. Collet, M. Formentin, and D. Tovazzi. Rhythmic behavior in a two-population mean-field Ising model. *Phys. Rev. E* 94 (2016), 042139.
- [CP14] J.T. Cox and E.A. Perkins. A complete convergence theorem for voter model perturbations. *Ann. Appl. Probab.* 24(1) (2014), 150–197.
- [CS73] P. Clifford and A. Sudbury. A model for spatial conflict. *Biometrika* 60(3) (1973), 581–588.
- [DFR13] P. Dai Pra, M. Fischer, and D. Regoli. A Curie-Weiss model with dissipation. *J. Stat. Phys.* 152(1) (2013), 37–53.
- [DLZ14] R. Durrett, T. Liggett, and Y. Zhang. The contact process with fast voting. *Electron. J. Probab.* 19 (2014), Paper No. 28, 1–19.
- [DN08] R.W.R. Darling and J.R. Norris. Differential equation approximations for Markov chains. *Probab. Surveys* 5 (2008), 37–79.
- [DST17] H. Duminil-Copin, V. Sidoravicius, and V. Tassion. Continuity of the phase transition for planar random-cluster and Potts models with $1 \leq q \leq 4$. *Commun. Math. Phys.* 349(1) (2017), 47–107.
- [Dur88] R. Durrett. *Lecture notes on particle systems and percolation*. Wadsworth & Brooks/Cole, Pacific Grove, 1988.
- [Dur91] R. Durrett. A new method for proving the existence of phase transitions. *Spatial Stochastic Processes, Festschr. in Honor of Ted Harris 70th Birthday* (K.S. Alexander and J.C. Watkins eds.) 141–169. *Prog. Probab.* 19, Birkhäuser, Boston, 1991.
- [Dur92] R. Durrett. Stochastic growth models: Bounds on critical values. *J. Appl. Probab.* 29(1) (1992), 11–20.
- [EK86] S.N. Ethier and T.G. Kurtz. *Markov Processes: Characterization and Convergence*. John Wiley & Sons, New York, 1986.
- [FM01] J.A. Fill and M. Machida. Stochastic monotonicity and realizable monotonicity. *Ann. Probab.* 29(2), (2001), 938–978.
- [GB07] T. Gobron and I. Merola. First-order phase transition in Potts models with finite-range interactions. *J. Stat. Phys.* 126(3) (2007), 507–583.

- [GPA01] M. Ghaemi, G.A. Parsafar, and M. Ashrafizaadeh. Calculation of the critical temperature for 2- and 3-dimensional Ising models and for 2-dimensional Potts models using the transfer matrix method. *J. Phys. Chem. B* 105(42) (2001), 10355–10359.
- [Gra86] L. Gray Duality for general attractive spin systems with applications in one dimension. *Ann. Probab.* 14(2) (1986), 371–396.
- [Har74] T.E. Harris. Contact interactions on a lattice. *Ann. Probab.* 2 (1974), 969–988.
- [Har76] T.E. Harris. On a class of set-valued Markov processes. *Ann. Probab.* 4 (1976), 175–194.
- [Isi25] E. Ising. Beitrag zur Theorie des Ferromagnetismus. *Zeitschrift für Physik* 31 (1925), 253–258.
- [JK14a] B. Jahnke and C. Külske. A class of nonergodic interacting particle systems with unique invariant measure. *Ann. Appl. Probab.* 24(6) (2014), 2595–2643.
- [JK14b] S. Jansen and N. Kurt. On the notion(s) of duality for Markov processes. *Prob. Surveys* 11 (2014), 59–120.
- [Kal97] O. Kallenberg. *Foundations of modern probability*. Springer, New York, 1997.
- [Kol56] A.N. Kolmogorov. On Skorohod convergence. *Theor. Probability Appl.* 1 (1956), 213–222.
- [KKO77] T. Kamae, U. Krengel, and G.L. O’Brien. Stochastic inequalities on partially ordered spaces. *Ann. Probab.* 5 (1977), 899–912.
- [Kro99] S. Krone. The two-stage contact process. *Ann. Appl. Probab.* 9(2) (1999), 331–351.
- [Law05] G.F. Lawler. *Conformally Invariant Processes in the Plane*. Mathematical Surveys and Monographs 114. American Mathematical Society, Providence, 2005.
- [Lig85] T.M. Liggett. *Interacting Particle Systems*. Springer-Verlag, New York, 1985.
- [Lig95] T.M. Liggett. Improved upper bounds for the contact process critical value. *Ann. Probab.* 23 (1995), 697–723.

- [Lig99] T.M. Liggett. *Stochastic interacting systems: contact, voter and exclusion processes*. Springer-Verlag, Berlin, 1999.
- [Lig10] T.M. Liggett. *Continuous time Markov processes. An introduction*. AMS, Providence, 2010.
- [LS21] J.N. Latz and J.M. Swart. Commutative monoid duality. Preprint (2021), arXiv:2108.01492.
- [LSS97] T.M. Liggett, R.H. Schonmann and A.M. Stacey. Domination by product measures. *Ann. Probab.* 25(1) (1997), 71–95.
- [Loe63] M. Loève. *Probability Theory, 3rd edition*. Van Nostrand, Princeton, 1963.
- [Mou95] T.S. Mountford. A coupling of infinite particle systems. *J. Math. Kyoto Univ.* 35(1) (1995), 43–52.
- [MSS20] T. Mach, A. Sturm, and J.M. Swart. Recursive tree processes and the mean-field limit of stochastic flows. *Electron. J. Probab.* 25 (2020) paper No. 61, 1–63.
- [Neu94] C. Neuhauser. A long range sexual reproduction process. *Stochastic Processes Appl.* 53(2) (1994), 193–220.
- [Nob92] C. Noble. Equilibrium behavior of the sexual reproduction process with rapid diffusion. *Ann. Probab.* 20(2) (1992), 724–745.
- [NP99] C. Neuhauser and S.W. Pacala. An explicitly spatial version of the Lotka-Volterra model with interspecific competition. *Ann. Appl. Probab.* 9(4) (1999), 1226–1259.
- [Ons44] L. Onsager. Crystal statistics, I. A two-dimensional model with an order-disorder transition. *Phys. Rev.* 65 (1944), 117–149.
- [Pei36] R. Peierls. On Ising’s model of ferromagnetism. *Proc. Cambridge Philos. Soc.* 36 (1936), 477–481.
- [Pre74] C.J. Preston A generalization of the FKG inequalities. *Comm. Math. Phys.* 36(3) (1974), 233–241.
- [RS03] F. Ritort and P. Sollich. Glassy dynamics of kinetically constrained models. *Adv. Phys.* 52(4) (2003), 219–342.

- [SL95] A. Sudbury and P. Lloyd. Quantum operators in classical probability theory. II: The concept of duality in interacting particle systems. *Ann. Probab.* 23(4) (1995), 1816–1830.
- [SL97] A. Sudbury and P. Lloyd. Quantum operators in classical probability theory. IV: Quasi-duality and thinnings of interacting particle systems *Ann. Probab.* 25(1) (1997), 96–114.
- [SS08] A. Sturm and J.M. Swart. Voter models with heterozygosity selection. *Ann. Appl. Probab.* 18(1) (2008), 59–99.
- [SS15a] A. Sturm and J.M. Swart. A particle system with cooperative branching and coalescence. *Ann. Appl. Probab.* 25(3) (2015), 1616–1649.
- [SS15b] A. Sturm and J.M. Swart. Monotone and additive Markov process duality. *J. Theor. Probab.* 31(2) (2018), 932–983.
- [Sud97] A. Sudbury. The convergence of the biased annihilating branching process and the double-flipping process in \mathbb{Z}^d . *Stoch. Proc. Appl.* 68 (1997), 25–264.
- [Sud99] A. Sudbury. Hunting submartingales in the jumping voter model and the biased annihilating branching process. *Adv. Appl. Probab.* 31(3) (1999), 839–854.
- [Sud00] A. Sudbury. Dual families of interacting particle systems on graphs. *J. Theor. Probab.* 13(3) (2000), 695–716.
- [Swa13a] J.M. Swart. Noninvadability implies noncoexistence for a class of cancellative systems. *Electron. Commun. Probab.* 18 (2013), paper No. 38, 1–12.
- [Swa13b] J.M. Swart. Duality and Intertwining of Markov Chains. Lecture notes for a course at the ALEA in Europe School, October 21–25 (2013) at CIRM, Luminy (Marseille). Available at http://staff.utia.cas.cz/swart/posters_index.html.
- [Tov19] D. Tovazzi. *Self-sustained periodic behaviors in interacting systems: macroscopic limits and fluctuations*. PhD Thesis University of Padua, 2019.

Index

- $\| \cdot \|_1$, 10
- $\| \cdot \|_\infty$, 10, 67
- $|\cdot|$, 11, 124
- \rightsquigarrow , 83
- $\underline{0}, \underline{1}$, 112
- $B(E)$, 67
- $\mathcal{B}(E)$, 67
- $\mathcal{C}(E)$, 67
- \mathcal{C}_{fin} , 90
- \mathcal{C}_{sum} , 90
- $\mathcal{D}(m)$, 78
- \mathcal{E} , 10
- \mathcal{I} , 114
- \mathcal{I}_e , 114
- Kf , 68
- $\mathcal{M}_1(E)$, 67
- $M_{x,i}$, 100
- \mathcal{N}_i , 10
- $N_{x,i}$, 17
- P_tf , 68
- $\mathcal{R}(f)$, 78
- $\mathcal{S}(\Lambda), \mathcal{S}_{\text{fin}}(\Lambda)$, 129
- T_i , 157
- $\mathbf{X}_{s,t}$, 88
- $\mathbf{X}_{s,u}$, 82, 88
- $\mathbf{X}_{s,u}^\pm$, 130
- X_t^x , 158
- ann_{ij} , 25
- bra_{ij} , 13
- bran_{ij} , 133
- coop_{ijk} , 28
- death_i , 14
- death_{ij} , 119
- excl_{ij} , 26
- kill_{ij} , 28
- $m_*(\beta)$, 19
- $m[i]$, 78
- \mathbf{m}_t^ω , 41, 81
- rebel_{ijk} , 136
- rw_{ij} , 25
- vot_{ij} , 11
- $x_{\text{upp}}(\beta)$, 51
- λ_c , 162
- μK , 68
- μP_t , 68
- μf , 68
- $\omega_{s,u}$, 41
- ψ , 139
- ψ_{add} , 129
- ψ_{canc} , 134
- ρ , 158, 161
- $\theta(\lambda)$, 15, 115, 157
- $\xi_s^{(i,t)}$, 155
- additive
 - interacting particle system, 124
 - map, 124
- annihilating branching map, 133
- annihilating random walk map, 25
- antiferromagnetic, 18
- atom of a measure, 36
- atomless measure, 36
- autonomous Markov process, 56
- backward in time process, 123
- Bernoulli
 - random variable, 37

- biased annihil. branching process, 29
- biased voter model, 23
- bounded linear operator, 70
- branching map, 13
- cadlag, 69, 130
- cancellative
 - duality function, 134
 - interacting particle system, 132
 - map, 132
- Cauchy equation, 71
- closable linear operator, 70
- closed linear operator, 70
- closure of linear operator, 70
- cluster, 11
- clustering, 11, 19
- coalescing random walk map, 25
- conservative Feller semigroup, 76
- contact process, 13
 - mean-field, 59
- cooperative branching map, 28
- core of closed operator, 73
- counting measure, 36
- coupling, 107
- critical
 - infection rate, 15, 162
 - point, 15, 19, 20
- critical behavior, 23
- critical point, 162
- Curie-Weiss model, 48
- death map, 14
- death rate, 15
- deterministic
 - probability kernel, 38, 68
- directed percolation, 166
- distribution determining, 103
- Dobrushin, 5
- domain of a linear operator, 70
- double death map, 119
- drift function, 49
- dual
 - lattice, 167
- dual map, 139, 140
- duality function, 139, 140
- edge set, 10
- embedded Markov chain, 39
- ergodic interacting particle system, 98
- exclusion map, 26
- exclusion process, 27
- explosion
 - of Markov processes, 44
- explosion time, 46
- explosive
 - Markov process, 46
- extremal invariant law, 114
- f -relevant, 78
- Feller
 - semigroup, 69
- ferromagnetic, 18
- generator, 9
 - Feller semigroup, 71
- Glauber dynamics, 17
- graph of an operator, 70
- graphical construction, 82
- graphical representation, 82
- Hölder's inequality, 159
- homogeneous, 157
- incident, 82
- increasing set, 109
- independent increments
 - of a stochastic flow, 42
- infection rate, 14
- integer lattice, 10
- intensity
 - of a Poisson point measure, 36
 - of a random measure, 35

- interacting particle system, 9
- intertwining of Markov processes, 148
- invariant law, 96
- Ising model, 16
 - mean-field, 48
- k -dependence, 172
- killing map, 28
- Laplace equation, 72
- lattice, 9, 65, 77
- Liggett, 6
- local state space, 9
- local map, 77
- local probability kernel, 77
- local state, 9
- local state space, 65, 77
- locally finite measure, 35
- lower invariant law, 112
- lower semi-continuous function, 141
- magnetization
 - spontaneous, 19
- Markov semigroup, 38
- maximum principle
 - positive, 72
- metastable behavior, 53
- monotone
 - function, 107
 - interacting particle system, 110
 - probability kernel, 109
- monotone representability, 109, 110
- nearest neighbor, 10, 11
- nontrivial
 - law, 116, 158
- one-point compactification, 45
- open path
 - in graphical representation, 127
 - oriented percolation, 166
- oriented percolation, 166
- path
 - of potential influence, 83
 - open
 - oriented percolation, 166
- Peierls argument, 167
- percolation
 - directed, 166
 - oriented, 166
 - parameter, 166
- phase transition, 20
- piecewise constant, 39
- Poisson point measure, 36
- Poisson point set, 36
- positive maximum principle, 72, 75
- Potts model, 16
- probability kernel, 38, 67
- product topology, 66
- random mapping representation
 - of a generator, 40, 109
 - of a probability kernel, 38, 68
 - of probability kernel, 109
- range of a particle system, 11
- rate of jumps, 39
- rebellious map, 136
- relevant
 - coordinate for function, 78
- renormalization group theory, 22
- separation of points, 103
- simple counting measure, 36
- site, 9
- Skorohod topology, 69
- Spitzer, 5
- spontaneous magnetization, 19, 118
- state space, 9
- stationary process, 97
- stochastic flow, 42
- stochastic order, 108
- subprobability kernel, 76
- summable variation, 91

- supremumnorm, 67
- thinning, 37, 149
- threshold voter model, 137
- transition probability
 - continuous, 68
- translation
 - invariant, 157
 - operator, 157
- two-stage contact process, 16
- type of a site, 11
- upper invariant law, 112
- variation
 - of a function on S^Λ , 90
- voter model, 11
- voter model map, 11
- weak convergence, 67
- wet sites, 167
- Wright-Fisher diffusion, 62, 74